

Dealing with *Indigofera Nivea* (Leguminosae) — A New Name for Madagascar and a New Combination for Africa

Authors: Callmander, Martin W., Labat, Jean-Noël, and Schrire, Brian D.

Source: *Candollea*, 65(2) : 363-364

Published By: The Conservatory and Botanical Garden of the City of Geneva (CJBG)

URL: <https://doi.org/10.15553/c2010v652a12>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

I. CALLMANDER, Martin W., Jean-Noël LABAT & Brian D. SCHRIRE:

Dealing with *Indigofera nivea* (Leguminosae) – a new name for Madagascar and a new combination for Africa

Introduction

This first note deals with the name “*Indigofera nivea*” which comprises two homonyms (one based on material from Madagascar and the other on material from southern Africa). A new species name is required for the illegitimate later homonym published by VIGUIER (1949) for which no valid name is available. The name *I. viguieri* Callm. & Labat is published here for this Malagasy species. *Indigofera nivea* Spreng. was validly published by SPRENGEL in 1826, but has been generally overlooked or confused in the literature. It is an earlier name for the southern African plant that has been known as *Indigastrum argyraeum* (Eckl. & Zeyh.) Schrire (SCHRIRE, 1992). A new combination in the genus *Indigastrum* Jaub. & Spach is required, which is published here as *I. niveum* (Spreng.) Schrire & Callm. Another tentative use of the specific epithet *nivea* was as a misapplied synonym of *Indigofera glomerata* E. Mey. (MEYER, 1832). He placed a question mark in his synonymy indicating his doubt about the identification of the Sprengel name.

Taxonomy and nomenclature of *Indigofera nivea* Spreng.

Indigofera viguieri Callm. & Labat, **nom. nov.**

≡ *Indigofera nivea* R. Vig. in Notul. Syst. (Paris) 13: 368. 1949 [non Spreng., Syst. Veg. 3: 273. 1826].

Typus: MADAGASCAR. **Prov. Mahajunga:** Ambongo, rive droite de la Mahavavy, Itampika, 13°38'23"S 48°40'06"E, V.1902, fl., *Perrier de la Bâthie 1434* (holo-: P [P0007766]!; iso-: K [K000392972]!, P [P0007767, P0007768]!).

Observations. – *Indigofera nivea* R. Vig., published in VIGUIER (1949), is endemic to Madagascar and was accepted as a valid name in “Leguminosae of Madagascar” by DU PUY & al. (2002). Viguier’s name is in fact a later homonym of *I. nivea* Spreng., and no valid name is available for the Malagasy species. The species is a small subshrub with densely white-woolly leaves and stems. *Indigofera viguieri* can be distinguished from the most closely related species, *I. kirkii* Oliv., by its densely white-woolly leaves (vs. the thinly white strigose leaves in *I. kirkii*), the more coriaceous texture of the leaves, and pods with 3-5 (vs. 1-3) seeds (DU PUY & al., 2002). It is known from only two collections made by Perrier de la Bâthie in northwestern Madagascar on sand dunes bordering the Betsiboka and Mahavavy rivers.

Etymology. – We name this species after René Viguier (1880-1931), from Caen (France), who made a major contribution to the knowledge of *Leguminosae* in Madagascar. René Viguier undertook a full revision of the family that he left unpublished when he died in 1931. Most of Viguier’s work was validated posthumously thanks to H. Humbert (VIGUIER, 1949, 1950, 1952) and his unpublished manuscript, “les Légumineuses de Madagascar” served as the basis for the “Leguminosae of Madagascar” (DU PUY & al., 2002).

Conservation status. – With an EOO of 18 km² and only two collections known, both dating back over one hundred years, and these comprising only two subpopulations, none of which occur in a protected network, *I. viguieri* is best assigned a preliminary status of “Critically Endangered” (CR A3c; B2ab[iii]) following the IUCN Red List Categories and Criteria (IUCN, 2001) (calculation following CALLMANDER & al., 2007).

Addresses of the authors: MWC: Missouri Botanical Garden, P.O. Box 299, St. Louis, Missouri, 63166-0299, U.S.A. and Conservatoire et Jardin botaniques de la Ville de Genève, ch. de l'Impératrice 1, C. P. 60, 1292 Chambésy, Switzerland. Email: martin.callmander@mobot.org

JNL: Muséum national d'Histoire Naturelle, Département Systématique et Evolution, UMR 7205, case postale 39, rue Cuvier 57, 75231 Paris, cedex 05, France.

BDS: Herbarium, Library, Art & Archives, Royal Botanic Gardens Kew, Richmond, Surrey, TW9 3AB, United Kingdom.

Indigastrum niveum (Spreng.) Schrire & Callm., **comb. nova**

= *Indigofera nivea* Spreng., *Syst. Veg.* 3: 273. 1826.

Typus: SOUTH AFRICA. **W. Cape Prov.:** *Willdenow 13890* (holo-: B-W!).

= *Indigofera argyraea* Eckl. & Zeyh., *Enum. Pl.*: 239. 1836. = *Indigastrum argyraeum* (Eckl. & Zeyh.) Schrire in *Bothalia* 22: 168. 1992. **Typus:** SOUTH AFRICA. **E. Cape Prov.:** Tembuland, nr. Klipplaat R. & Swart Kei R., by Shiloh Mission, 1060-1200 m, I-III.1832, *Ecklon & Zeyher 1595* (lecto-: S!; isolecto-: B! G! M! SAM! TCD!) (lectotypified by SCHRIRE, 1992).

Observations. – Taxonomic research by SCHRIRE (1991, 1992, 1995) and molecular phylogenies (SCHRIRE & al., 2003, 2009), provided the background that justified the resurrection of the genera *Microcharis* Benth. and *Indigastrum*, which had generally been placed in synonymy under *Indigofera* L. (SCHRIRE, 1992). Among others, the taxon *Indigofera argyraea* was therefore transferred to the genus *Indigastrum*. In “Plants of southern Africa: names and distribution” (ARNOLD & DE WET, 1993), Schrire placed *Indigofera nivea* in synonymy under *Indigastrum argyraea*, based on unpublished information before he was able to study the type. More recently after studying the type in the Willdenow collection in Berlin, the identity of the Sprengel name was confirmed, and it became apparent that *Indigofera argyraea* and *I. nivea* are indeed conspecific, the necessary new combination being made here.

Acknowledgements

The first author thanks Cyrille Chatelain and Pierre-André Loizeau for support at the Conservatoire et Jardin botaniques de la Ville de Genève (CJBG). The authors are also grateful to Patrick Perret for his assistance with nomenclatural matters at the CJBG. Financial support was provided by grants from the U.S. National Science Foundation (0743355) and the Andrew W. Mellon Foundation.

References

- ARNOLD, T. H. & B. C. DE WET (1993). Plants of southern Africa: names and distribution. *Mem. Bot. Survey S. Africa* 62.
- CALLMANDER, M. W., G. E. SCHATZ, P. P. LOWRY II, M. O. LAIVAO, J. RAHARIMAMPIONONA, S. ANDRIAMBOLOLONERA, T. RAMINOSOA & T. CONSIGLIO (2007). Application of IUCN Red List criteria and assessment of Priority Areas for Plant Conservation in Madagascar: rare and threatened Pandanaceae indicate new sites in need of protection. *Oryx* 41: 168-176.
- DU PUY, D. J., J.-N. LABAT, R. RABEVOHITRA, J. F. VILLIERS, J. BOSSER & J. MOAT (2002). *The Leguminosae of Madagascar*. Royal Botanic Gardens, Kew.
- IUCN (2001). *IUCN Red List Categories and Criteria: version 3.1*. IUCN Species Survival Commission. IUCN, Gland and Cambridge.
- MEYER, E. H. F. (1832). *Plantae Ecklonianae*. *Linnaea* 7: 145-173.
- SCHRIRE, B. D. (1991). *Systematic studies in African Indigoferaeae (Leguminosae-Papilionoideae)*. Ph.D. thesis, University of Natal, South Africa.
- SCHRIRE, B. D. (1992). New combinations and resurrected names in *Microcharis* and *Indigastrum* (Leguminosae-Papilionoideae). *Bothalia* 22: 165-170.
- SCHRIRE, B. D. (1995). Cladistic analysis of the tribe Indigoferaeae (Leguminosae). In: CRISP, M. D. & J. J. DOYLE (ed.), *Advances in Legume Systematics, Part 7, Phylogeny*: 161-244. Royal Botanic Gardens, Kew.
- SCHRIRE, B. D., M. LAVIN, N. P. BARKER, H. CORTES-BURNS, I. VON SENGER & J.-H. KIM (2003). Towards a phylogeny of *Indigofera* (Leguminosae-Papilionoideae): identification of major clades and relative ages. In: KLITGAARD, B. B. & A. BRUNEAU (ed.), *Advances in Legume Systematics, part 10, Higher level systematics*: 269-302. Royal Botanic Gardens, Kew.
- SCHRIRE, B. D., M. LAVIN, N. P. BARKER & F. FOREST (2009). Phylogeny of the tribe Indigoferaeae (Leguminosae-Papilionoideae): Geographically structured more in succulent-rich and temperate settings than in grass-rich environments. *Amer. J. Bot.* 96: 816-852.
- SPRENGEL, C. (1826). *Systema vegetabilium*. Vol. 3. Göttingen.
- VIGUIER, R. (1949). *Leguminosae Madagascarienses Novae*. *Notul. Syst. (Paris)* 13: 333-369.
- VIGUIER, R. (1950). *Leguminosae Madagascarienses Novae* (suite 1). *Notul. Syst. (Paris)* 14: 62-74.
- VIGUIER, R. (1952). *Leguminosae Madagascarienses Novae* (suite 2). *Notul. Syst. (Paris)* 14: 168-187.