



An overlooked name provides the basionym for a new combination in Malagasy Tarenna (Rubiaceae)

Author: Turner, Ian M.

Source: Candollea, 77(2) : 237-240

Published By: The Conservatory and Botanical Garden of the City of Geneva (CJBG)

URL: <https://doi.org/10.15553/c2022v772a8>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

An overlooked name provides the basionym for a new combination in Malagasy *Tarenna* (Rubiaceae)

Ian M. Turner

Abstract

TURNER, I.M. (2022). An overlooked name provides the basionym for a new combination in Malagasy *Tarenna* (Rubiaceae). *Candollea* 77: 237–240. In English, English and French abstracts. DOI: <http://dx.doi.org/10.15553/c2022v772a8>

Pavetta madagascariensis Ten. (Rubiaceae) is an overlooked name described from material grown in the Naples Botanical Garden. The name is here transferred to *Tarenna* Gaertn. as *T. madagascariensis* (Ten.) I.M. Turner, and it becomes the correct name for the species currently referred to as *T. grevei* (Drake) Homolle. Neotypes are designated for *Pavetta madagascariensis* and *P. gracilis* A. Rich. ex DC. respectively and a lectotype for *Ixora grevei* Drake.

Résumé

TURNER, I.M. (2022). Un nom oublié fournit le basionyme d'une nouvelle combinaison de *Tarenna* (Rubiaceae) à Madagascar. *Candollea* 77: 237–240. En anglais, résumés anglais et français. DOI: <http://dx.doi.org/10.15553/c2022v772a8>

Pavetta madagascariensis Ten. (Rubiaceae) est un nom oublié décrit à partir de matériel cultivé dans le jardin botanique de Naples. Le nom est ici transféré à *Tarenna* Gaertn. sous le nom de *T. madagascariensis* (Ten.) I.M. Turner, et il devient le nom correct pour l'espèce actuellement désignée sous le nom de *T. grevei* (Drake) Homolle. Des néotypes pour *Pavetta madagascariensis* et *P. gracilis* A. Rich. ex DC. sont désignés respectivement et un lectotype pour *Ixora grevei* Drake.

Keywords

RUBIACEAE – *Pavetta* – *Tarenna* – Madagascar – Naples – Giovanni Gussone – Michele Tenore – neotype

Address of the author:

Royal Botanic Gardens Kew, Richmond, Surrey TW9 3AB, UK and Singapore Botanic Gardens, National Parks Board, 1 Cluny Road, Singapore 259569, Singapore. E-mail: i.turner@kew.org

First published online on November 23, 2022.

ISSN: 0373-2967 – Online ISSN: 2235-3658 – *Candollea* 77(2): 237–240 (2022)

© CONSERVATOIRE ET JARDIN BOTANIQUES DE GENÈVE 2022

Introduction

Nomenclatural databases are rarely perfect. The vagaries of fate can result in omissions from the best-organised compilations. So it is with the Index Kewensis and its successor, the International Plant Names Index, and an example of such an omission is discussed in this note. It concerns the validly published name: *Pavetta madagascariensis* Ten. (TENORE, 1845). Despite RIPPA (1935) drawing attention to the absence of this name from Index Kewensis, it is still missing. RIPPA (1935) noted that the species was described by Michele Tenore (1780–1861) from plants growing in the Naples Botanical Garden. These were cultivated from material introduced from Madagascar by a certain “Dr. Hardy” in 1839. According to RIPPA (1935), material of this plant remained in cultivation in Naples for many years. Rippa identified the species as *P. gracilis* A. Rich. ex DC., which is an older name than *P. madagascariensis* Ten., and was evidently considered to be conspecific. However, since Rippa’s time, generic concepts have changed and this species is now considered to belong in *Tarenna* Gaertn. (VERDCOURT, 1983; DE BLOCK, 2005). The direct transfer of *Pavetta gracilis* to *Tarenna* is blocked by the name of a West African species, *T. gracilis* (Stapf) Keay. Verdcourt therefore published a nomen novum, *T. richardii* Verdc., for *Pavetta gracilis*. DE BLOCK (2005) noted that there were several heterotypic synonyms available that could provide names in *Tarenna* that had priority over *T. richardii*. These included *T. grevei* (Drake) Homolle, that was considered the correct name for the species by De Block. *Ixora grevei* Drake, the basionym of *Tarenna grevei*, was published in 1897, while Tenore’s *Pavetta madagascariensis* was published more than 50 years earlier and the name is not blocked in *Tarenna*.

The new combination *Tarenna madagascariensis* (Ten.) I.M. Turner is therefore proposed below.

Nomenclature

Tarenna madagascariensis (Ten.) I.M. Turner, **comb. nov.**

= *Pavetta madagascariensis* Ten., Cat. Orto Bot. Napoli: 90. 1845.

Neotypus (designated here): [MADAGASCAR]: cultivated in the Naples Botanical Garden, 14.VII.1850, Gussone s.n. (NAP [NAP0000515] image!).

= *Pavetta gracilis* A. Rich. ex DC., Prodr. 4: 492. 1830.
 = *Ixora gracilis* (A. Rich. ex DC.) Kuntze, Revis. Gen. Pl. 1: 287. 1891. = *Enterospermum gracile* (A. Rich. ex DC.) Bremek. in Repert. Spec. Nov. Regni Veg. 37: 193. 1934. = *Tarenna richardii* Verdc. in Kew Bull. 37: 561. 1983, nom nov. [non *T. gracilis* (Stapf) Keay, 1957], **syn. nov. Neotypus** (designated here): MADAGASCAR: “des côtes orient. de Madag.”, s.d., Richard 906 (P [P03919043] image!).

= *Ixora grevei* Drake in Bull. Mens. Soc. Linn. Paris 2: 1308. 1897. = *Tarenna grevei* (Drake) Homolle in Bull. Soc. Bot. France 85: 606. 1939, **syn. nov. Lectotypus** (designated here): MADAGASCAR: sine loco, s.d., Grevé 261 (P [P00173282] image!; isolecto-: P [P00115297, P00173281, P00173283, P03919323] images!, TAN [TAN000755] image!).

Notes. – There is no specimen of *Pavetta madagascariensis* in Tenore’s herbarium in Naples (R. Vallariello, pers. comm.), but, there is a specimen (Fig. 1) in the herbarium of Giovanni Gussone (1787–1811), a Neapolitan botanist contemporary of Tenore. The specimen accords well with Tenore’s description, with one notable exception. In the protologue, Tenore stated ‘calycibus 5-dentatis’ but the Gussone specimen has tetramerous flowers. As the specimen is a good match to *Tarenna grevei* and it is labelled with Tenore’s name in Gussone’s hand, I consider the discrepancy in the number of calyx lobes to be a lapse on Tenore’s part and accept Rippa’s interpretation of the name. The Gussone specimen [NAP0000515] is designated here as the neotype of *Pavetta madagascariensis*, and a new combination for this name is proposed in *Tarenna*.

Pavetta gracilis was described separately by Augustin Pyramus de Candolle (1778–1841) and by Achille Richard (1794–1852) in 1830 when they were both studying the family Rubiaceae. Candolle’s fourth volume of the *Prodromus* was published two months earlier than Richard’s *Mémoire* (see KIRKBRIDE & WIERSEMA, 2022). Among the 72 Rubiaceae names ascribed to Richard in the *Prodromus*, several were not associated to any specimen because Candolle copied Richard’s descriptions and no original material was available to him (KIRKBRIDE & WIERSEMA, 2022). A neotype is therefore designated here on the best material seen by Richard now deposited at P [P03919042].

The best preserved material of Grevé 261 is designated here as the lectotype of *Ixora grevei*.

Acknowledgements

I am very grateful to Annalisa Santangelo and Roberta Vallariello of Università degli Studi di Napoli Federico II for their kind assistance in searching for specimens and providing images and to Petra De Block for sharing images of Grevé 261 and advice on the designation of the lectotype.




Fig. 1. – Neotype of *Pavetta madagascariensis* Ten. in the herbarium of Giovanni Gussone. [NAP0000515; © Università degli Studi di Napoli Federico II]

References

- DE BLOCK, P. (2005). Notes on *Tarenna* (Pavetteae) as a precursor to a checklist of Rubiaceae of Madagascar and the Comores. *Syst. Geogr. Pl.* 75: 107–116.
- KIRKBRIDE, J.H. JR. & J.H. WIERSEMA (2022). Authorship and typification of Rubiaceae names published by A.P. de Candolle and A. Richard revisited. *Taxon* [online version]: <https://doi.org/10.1002/tax.12724>
- RIPPA, A. (1935). Notizie sulla Pavetta madagascariensis Ten. *Bull. Orto Bot. Regia Univ. Napoli* 13: 7–9.
- TENORE, M. (1845). *Catalogo delle piante che si coltivano nel R. Orto Botanico di Napoli: corredato della pianta del medesimo, e di annotazioni*. Tipografia dell' Aquila di V. Puzziello, Napoli.
- VERDCOURT, B. (1983). Notes on Mascarene Rubiaceae. *Kew Bull.* 37: 521–574.