

Review of the Dauer Hypothesis: What Non-Parasitic Species Can Tell us about the Evolution of Parasitism

Authors: Bubrig, Louis T., and Fierst, Janna L.

Source: Journal of Parasitology, 107(5) : 717-725

Published By: American Society of Parasitologists


URL: <https://doi.org/10.1645/21-40>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



REVIEW OF THE DAUER HYPOTHESIS: WHAT NON-PARASITIC SPECIES CAN TELL US ABOUT THE EVOLUTION OF PARASITISM

Louis T. Bubrig¹ and Janna L. Fierst²

¹ Department of Biology, University of Virginia, 485 McCormick Road, Charlottesville, Virginia 22904.

² Department of Biological Sciences, University of Alabama, 300 Hackberry Lane, Tuscaloosa, Alabama 35487-0344. Correspondence should be sent to Louis T. Bubrig (<http://orcid.org/0000-0002-2578-8397>) at: ltb4pv@virginia.edu

KEY WORDS ABSTRACT

Evolution	Parasitic lineages have acquired suites of new traits compared to their nearest free-living relatives.
Nematodes	When and why did these traits arise? We can envision lineages evolving through multiple stable intermediate steps such as a series of increasingly exploitative species interactions. This view allows us to use non-parasitic species that approximate those intermediate steps to uncover the timing and original function of parasitic traits, knowledge critical to understanding the evolution of parasitism.
Ecology	The dauer hypothesis proposes that free-living nematode lineages evolved into parasites through two intermediate steps, phoresy and necromeny. Here we delve into the proposed steps of the dauer hypothesis by collecting and organizing data from genetic, behavioral, and ecological studies in a range of nematode species. We argue that hypotheses on the evolution of parasites will be strengthened by complementing comparative genomic studies with ecological studies on non-parasites that approximate intermediate steps.
Behavior	
Species interactions	
Dauer	
Parasite	
Phoresy	
Necromeny	

Parasitism has evolved hundreds of times (Geary and Thompson, 2001; Johnson et al., 2004; Weinstein and Kuris, 2016). The most intimate types of parasitism in which the parasite lives on or in the host are mediated by gene family expansions (Tsai et al., 2013), horizontal gene transfers (Davis et al., 2000), new behaviors (Poinar, 1983; Chaisson and Hallem, 2012), and novel molecular defenses (Warburton and Zelmer, 2010; Haegeman et al., 2011). Within the large number of changes in highly derived parasite lineages is a subset of core traits that mediate the fundamental components of the parasitic lifestyle. Because these components are numerous (e.g., host finding, host attachment, immune system evasion), the parasitic lifestyle seems prohibitively irreducible (Dieterich and Sommer, 2009). If an entire suite of traits is required for successful parasitism, how does any individual trait increase in frequency in a free-living population?

One hypothesis is that many “parasitism” traits arose well before the transition to parasitism and were co-opted from their ancestral functions. Specifically, lineages acquired parasitic traits by establishing and shifting between other types of species interactions like mutualisms or commensalisms (Poinar, 1983; West et al., 1996; Johnson et al., 1997; Weinstein and Kuris, 2016). Phylogenies support this hypothesis: parasitic lineages’ closest non-parasitic relatives often associate with similar host species in non-parasitic ways (Proctor and Owens, 2000; Kikuchi et al., 2011; Weinstein and Kuris, 2016).

This hypothesis suggests that studying parasitic lineages alone could overlook important aspects of their evolutionary history.

When inferring when a given trait arose, we can be led astray by assuming its ancestral function was the same as its current one. For instance, plant-parasitic nematodes have acquired cellulase enzymes horizontally (Haegeman et al., 2011). These enzymes enable nematodes to burrow through host plant tissue, suggesting that their acquisition occurred concurrently with the transition to parasitism (Jones, 1981). Alternatively, the enzymes could have been acquired much earlier for a different purpose before being co-opted for parasitism. To investigate this possibility, we can see if any non-parasitic species have cellulase genes that serve non-parasitic functions. The free-living nematode *Pristionchus pacificus* has 7 cellulase genes (Mayer et al., 2011), challenging the assumption that these are exclusive to plant parasites and suggesting that cellulases may have been acquired well before the transition to parasitism—possibly to break down bacterial biofilms (Romeo, 2008), as in *P. pacificus*—and later co-opted for plant tissue invasion. In short, non-parasites demonstrate alternative uses for so-called “parasitism traits,” which helps us investigate their origins.

An ideal system for studying the evolution of parasitism should therefore be rich in interaction types with readily available ecological, behavioral, genetic, and developmental data. Nematodes are popular for these reasons, but genetic and developmental data in this group (and many other parasitism study systems) have outpaced basic understanding of behavior and ecology (West et al., 1996; Whitfield, 1998; Proctor and Owens, 2000; Félix and Braendle, 2010; Adams et al., 2020). Comparative

genomic studies have attempted to unlock the evolution of parasitism—for example, by studying convergence in molecular pathways (Crook, 2014)—but without an ecological understanding of species interactions, these models of parasitism evolution are limited. Here we argue that studying parasitism requires collecting seemingly disparate ecological data and reexamining them through the lens of parasitism evolution. We focus on 1 model, the so-called dauer hypothesis in nematodes, to update our knowledge of the hypothesis and illustrate how an ecological approach can benefit any parasitic study system.

THE DAUER HYPOTHESIS

The dauer hypothesis developed from the observation that free-living and parasitic nematodes have remarkably conserved life cycles (Fig. 1). All nematodes proceed through 4 larval stages separated by molts. Some species have an extra component to the life cycle; instead of passing through one of the normal larval stages, they may enter an alternative life stage with unique properties and functions (Rogers and Sommerville, 1963). In some species, the alternative stage (called the “dauer”) facilitates dormancy, a halting of development until conditions are favorable (Poinar, 1983). In others, the alternative stage (also called the “dauer”) is used for dispersing between habitat patches via larger carrier animals (Kiontke and Sudhaus, 2006; Félix and Braendle, 2010). In parasitic species, the alternative stage (called the “infective juvenile”) facilitates transmission between hosts (Anderson, 1984). Though their functions differ, these alternative stages survive harsh external stressors (Ellenby, 1968; Klass and Hirsh, 1976; Stasiuk et al., 2012), tie development to external cue detection (Rogers and Sommerville, 1963; Cassada and Russell, 1975; Stasiuk et al., 2012), have similar morphological structure and neuronal regulation, and almost always split from the main life cycle at the L2–L3 molt (Crook, 2014). The dauer hypothesis suggests that these 3 alternative stages are homologous (Rogers and Sommerville, 1963) and that the infective juveniles of parasites are derived from the more ancestral dauer larva (Hotze et al., 1993; Crook, 2014).

This broad version of the dauer hypothesis applies to any of the 15 or more origins of parasitism in Nematoda, including transitions to both plant and animal parasitism (Blaxter and Koutsovoulos, 2015). Over the years, researchers have detailed more specific versions for the evolution of different types of parasites. The best-known version lays out a 4-step evolutionary pathway for transitions to animal parasitism. The proposed evolutionary sequence is 1) free-living ancestors that do not associate with any larger species, 2) phoretic relationships in which nematodes superficially attach to a larger animal for dispersal, 3) necromeny, in which nematodes may feed on their dead hosts without directly contributing to the death themselves, and 4) parasitism (Crook, 2014). Many extant nematode species approximate these 4 steps (Poinar, 1983), so studying them can reveal the origins of parasitism traits, their ancestral benefits, and when they were co-opted for new purposes.

STEP ONE: NON-ASSOCIATING

The dauer hypothesis proposed that the road to parasitism begins early, and the ecological conditions at the non-associating




Figure 1. Newly hatched nematodes assess their environment and then develop as an L3 (normal larva) or a dauer (resistant larva). L1–L4: larval stage 1–4. Color version available online.

step profoundly impact whether and how lineages evolve towards parasitism.

Free-living nematodes live in a variety of habitats, such as marine sediments or terrestrial soil, and eat different diets, like bacteria, fungus, yeast, or micro-animals (Poinar, 1983). This variation imposes unique stressors and selects different traits that can influence later developments. For instance, desiccation is a major selective force in terrestrial but not marine environments (Rebecchi et al., 2020). The dauer larva is thought to combat this stressor because it retains water well (Rogers and Sommerville, 1963; Ellenby, 1968; McSorley, 2003), remains dormant for a long time (Klass and Hirsh, 1976), and exists only in terrestrial lineages (Anderson, 1988, 1996). If the dauer precedes parasitism as suggested by the dauer hypothesis, we might expect that parasitism has only evolved in terrestrial lineages. Consistent with this prediction, phylogenetic analysis shows that parasitic lineages overwhelmingly derive from terrestrial ancestors; even nematodes that parasitize marine hosts like fish are more closely related to terrestrial nematode lineages than to marine ones (Anderson, 1984, 1996; Poulin, 2007), despite marine lineages being far older (Anderson, 1996; Blaxter and Koutsovoulos, 2015). In short, lineages have unequal probabilities of evolving parasitism depending on their ecological context.

Traits in the non-associating step may also predict what kind of parasitism evolves. Animal parasites are most closely related to bacteria-eating nematodes with narrow pharynxes (Poinar, 1983; Campbell and Lewis, 2002), whereas plant parasites derive from fungivorous nematodes with long hollow stylets (Poinar, 1983; Luc et al., 1987) (but see Holterman et al., 2009). The traits that facilitate each diet are more easily co-opted to parasitize one type of host over the other. Again, even the most basic early traits like diet can have far-reaching impacts on future developments.

STEP TWO: PHORESY

Under the dauer hypothesis, the next step in parasitism evolution is phoresy, a transport relationship between a small carried species, the phoront, and a more mobile species, the vector (Crook, 2014; White et al., 2017). Phoresy is advantageous for exploiting patchy resources as opposed to more continuous




Figure 2. Phoretic nematodes can improve their dispersal success at multiple steps. Color version available online.

resources. For example, many lineages proliferate on dense blooms of microbes found on substrates like rotting fruit or carrion (Félix and Braendle, 2010). Such resources are far apart and short-lived, which selects for fast life cycles and phoresy (Young, 1961; Southwood, 1977; Bartlow and Agosta, 2021).

Phoresy as a step toward parasitism is not limited to the dauer hypothesis; it shows up in taxa such as mites (Houck and OConnor, 1991; Athias-Binche and Morand, 1993). Two aspects of phoresy contribute to its potential significance in the evolution of parasitism. Unlike species interactions that involve passing contact between the 2 species, phoresy necessitates intimate and long-lasting physical contact between the phoront and its vector, which may lend itself to future exploitation. Phoresy also constrains the sizes of the participants; it only works if the phoront is much smaller than the vector. This same size difference characterizes host-parasite interactions (Lafferty and Kuris, 2002), so phoresy may be more likely to foster parasitism than interactions that do not constrain participant size.

We might predict that lineages of adept dispersers (or those with no need to disperse) would have no need for phoresy and therefore not acquire the vector-finding and vector-attachment traits required by parasitism. We find evidence supporting this prediction in the difference between marine vs. terrestrial lineages. Marine nematodes are poor directional swimmers (Sherman et al., 1983) but can easily ride currents to new locations (Heip et al., 1985) or attach to floating driftwood and algae (Gerlach, 1977; Highsmith, 1985). When marine sediments are disturbed in experimental plots, nematodes swiftly recolonize them via these modes of dispersal (Sherman and Coull, 1980; Chandler and Fleeger, 1983; Sherman et al., 1983). Marine nematodes also have less need to disperse since their habitats are continually replenished with organic material (Heip et al., 1985; Allredge and Silver, 1988). Therefore, marine nematodes would probably receive little benefit from phoretic relationships, which explains their rarity (Poulin, 2007). Even outside of Nematoda, marine phoronts are rarer than terrestrial phoronts, though under-sampling cannot be ruled out (Bartlow and Agosta, 2021). In terrestrial lineages, ephemeral habitats necessitate frequent dispersal via a vector (OConnor, 1982; Félix and Braendle, 2010); artificial habitats are rarely colonized by nematodes when vectors are excluded (M. Rockman, pers. comm.). Therefore, the

spatial structure of terrestrial habitats can select for traits that increase dispersal ability through phoresy (Houck and OConnor, 1991).

Phoresy adaptations

Phoresy selects for traits that promote dispersal by increasing the frequency and success of transport; these same traits may subsequently promote host finding in parasitism. Phoronts can increase dispersal success in 4 ways: preparing for dispersal in the starting habitat, embarking on the vector, surviving during travel, and disembarking into the new habitat (Fig. 2).

How does a phoretic organism know when to stay in the starting habitat and when to disperse? Individuals that disperse too early miss out on resources and reproductive opportunities still available in the starting habitat. Individuals that fail to notice a swiftly degrading habitat may miss their chance to leave (Viney and Harvey, 2017). The most obvious solution is to tie dispersal to environmental cues. The dauer stage of non-associating nematodes is induced by poor environmental conditions (Cassada and Russell, 1975), so lineages that used the dauer stage for dispersal could immediately co-opt that sensory machinery to temporarily limit dispersal attempts. Subsequently, this machinery could be modified in parasites to find their hosts.

Several traits make embarkment on the vector more predictable. Phoronts that sense and move toward vectors increase their chance of being picked up relative to phoronts that encounter vectors randomly (Haas, 2003). Potential chemotaxis cues include vector pheromones, metabolites, or temperature (Rogers and Sommerville, 1963; Granzer and Haas, 1991). These cues may be general—produced by a wide range of vector species—or specific to particular taxa. For instance, many phoretic and parasitic nematodes are attracted to carbon dioxide, a general cue given off by exhaling animals (Sciaccia et al., 2002; Haas, 2003; Harbison et al., 2009). In some cases, vector seeking is temporally limited because only the dauer stage is attracted to carbon dioxide (Hallem et al., 2011). Some phoretic species even sense specific cues like pheromones or excreted compounds in vector waste (Haas, 2003; Hong and Sommer, 2006). More chemosensation and chemotaxis studies should be performed to understand the relationship between chemosensation and the range of vectors a given nematode uses in the wild.

Phoronts also use structures or behaviors to maximize the chance of vector attachment (White et al., 2017). In parasitic species, this translates to an increasing chance of infection after exposure to the host. Heritable intraspecific variation in such traits could allow selection to increase the likelihood of attachment and solidify the relationship between the phoront and vector (Durkin and Luong, 2018; Durkin et al., 2020). Unlike some taxa that require complex structures like hooks or claws (Bartlow and Agosta, 2021), most nematodes only need a film of water to stick to passing organisms and inanimate objects (Stiernagle, 2006). To ensure contact, dauers climb fungal stalks or other surface irregularities, hang on by their tails, and wave (Reed and Wallace, 1965; Croll and Matthews, 1977); this behavior (“nictation”) is hypothesized to reduce the dauer’s contact with its substrate and better attach to its vectors (Campbell and Gaugler, 1993). Lab studies show that in both non-parasitic and parasitic species, individuals that nictate attach to vectors at significantly higher rates than non-nictating species or nictation-defective mutants of nictating species (Granzer and Haas, 1991; Campbell and Gaugler, 1993; Brown et al., 2011; Lee et al., 2012). Some parasitic species nictate faster when detecting carbon dioxide, presumably to increase the chance of embarkment when a host is nearby (Rogers and Sommerville, 1963; Granzer and Haas, 1991; Gang and Hallem, 2016). The nictation behavior itself may have been co-opted from the non-associating step where a stand-and-wave technique would be useful in traversing wide pores in the soil matrix (Reed and Wallace, 1965; Griffin, 2012). Such behaviors are physically demanding. Normally, dauers that sit and wait for better environments retain their old cuticle to provide a double layer of desiccation resistance (Ellenby, 1968; Poinar, 1983). Phoretic species cast off their old cuticle immediately after the molt (Cassada and Russell, 1975; Kiontke and Sudhaus, 2006), possibly to facilitate the acrobatics of host seeking, climbing, and nictation.

Once on the vector, the phoront must endure several environmental stressors, notably desiccation, starvation, and time. These same stressors affect parasitic species when they are searching for new hosts. The dauer stage lives off stored nutrients for months (Klass and Hirsh, 1976; Wadsworth and Riddle, 1989) so they are not as food limited as other life stages would be. To limit desiccation, dauers seek out humid regions near the vector’s core or around openings: on beetles, they line up under the elytra (Blaxter, 2003; Moser et al., 2005), and on isopods, in gaps between segments (Kiontke and Fitch, 2013). Though the association is physically intimate, phoronts are not thought to interfere with their vector’s mobility.

When the vector reaches a new habitat, the phoront must disembark to resume development, feed, and reproduce. Disembarkment is less relevant to parasitism from a trait perspective and more relevant as an evolutionary constraint. Because phoronts resume feeding and reproduction only in new food patches, mutations that cause phoronts to avoid disembarking are quickly weeded out (Farish and Axtell, 1971; Houck and O’Connor, 1991). The fitness of the phoront is aligned with the fitness of the vector during transit, stabilizing the species interaction and blocking exploitation as long as the phoront still needs to arrive at a new habitat for food and reproduction. If, however, the phoront finds a way to break the coupling between phoront and vector fitness, exploitation can occur.

STEP THREE: NECROMENY

Necromenic nematodes use a vector to disperse, as in phoresy, but have an added feeding component: if the vector dies in transit, they may feed and develop on the microbes that proliferate on the cadaver (Crook, 2014). Necromeny thus builds on phoresy by selecting for traits that reinterpret the vector as a potential habitat by detecting vector health and tying development to these signals (Wirth, 2009). Parasitism can further evolve from this host detection and development. Additionally, necromeny breaks the coupling between an organism’s fitness and its vector’s fitness since the organism can survive to reproduce whether the vector lives or dies.

The transition from phoresy to necromeny can be facilitated or constrained by vector traits as well as phoront traits. While necromeny provides nematodes with an alternative food source and reproductive ground, it does not eliminate the need for further dispersal via another vector. In some vector species, death pheromones encourage conspecifics to aggregate around the cadaver or physically manipulate it, which can transfer necromenic organisms to a live vector, continuing the cycle (Wilson et al., 1958; Wirth, 2009; Yao et al., 2009). In contrast, some vector species are strongly repelled by conspecific death pheromones, which could bar the development of necromeny since the vector habitat is effectively a dead end (Yao et al., 2009). Alternatively, avoidance of dead conspecifics could select for more generalist nematodes that rely on pick up by other vector species.

Necromeny adaptations

Though necromeny has been noted for decades (Schulte, 1989), we know surprisingly little about it. Identifying necromenic species is challenging—many species appear necromenic in the lab environment but not in the wild (Blaxter, 2003; Félix et al., 2018). The context-specific nature of necromeny implies that the relevant changes are likely in signaling and communication rather than the capability of resource use. By this argument, necromeny selects for traits that reinterpret vector environments as sites of food and reproduction (Wirth, 2009). We focus on 4 major areas that could facilitate this shift (Fig. 3): detecting vector cues, detecting the vector microbiome, intraspecific communication between nematodes, and sensing the abiotic context.

Nematode-vector communication could contribute to necromenic behavior and prime the nematode for parasitism. In particular, nematodes may develop the ability to monitor vector health by detecting cues produced by sick or dead vectors. For instance, *Caenorhabditis elegans* (a suspected necromenic species) emerges from the dauer stage faster when detecting oleic acid (Kaul et al., 2014), a compound released by many arthropods after death (Wilson et al., 1958; Yao et al., 2009). Some cues are produced constitutively by healthy vectors, and their absence signals sickness or death. *Pristionchus* spp. are kept in the dauer stage by a constitutively produced beetle pheromone and resume development only once their vector dies and ceases pheromone production (Cinkornpumin et al., 2014). In theory, necromeny links nematode development to its vector. The linkage can strengthen over time, as demonstrated by the many parasitic species that cannot develop properly until sensing one or more host cues (Rogers and Sommerville, 1963).

Vectors are also defined by a surface microbiome during life and the necrobiome that proliferates after death (Schulte, 1989).




Figure 3. In deciding whether to resume development on a dead host, food signals may be integrated with environmental signals, host signals, and intraspecific communication. Modifications in sensitivity to these signals or changes in integration may be necessary for the development of necromeny. Color version available online.

The vector microbiome likely differs from the microbiome on the phoront's patchy habitat. Nematodes distinguish between bacteria with impressive resolution (Grewal and Wright, 1992), and dauers (at least in *C. elegans*) use this information when deciding whether to resume development (Bubrig et al., 2020). Non-necromenic species might confine development to environments with their preferred bacterial community and might not recognize vector bacteria as a legitimate food source. Shifting food preferences could occur with changes in odorant receptors, which are highly mutable (Quignon et al., 2005), or in their expression.

Nematodes regulate and coordinate development via the exchange of pheromones among conspecifics (Golden and Riddle, 1984; Ludewig et al., 2019). A nematode's dauer state is maintained by nearby dauers. Nematodes that develop to adulthood signal nearby dauers to resume development (Ludewig et al., 2019). Intraspecific communication can therefore reinforce habitat use by coordinating development as a group. The habitat shift seen in necromeny could result from adjusting the production of pheromone components, which are known to be highly genotype and environment specific in both free-living and parasitic species (Kaplan et al., 2011; Mayer and Sommer, 2011; Stasiuk et al., 2012; Diaz et al., 2014). Similar group coordination may increase infection success in parasitic lineages (Campbell et al., 1999).

Chemosensation in nematodes depends on the environmental context (Bargmann, 2006), so interpreting vector bacteria as "food" may require integration with cues from the preferred substrate. For instance, nematodes living on rotting fruit might scan for plant oils, sugars, or high moisture in addition to bacterial food signals. Integrating vector-specific environmental cues or lessening the importance of environmental cues could play a role in necromeny.

STEP FOUR: PARASITISM

Parasites differ from phoretic or necromenic stages in that parasites feed and develop on living hosts. Necromeny may facilitate this exploitation by uncoupling vector and nematode fitness; the nematodes do not depend solely on their ancestral habitat. In some cases, parasites have free-living stages that still feed in the environment either facultatively like *Pristionchus lheritieri* (Geraert et al., 1989) or as part of an alternating life cycle like *Strongyloides stercoralis* (Yamada et al., 1991). In other cases, the facultative strategy may be used as a stepping stone to obligate parasitism and the abandonment of the ancestral habitat (Durkin and Luong, 2018, 2020; Luong and Mathot, 2019).

Parasitic nematodes use 2 main methods to feed on their host (direct feeding and indirect feeding), which suggests multiple ways a lineage could evolve host feeding. In indirect feeding, nematodes do not feed on host tissue. Instead, they weaponize their ancestral food source to kill the host. *Heterorhabditis* spp. harbor symbiotic bacteria that are harmless to them but highly pathogenic to their hosts. The bacteria kill the host and proliferate on the cadaver, after which the nematodes feed on the bacteria to grow and develop (Poinar, 1990). Traits of non-parasitic nematodes could facilitate the evolution of this strategy. Larvae developing into dauers plug both ends of their gut, which can trap bacteria inside (Riddle et al., 1981; Rae et al., 2008). If the bacterial species is asymptomatic to the nematode but pathogenic to the carrier species, dauers may inadvertently weaponize them when they expel the bacteria (Chantanao and Jensen, 1969). The relationship could be strengthened by selecting for bacteria that are less pathogenic to the nematode and more pathogenic to the carrier species, more likely to survive the nematode gut, and more likely to reassociate with the nematode again, such as if the bacterium regularly out-competes other bacterial species (Ffrench-Constant et al., 2000).

In direct feeding, nematodes switch from their ancestral food source (e.g., bacteria) to the host's tissue, likely via digestive enzymes (Geary and Thompson, 2001; Poulin and Randhawa, 2015). Animal parasitic nematodes co-opt endogenous digestive enzymes and secrete them into the environment rather than retain them internally (Blaxter, 2003). Plant-parasitic nematodes require additional enzymes like cellulases that are not endogenous (Danchin et al., 2010).

Regardless of the evolutionary pathway, some limited set of changes must facilitate basic host exploitation (Geary and Thompson, 2001). Once that threshold is crossed, newly evolved parasites encounter a new set of challenges like the host immune system, and many aspects of the ancestral lifestyle may no longer apply (Blaxter, 2003; Poulin and Randhawa, 2015). Over time, parasites might even target new tissues (Sukhdeo et al., 1997), jump to new hosts, or integrate new hosts into a complex life cycle (Sudhaus, 2018). The majority of parasitism genes we find in comparative genomic studies have been acquired in the millions of years after the initial transition to parasitism, potentially disguising the core set of genes that enabled initial exploitation (Blaxter, 2003; Opperman et al., 2008; Kikuchi et al., 2011). The original genes may not even be present anymore, their functions transferred and diluted to dozens of other genes (Viney, 2017). Therefore, studying the transition to parasitism becomes harder

the farther back exploitation began. To solve this, we can shift our attention from prototypical parasites and study extant species that are facultatively parasitic (Luong and Mathot, 2019) or in transitional phases (Stevens et al., 2020). Such cases help us distinguish between critical changes for parasitism and subsequent adaptation.

CONCLUSION

The problem with models of parasitism evolution is that they cannot be tested directly short of experimentally evolving free-living species into parasites (Viney, 2017). We can gain insight into the transition to parasitism by aggregating and interpreting data from related non-parasitic species. The process gives clues about when key parasitism traits likely arose and what their ancestral function may have been. While this article focused on 1 nematode model, the approach can benefit any models—such as alternative dauer-based hypotheses in Nematoda (Sudhaus, 2018) or those in other taxa like mites (Houck and OConnor, 1991)—provided the model's steps are well defined and we can find non-parasitic species that approximate them.

Our interpretations are only as good as the ecological data for non-parasitic species. In model nematodes, ecology and natural history have been vastly outpaced by genetics, genomics, and developmental biology (Kammenga et al., 2008; Félix and Braendle, 2010). This pattern is hardly a quirk of Nematoda (West et al., 1996; Whitfield, 1998; Proctor and Owens, 2000) and reflects a broader skew toward applied parasitology and away from evolutionary and ecological studies (Keymer and Read, 1990; Vickerman, 2009; Jackson, 2015). We can understand the evolution of parasitism only by tuning more into the basic ecology and natural history of parasitic species and their non-parasitic relatives.

ACKNOWLEDGMENTS

The authors would like to thank A. Gibson for helpful comments and guidance as well as M. Rockman for sharing unpublished results. LTB was supported by the Graduate Council Fellowship through the University of Alabama Graduate School. JLF was supported by the US National Science Foundation awards EF 1921585 and DEB 1941854.

LITERATURE CITED

- ADAMS, P. E., L. T. BUBRIG, AND J. L. FIERST. 2020. Genome evolution: On the road to parasitism. *Current Biology* 30: R272–R274.
- ALLDREDGE, A. L., AND M. W. SILVER. 1988. Characteristics, dynamics and significance of marine snow. *Progress in Oceanography* 20:41–82.
- ANDERSON, R. C. 1984. The origins of zooparasitic nematodes. *Canadian Journal of Zoology—Revue Canadienne de Zoologie* 62: 317–328.
- ANDERSON, R. C. 1988. Nematode transmission patterns. *Journal of Parasitology* 74: 30–45.
- ANDERSON, R. C. 1996. Why do fish have so few roundworm (nematode) parasites? *Environmental Biology of Fishes* 46: 1–5.
- ATHIAS-BINCHE, F., AND S. MORAND. 1993. From phoresy to parasitism: The example of mites and nematodes. *Research and Reviews in Parasitology* 53:73–79.
- BARGMANN, C. I. 2006. Chemosensation in *C. elegans*. In *WormBook*, E. Jorgensen (ed.). The *C. elegans* Research Community, WormBook, p. 1–29. doi:10.1895/wormbook.1.123.1.
- BARTLOW, A. W., AND S. J. AGOSTA. 2021. Phoresy in animals: Review and synthesis of a common but understudied mode of dispersal. *Biological Reviews* 96: 223–246.
- BLAXTER, M. L. 2003. Nematoda: Genes, genomes and the evolution of parasitism. *Advances in Parasitology* 54: 101–195.
- BLAXTER, M., AND G. KOUTSOVOULOS. 2015. The evolution of parasitism in Nematoda. *Parasitology* 142(Suppl. 1): S26–S39.
- BROWN, F. D., I. D'ANNA, AND R. J. SOMMER. 2011. Host-finding behaviour in the nematode *Pristionchus pacificus*. *Proceedings of the Royal Society B—Biological Sciences* 278: 3260–3269.
- BUBRIG, L. T., J. M. SUTTON, AND J. L. FIERST. 2020. *Caenorhabditis elegans* dauers vary recovery in response to bacteria from natural habitat. *Ecology and Evolution* 10: 9886–9895.
- CAMPBELL, J. F., AND R. GAUGLER. 1993. Nictation behavior and its ecological implications in the host search strategies of entomopathogenic nematodes (Heterorhabditidae and Steinernematidae). *Behaviour* 126: 155–169.
- CAMPBELL, J. F., A. M. KOPPENHOFER, H. K. KAYA, AND B. CHINNASRI. 1999. Are there temporarily non-infectious dauer stages in entomopathogenic nematode populations: A test of the phased infectivity hypothesis. *Parasitology* 118: 499–508.
- CAMPBELL, J. F., AND E. E. LEWIS. 2002. Entomopathogenic nematode host-search strategies. In *The Behavioural Ecology of Parasites*, E. E. Lewis, J. F. Campbell, and M. V. K. Sukhdeo (eds.). CABI Publishing, New York, New York, p. 13–38.
- CASSADA, R. C., AND R. L. RUSSELL. 1975. Dauerlarva, a post-embryonic developmental variant of nematode *Caenorhabditis elegans*. *Developmental Biology* 46:326–342.
- CHAISSON, K. E., AND E. A. HALLEM. 2012. Chemosensory behaviors of parasites. *Trends in Parasitology* 28: 427–436.
- CHANDLER, G. T., AND J. W. FLEEGER. 1983. Meiofaunal colonization of azoic estuarine sediment in Louisiana—Mechanisms of dispersal. *Journal of Experimental Marine Biology and Ecology* 69: 175–188.
- CHANTANAO, A., AND H. J. JENSEN. 1969. Saprozoic nematodes as carriers and disseminators of plant pathogenic bacteria. *Journal of Nematology* 1: 216–218.
- CINKORNPUMIN, J. K., D. R. WISIDAGAMA, V. RAPOPORT, J. L. GO, C. DIETERICH, X. Y. WANG, R. J. SOMMER, AND R. L. HONG. 2014. A host beetle pheromone regulates development and behavior in the nematode *Pristionchus pacificus*. *Elife* 3: e03229. doi:10.7554/eLife.03229.
- CROLL, N. A., AND B. E. MATTHEWS. 1977. *Biology of Nematodes*. Wiley, New York, New York, 201 p.
- CROOK, M. 2014. The dauer hypothesis and the evolution of parasitism: 20 years on and still going strong. *International Journal for Parasitology* 44: 1–8.
- DANCHIN, E. G. J., M. N. ROSSO, P. VIEIRA, J. DE ALMEIDA-ENGLER, P. M. COUTINHO, B. HENRISSAT, AND P. ABAD. 2010. Multiple lateral gene transfers and duplications have promoted plant parasitism ability in nematodes. *Proceedings of*

- the National Academy of Sciences of the United States of America 107: 17651–17656.
- DAVIS, E. L., R. S. HUSSEY, T. J. BAUM, J. BAKKER, AND A. SCHOTS. 2000. Nematode parasitism genes. *Annual Review of Phytopathology* 38: 365–396.
- DIAZ, S. A., V. BRUNET, G. C. LLOYD-JONES, W. SPINNER, B. WHARAM, AND M. VINEY. 2014. Diverse and potentially manipulative signalling with ascarosides in the model nematode *C. elegans*. *BMC Evolutionary Biology* 14: 46. doi:10.1186/1471-2148-14-46.
- DIETERICH, C., AND R. J. SOMMER. 2009. How to become a parasite—Lessons from the genomes of nematodes. *Trends in Genetics* 25: 203–209.
- DURKIN, E. S., AND L. T. LUONG. 2018. Experimental evolution of infectious behaviour in a facultative ectoparasite. *Journal of Evolutionary Biology* 31: 362–370.
- DURKIN, E. S., A. M. ROTH, AND C. N. KEISER. 2020. Parasitic personalities: Consistent individual differences in behavior in a facultatively parasitic mite. *Journal of Insect Behavior* 33: 14–19.
- ELLENBY, C. 1968. Desiccation survival of infective larva of *Haemonchus contortus*. *Journal of Experimental Biology* 49: 469–475.
- FARISH, D. J., AND R. C. AXTELL. 1971. Phoresy redefined and examined in *Macrocheles muscaedomesticae* (Acarina: Macrochelidae). *Acarologia* 13: 16–29.
- FÉLIX, M.-A., M. AILION, J.-C. HSU, A. RICHAUD, AND J. WANG. 2018. *Pristionchus* nematodes occur frequently in diverse rotting vegetal substrates and are not exclusively necromenic, while *Panagrellus redivivoides* is found specifically in rotting fruits. *PLoS ONE* 13: e0200851. doi:10.1371/journal.pone.0200851.
- FÉLIX, M.-A., AND C. BRAENDLE. 2010. The natural history of *Caenorhabditis elegans*. *Current Biology* 20: R965–R969.
- FFRENCH-CONSTANT, R. H., N. WATERFIELD, V. BURLAND, N. T. PERNA, P. J. DABORN, D. BOWEN, AND F. R. BLATTNER. 2000. A genomic sample sequence of the entomopathogenic bacterium *Photorhabdus luminescens* W14: Potential implications for virulence. *Applied and Environmental Microbiology* 66: 3310–3329.
- GANG, S. S., AND E. A. HALLEM. 2016. Mechanisms of host seeking by parasitic nematodes. *Molecular and Biochemical Parasitology* 208: 23–32.
- GEARY, T. G., AND D. P. THOMPSON. 2001. *Caenorhabditis elegans*: How good a model for veterinary parasites? *Veterinary Parasitology* 101: 371–386.
- GERAERT, E., Z. MRACEK, AND W. SUDHAUS. 1989. The facultative insect parasite *Pristionchus lheritieri* (syn. *P. pseudolheritieri*) (Nematoda: Diplogasterida). *Nematologica* 35: 133–141.
- GERLACH, S. A. 1977. Means of meiofauna dispersal. *Mikrofauna Meeresboden* 61: 89–103.
- GOLDEN, J. W., AND D. L. RIDDLE. 1984. The *Caenorhabditis elegans* dauer larva: Developmental effects of pheromone, food, and temperature. *Developmental Biology* 102: 368–378.
- GRANZER, M., AND W. HAAS. 1991. Host-finding and host recognition of infective *Ancylostoma caninum* larvae. *International Journal for Parasitology* 21: 429–440.
- GREWAL, P. S., AND D. J. WRIGHT. 1992. Migration of *Caenorhabditis elegans* (Nematoda: Rhabditidae) larvae towards bacteria and the nature of the bacterial stimulus. *Fundamental and Applied Nematology* 15: 159–166.
- GRIFFIN, C. T. 2012. Perspectives on the behavior of entomopathogenic nematodes from dispersal to reproduction: Traits contributing to nematode fitness and biocontrol efficacy. *Journal of Nematology* 44: 177–184.
- HAAS, W. 2003. Parasitic worms: Strategies of host finding, recognition and invasion. *Zoology* 106: 349–364.
- HAEGEMAN, A., J. T. JONES, AND E. G. J. DANCHIN. 2011. Horizontal gene transfer in nematodes: A catalyst for plant parasitism? *Molecular Plant-Microbe Interactions* 24: 879–887.
- HALLEM, E. A., A. R. DILLMAN, A. V. HONG, Y. J. ZHANG, J. M. YANO, S. F. DEMARCO, AND P. W. STERNBERG. 2011. A sensory code for host seeking in parasitic nematodes. *Current Biology* 21: 377–383.
- HARBISON, C. W., M. V. JACOBSEN, AND D. H. CLAYTON. 2009. A hitchhiker's guide to parasite transmission: The phoretic behaviour of feather lice. *International Journal for Parasitology* 39: 569–575.
- HEIP, C., M. VINCX, AND G. VRANKEN. 1985. The ecology of marine nematodes. In *Oceanography and Marine Biology: Annual Review*, M. Barnes (ed.). Aberdeen University Press, Aberdeen, U.K., p. 399–489.
- HIGHSMITH, R. C. 1985. Floating and algal rafting as potential dispersal mechanisms in brooding invertebrates. *Marine Ecology Progress Series* 25: 169–179.
- HOLTERMAN, M., G. KARSSSEN, S. VAN DEN ELSEN, H. VAN MEGEN, J. BAKKER, AND J. HELDER. 2009. Small subunit rDNA-based phylogeny of the Tylenchida sheds light on relationships among some high-impact plant-parasitic nematodes and the evolution of plant feeding. *Phytopathology* 99: 227–235.
- HONG, R. L., AND R. J. SOMMER. 2006. Chemoattraction in *Pristionchus* nematodes and implications for insect recognition. *Current Biology* 16: 2359–2365.
- HOTEZ, P., J. HAWDON, AND G. A. SCHAD. 1993. Hookworm larval infectivity, arrest and amphiparatenesis—The *Caenorhabditis elegans* daf-c paradigm. *Parasitology Today* 9: 23–26.
- HOUCK, M.-A., AND B. M. OCONNOR. 1991. Ecological and evolutionary significance of phoresy in the Astigmata. *Annual Review of Entomology* 36: 611–636.
- JACKSON, A. P. 2015. Preface. The evolution of parasite genomes and the origins of parasitism. *Parasitology* 142(Suppl. 1): S1–S5.
- JOHNSON, K. P., K. YOSHIZAWA, AND V. S. SMITH. 2004. Multiple origins of parasitism in lice. *Proceedings of the Royal Society B—Biological Sciences* 271: 1771–1776.
- JOHNSON, N. C., J. H. GRAHAM, AND F. A. SMITH. 1997. Functioning of mycorrhizal associations along the mutualism-parasitism continuum. *New Phytologist* 135: 575–586.
- JONES, M. G. K. 1981. Host cell responses to endoparasitic nematode attack: Structure and function of giant cells and syncytia. *Annals of Applied Biology* 97: 353–372.
- KAMMENG, J. E., P. C. PHILLIPS, M. DE BONO, AND A. DOROSZUK. 2008. Beyond induced mutants: Using worms to study natural variation in genetic pathways. *Trends in Genetics* 24: 178–185.
- KAPLAN, F., J. SRINIVASAN, P. MAHANTI, R. AJREDINI, O. DURAK, R. NIMALENDRAN, P. W. STERNBERG, P. E. TEAL, F. C. SCHROEDER, A. S. EDISON, ET AL. 2011. Ascaroside expression

- in *Caenorhabditis elegans* is strongly dependent on diet and developmental stage. *PLoS ONE* 6: e17804. doi:10.1371/journal.pone.0017804.
- KAUL, T. K., P. R. RODRIGUES, I. V. OGUNGBE, P. KAPAHI, AND M. S. GILL. 2014. Bacterial fatty acids enhance recovery from the dauer larva in *Caenorhabditis elegans*. *PLoS ONE* 9: e86979. doi:10.1371/journal.pone.0086979.
- KEYMER, A. E., AND A. F. READ. 1990. The evolutionary biology of parasitism. *Parasitology Today* 6: 2–3.
- KIKUCHI, T., J. A. COTTON, J. J. DALZELL, K. HASEGAWA, N. KANZAKI, P. McVEIGH, T. TAKANASHI, I. J. TSAI, S. A. ASSEFA, P. J. A. COCK, ET AL. 2011. Genomic insights into the origin of parasitism in the emerging plant pathogen *Bursaphelenchus xylophilus*. *PLoS Pathogens* 7(9): e1002219. doi:10.1371/journal.ppat.1002219.
- KIONTKE, K., AND D. H. A. FITCH. 2013. Nematodes. *Current Biology* 23: R862–R864.
- KIONTKE, K., AND W. SUDHAUS. 2006. Ecology of *Caenorhabditis* species. In *WormBook: The Online Review of C. elegans Biology*, D. H. A. Fitch (ed.). The *C. elegans* Research Community, p. 1–14. doi:10.1895/wormbook.1.37.1.
- KLASS, M., AND D. HIRSH. 1976. Non-aging developmental variant of *Caenorhabditis elegans*. *Nature* 260: 523–525.
- LAFFERTY, K. D., AND A. M. KURIS. 2002. Trophic strategies, animal diversity and body size. *Trends in Ecology & Evolution* 17: 507–513.
- LEE, H., M.-K. CHOI, D. LEE, H.-S. KIM, H. HWANG, H. KIM, S. PARK, Y.-K. PAIK, AND J. LEE. 2012. Nictation, a dispersal behavior of the nematode *Caenorhabditis elegans*, is regulated by IL2 neurons. *Nature Neuroscience* 15: 107–112.
- LUC, M., A. R. MAGGENTI, R. FORTUNER, D. J. RASKI, AND E. GERAERT. 1987. A reappraisal of Tylenchina (Nemata) 1. For a new approach to the taxonomy of Tylenchina. *Revue de Nématologie* 10:127–134.
- LUDEWIG, A. H., A. B. ARTYUKHIN, E. Z. APRISON, P. R. RODRIGUES, D. C. PULIDO, R. N. BURKHARDT, O. PANDA, Y. K. ZHANG, P. GUDIBANDA, I. RUVINSKY, ET AL. 2019. An excreted small molecule promotes *C. elegans* reproductive development and aging. *Nature Chemical Biology* 15: 838–845.
- LUONG, L. T., AND K. J. MATHOT. 2019. Facultative parasites as evolutionary stepping-stones towards parasitic lifestyles. *Biology Letters* 15: 20190058. doi:10.1098/rsbl.2019.0058.
- MAYER, M. G., AND R. J. SOMMER. 2011. Natural variation in *Pristionchus pacificus* dauer formation reveals cross-preference rather than self-preference of nematode dauer pheromones. *Proceedings of the Royal Society B—Biological Sciences* 278: 2784–2790.
- MAYER, W. E., L. N. SCHUSTER, G. BARTELMES, C. DIETERICH, AND R. J. SOMMER. 2011. Horizontal gene transfer of microbial cellulases into nematode genomes is associated with functional assimilation and gene turnover. *BMC Evolutionary Biology* 11:13. doi: 10.1186/1471-2148-11-13.
- McSORLEY, R. 2003. Adaptations of nematodes to environmental extremes. *Florida Entomologist* 86: 138–142.
- MOSER, J. C., H. KONRAD, T. KIRISITS, AND L. K. CARTA. 2005. Phoretic mites and nematode associates of *Scolytus multistriatus* and *Scolytus pygmaeus* (Coleoptera:Scolytidae) in Austria. *Agricultural and Forest Entomology* 7: 169–177.
- O'CONNOR, B. M. 1982. Evolutionary ecology of astigmatid mites. *Annual Review of Entomology* 27: 385–409.
- OPPERMAN, C. H., D. M. BIRD, V. M. WILLIAMSON, D. S. ROKHSAR, M. BURKE, J. COHN, J. CROMER, S. DIENER, J. GAJAN, S. GRAHAM, ET AL. 2008. Sequence and genetic map of *Meloidogyne hapla*: A compact nematode genome for plant parasitism. *Proceedings of the National Academy of Sciences in the United States of America* 105: 14802–14807.
- POINAR, G. O. 1983. *The Natural History of Nematodes*. Prentice Hall, Englewood Cliffs, New Jersey, 323 p.
- POINAR, G. O. 1990. Taxonomy and biology of Steinernematidae and Heterorhabditidae. In *Entomopathogenic Nematodes in Biological Control*, R. Gaugler and H. K. Kaya (eds.). CRC Press, Boca Raton, Florida, p. 23–58.
- POULIN, R. 2007. *Evolutionary Ecology of Parasites*. Princeton University Press, Princeton, New Jersey, 332 p.
- POULIN, R., AND H. S. RANDHAWA. 2015. Evolution of parasitism along convergent lines: From ecology to genomics. *Parasitology* 142(Suppl. 1): S6–S15.
- PROCTOR, H., AND I. OWENS. 2000. Mites and birds: Diversity, parasitism and coevolution. *Trends in Ecology & Evolution* 15: 358–364.
- QUIGNON, P., M. GIRAUD, M. RIMBAULT, P. LAVIGNE, S. TACHER, E. MORIN, E. RETOUT, A. S. VALIN, K. LINDBLAD-TOH, J. NICOLAS, ET AL. 2005. The dog and rat olfactory receptor repertoires. *Genome Biology* 6: R83. doi:10.1186/gb-2005-6-10-r83.
- RAE, R., M. RIEBESELL, I. DINKELACKER, Q. WANG, M. HERRMANN, A. M. WELLER, C. DIETERICH, AND R. J. SOMMER. 2008. Isolation of naturally associated bacteria of necromenic *Pristionchus* nematodes and fitness consequences. *Journal of Experimental Biology* 211: 1927–1936.
- REBECCHI, L., C. BOSCHETTI, AND D. R. NELSON. 2020. Extreme-tolerance mechanisms in meiofaunal organisms: A case study with tardigrades, rotifers and nematodes. *Hydrobiologia* 847: 2779–2799.
- REED, E. M., AND H. R. WALLACE. 1965. Leaping locomotion by an insect-parasitic nematode. *Nature* 206: 210–211.
- RIDDLE, D. L., M. M. SWANSON, AND P. S. ALBERT. 1981. Interacting genes in nematode dauer larva formation. *Nature* 290: 668–671.
- ROGERS, W. P., AND R. I. SOMMERVILLE. 1963. The infective stage of nematode parasites and its significance in parasitism. *Advances in Parasitology* 1: 109–177.
- ROMEO, T. 2008. *Bacterial Biofilms*. Springer, Berlin, Germany, 293 p.
- SCHULTE, F. 1989. The association between *Rhabditis necromena* Sudhaus & Schulte, 1989 (Nematoda: Rhabditidae) and native and introduced millipedes in south Australia. *Nematologica* 35: 82–89.
- SCIACCA, J., W. M. FORBES, F. T. ASHTON, E. LOMBARDINI, H. R. GAMBLE, AND G. A. SCHAD. 2002. Response to carbon dioxide by the infective larvae of three species of parasitic nematodes. *Parasitology International* 51: 53–62.
- SHERMAN, K. M., AND B. C. COULL. 1980. The response of meiofauna to sediment disturbance. *Journal of Experimental Marine Biology and Ecology* 46: 59–71.
- SHERMAN, K. M., J. A. REIDENAUER, D. THISTLE, AND D. MEETER. 1983. Role of a natural disturbance in an assemblage of

- marine free-living nematodes. *Marine Ecology Progress Series* 11: 23–30.
- SOUTHWOOD, T. R. E. 1977. Habitat, the templet for ecological strategies? *Journal of Animal Ecology* 46: 336–365.
- STASIUK, S. J., M. J. SCOTT, AND W. N. GRANT. 2012. Developmental plasticity and the evolution of parasitism in an unusual nematode, *Parastrongyloides trichosuri*. *EvoDevo* 3: 1. doi:10.1186/2041-9139-3-1.
- STEVENS, L., S. ROOKE, L. C. FALZON, E. M. MACHUKA, K. MOMANYI, M. K. MURUNGI, S. M. NJOROGE, C. O. ODINGA, A. OGENDO, J. OGOLA, E. M. FÈVRE, AND M. BLAXTER. 2020. The genome of *Caenorhabditis bovis*. *Current Biology* 30: 1023–1031.
- STIERNAGLE, T. 2006. Maintenance of *C. elegans*. In *C. elegans: A Practical Approach*, I. Hope (ed.). Oxford University Press, Oxford, U.K., p. 51–67.
- SUDHAUS, W. 2018. Various evolutionary avenues of Nematoda to parasitism in Gastropoda. *Soil Organisms* 90: 115–122.
- SUKHDEO, S. C., M. V. K. SUKHDEO, M. B. BLACK, AND R. C. VRIJENHOEK. 1997. The evolution of tissue migration in parasitic nematodes (Nematoda: Strongylida) inferred from a protein-coding mitochondrial gene. *Biological Journal of the Linnean Society* 61: 281–298.
- TSAI, I. J., M. ZAROWIECKI, N. HOLROYD, A. GARCIARRUBIO, A. SANCHEZ-FLORES, K. L. BROOKS, A. TRACEY, R. J. BOBES, G. FRAGOSO, E. SCIUTTO, ET AL. 2013. The genomes of four tapeworm species reveal adaptations to parasitism. *Nature* 496: 57–63.
- VICKERMAN, K. 2009. “Not a very nice subject.” Changing views of parasites and parasitology in the twentieth century. *Parasitology* 136: 1395–1402.
- VINEY, M. 2017. How can we understand the genomic basis of nematode parasitism? *Trends in Parasitology* 33: 444–452.
- VINEY, M., AND S. HARVEY. 2017. Reimagining pheromone signalling in the model nematode *Caenorhabditis elegans*. *PLOS Genetics* 13: e1007046. doi:10.1371/journal.pgen.1007046.
- WADSWORTH, W. G., AND D. L. RIDDLE. 1989. Developmental regulation of energy metabolism in *Caenorhabditis elegans*. *Development Biology* 132: 167–173.
- WARBURTON, E. M., AND D. A. ZELMER. 2010. Prerequisites for parasitism in Rhabditid nematodes. *Journal of Parasitology* 96: 89–94.
- WEINSTEIN, S. B., AND A. M. KURIS. 2016. Independent origins of parasitism in Animalia. *Biology Letters* 12: 20160324. doi:10.1098/rsbl.2016.0324.
- WEST, S. A., E. A. HERRE, D. M. WINDSOR, AND P. R. S. GREEN. 1996. The ecology and evolution of the New World non-pollinating fig wasp communities. *Journal of Biogeography* 23: 447–458.
- WHITE, P. S., L. MORRAN, AND J. DE ROODE. 2017. Phoresy. *Current Biology* 27: R578–R580.
- WHITFIELD, J. B. 1998. Phylogeny and evolution of host-parasitoid interactions in hymenoptera. *Annual Review of Entomology* 43: 129–151.
- WILSON, E. O., N. I. DURLACH, AND L. M. ROTH. 1958. Chemical releasers of necrophoric behavior in ants. *Psyche* 65: 108–114.
- WIRTH, S. 2009. Necromenic life style of *Histiostoma polypori* (Acari: Histiostomatidae). *Experimental and Applied Acarology* 49: 317–327.
- YAMADA, M., S. MATSUDA, M. NAKAZAWA, AND N. ARIZONO. 1991. Species-specific differences in heterogonic development of serially transferred free-living generations of *Strongyloides planiceps* and *Strongyloides stercoralis*. *Journal of Parasitology* 77: 592–594.
- YAO, M., J. ROSENFELD, S. ATTRIDGE, S. SIDHU, V. AKSENOV, AND C. D. ROLLO. 2009. The ancient chemistry of avoiding risks of predation. *Evolutionary Biology* 36: 267–281.
- YOUNG, E. C. 1961. Degeneration of flight-musculature in the corixidae and notonectidae. *Nature* 189: 328–329.