

SIMULTANEOUS DETECTION OF VAIRIMORPHA INVICTAE (MICROSPORIDIA: BURENELLIDAE) AND THELOHANIA SOLENOPSAE (MICROSPORIDIA: THELOHANIIDAE) IN FIRE ANTS BY PCR

Authors: Valles, Steven M., Oi, David H., Briano, Juan A., and Williams, David F.

Source: Florida Entomologist, 87(1) : 85-87

Published By: Florida Entomological Society

URL: [https://doi.org/10.1653/0015-4040\(2004\)087\[0085:SDOVIM\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2004)087[0085:SDOVIM]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SIMULTANEOUS DETECTION OF *VAIRIMORPHA INVICTAE* (MICROSPORIDIA: BURENELLIDAE) AND *THELOHANIA SOLENOPSAE* (MICROSPORIDIA: THELOHANIIDAE) IN FIRE ANTS BY PCR

STEVEN M. VALLES¹, DAVID H. OI¹, JUAN A. BRIANO² AND DAVID F. WILLIAMS¹

¹Center for Medical, Agricultural and Veterinary Entomology, USDA-ARS
1600 SW 23rd Drive, Gainesville, Florida, 32608, USA

²South American Biological Control Laboratory, USDA-ARS, Bolivar 1559 Hurlingham
Buenos Aires Province, Argentina

Microsporidia are obligate intracellular protozoan parasites of eukaryotes (Mathis 2000). Two species of microsporidia, *Thelohania solenopsae* (Knell et al. 1977) and *Vairimorpha invictae* (Jouvenaz and Ellis 1986) have been reported to be effective biological control agents against the fire ant, *Solenopsis invicta* (Williams et al. 1999, Briano and Williams 2002). Unfortunately, because the life cycles of these pathogens remain unknown, diagnosis is principally limited to microscopic examination of ant homogenates for the characteristic spore stage. This limitation has hampered epidemiological studies, the elucidation of potential intermediate hosts, and description of the complete life cycle. While a number of PCR-based methods have been developed for detection of *T. solenopsae* (Snowden et al. 2002, Valles et al. 2002) none are available for *V. invictae*. By exploiting nucleotide sequence differences in the 16S rRNA genes of *T. solenopsae* and *V. invictae*, we provide a PCR-based method capable of detecting infection of fire ants by either pathogen.

V. invictae-infected colonies of *S. invicta* were collected in Argentina (near San Javier, Santa Fe Province) in April 2003. Infections were determined by the observation of *V. invictae* spores in wet mount preparations of macerated adult ants under a phase-contrast microscope (400X, Briano and Williams 2002). In addition, 1 *S. invicta* (Santa Fe Province) and 2 *S. richteri* (Entre Rios Province) colonies with dual infections (*V. invictae* and *T. solenopsae*) were collected in Argentina in April 2003. *S. invicta* were keyed to species (Trager 1991) and verified as "invicta-like" by chemotaxonomy (Vander Meer and Lofgren 1990). Genomic DNA was extracted from adult ants as described by Valles et al. (2002).

PCR was carried out with primer pairs specific for the 16S rRNA gene of *T. solenopsae* (p1, 5'CGAAGCATGAAAGCGGAGC and p2, 5'CAGCATGTATATGCACTACTGGAGC) and *V. invictae* (p90, 5'CACGAAGGAGGATAACCACGGT and p93, CGCAATCAGTCTGTGAATCTCTTCA). The microsporidian-specific primers were designed by aligning the *T. solenopsae* (accession number AF134205) and *V. invictae* 16S rRNA gene sequences with the Vector NTI 7.1 program (Informax, Inc., Bethesda, MD) and choosing unique areas from each species. A published nucleotide sequence for

the 16S rRNA gene was available in GenBank for a *Vairimorpha* sp. thought to be *V. invictae* (accession number AF031539). To verify that this sequence corresponded to the *V. invictae* 16S rRNA gene, we amplified a fragment of the gene from *V. invictae* with primers p90 and p93. The 791 bp amplicon was purified by separation on a 1.2% agarose gel, ligated into pGEM-T easy (Promega, Madison, WI), and used to transform Solopack Gold supercompetent *E. coli* DH5α cells (Statagene, La Jolla, CA). Insert-positive clones were sequenced by the Interdisciplinary Center for Biotechnology Research, University of Florida. Three replicates were sequenced.

PCR was conducted by the hot start method in a PTC 100 thermal cycler (MJ Research, Waltham, MA) under the following optimized temperature regime: 1 cycle at 94°C for 2 min, then 35 cycles at 94°C for 15 sec, 55°C for 15 sec, and 68°C for 45 sec, followed by a final elongation step of 5 min at 68°C. The reaction was conducted in a 50-μl volume containing 2 mM MgCl₂, 200 μM dNTP mix, 1 unit of Platinum Taq DNA polymerase (Invitrogen, Carlsbad, CA), 0.4 μM of each primer, and 0.5 μl of the genomic DNA preparation (10 to 100 ng). PCR products were separated on a 1.2% agarose gel and visualized by ethidium bromide staining. For all experiments, positive and negative controls were run alongside treatments.

The fragment of the 16S rRNA gene that we amplified from *V. invictae* (host *S. invicta*) was identical to the sequence reported previously by Moser et al. (1998). Despite being found in *S. richteri*, they suspected that the microsporidian with which they were working was *V. invictae*. Indeed, Briano et al. (2002) reported that *T. solenopsae* and *V. invictae* could infect either ant species, *S. invicta* or *S. richteri*. This conclusion was confirmed by successful detection of *V. invictae* from *S. invicta* and *S. richteri* with *V. invictae*-specific primers, p90 and p93.

Figure 1 demonstrates the specificity of the primer pairs for each species 16S rRNA gene. As reported by Valles et al. (2002), the *T. solenopsae*-specific primer pair, p1 and p2, produced a 318 bp amplicon exclusively from *T. solenopsae*-infected *S. invicta* (column 2). Similar specificity was observed for the 16S rRNA gene of *V. invictae* with primers p90 and p93; a 791 bp amplicon was pro-




Fig. 1. Banding patterns on a 1.2% agarose gel after multiplex PCR with 16S rRNA-specific oligonucleotide primers. Column 1, molecular weight markers expressed as base pairs (bp); column 2, DNA prepared from *T. solenopsae*-infected *S. invicta*, oligonucleotide primers p1 and p2 (*T. solenopsae*-specific); column 3, DNA prepared from *T. solenopsae*-infected *S. invicta*, oligonucleotide primers p90 and p93 (*V. invictae*-specific); column 4, DNA prepared from *V. invictae*-infected *S. invicta*, oligonucleotide primers p1 and p2 (*T. solenopsae*-specific); column 5, DNA prepared from *V. invictae*-infected *S. invicta*, oligonucleotide primers p90 and p93 (*V. invictae*-specific); column 6, mixture of DNA prepared from *V. invictae*-infected *S. invicta* and *T. solenopsae*-infected *S. invicta*, oligonucleotide primers p1 and p2 (*T. solenopsae*-specific), and p90 and p93 (*V. invictae*-specific); column 7, DNA prepared from *T. solenopsae*- and *V. invictae*-infected *S. invicta*, oligonucleotide primers p1 and p2 (*T. solenopsae*-specific), and p90 and p93 (*V. invictae*-specific); column 8, DNA prepared from uninfected *S. invicta*, oligonucleotide primers p1 and p2, and p90 and p93.

duced exclusively from *V. invictae*-infected *S. invicta* (column 5). In cases where an ant colony was infected with both organisms, each microsporidian species could be discerned in a single multiplex reaction containing both primers sets (Fig. 1, lanes 6 and 7). Again, *V. invictae* was successfully detected in either *S. invicta* or *S. richteri*.

Microscopic detection of these microsporidia is labor intensive and limited to known stages of development. The multiplex PCR method to detect *T. solenopsae* and *V. invictae* offers a number of advantages over traditional microscopy, including, increased sensitivity, specificity, and the ability to identify all developmental stages. Thus, multiplex PCR decreases the risk of misidentification and will facilitate epizootiological studies concerned with these pathogens.

We thank Chuck Strong for technical assistance and R. Vander Meer for identifying the ant species by gas chromatography. We also thank R. M. Pereira and J. L. Capinera who provided helpful reviews of a previous version of the manuscript.

SUMMARY

A PCR-based method capable of detecting *The-lohania solenopsae* and/or *Vairimorpha invictae* infection in the red imported fire ant, *Solenopsis*

invicta, was developed. Multiplex PCR allows simultaneous detection of both species of microsporidia in a single reaction.

REFERENCES

- BRIANO, J. A., AND D. F. WILLIAMS. 2002. Natural occurrence and laboratory studies of the fire ant pathogen *Vairimorpha invictae* (Microsporida: Burenellidae) in Argentina. *Environ. Entomol.* 31: 887-894.
- BRIANO, J. A., D. F. WILLIAMS, D. H. OI, AND L. R. DAVIS, JR. 2002. Field host range of the fire ant pathogens *The-lohania solenopsae* (Microsporida: Thelohaniidae) and *Vairimorpha invictae* (Microsporida: Burenellidae) in South America. *Biol. Control* 24: 98-102.
- JOUVENAZ, D. P., AND E. A. ELLIS. 1986. *Vairimorpha invictae* n. sp. (Microsporida: Microsporida), a parasite of the red imported fire ant, *Solenopsis invicta* Buren (Hymenoptera: Formicidae). *J. Protozool.* 33: 457-461.
- KNELL, J. D., G. E. ALLEN, AND E. I. HAZARD. 1977. Light and electron microscope study of *The-lohania solenopsae* n. sp. (Microsporida: Protozoa) in the red imported fire ant, *Solenopsis invicta*. *J. Invertebr. Pathol.* 29: 192-200.
- MATHIS, A. 2000. Microsporida: emerging advances in understanding the basic biology of these unique organisms. *Int. J. Parasitol.* 30: 795-804.
- MOSER, B. A., J. J. BECNEL, J. MARUNIAK, AND R. S. PATTERSON. 1998. Analysis of the ribosomal DNA se-

- quences of the microsporidia *Thelohania* and *Vairimorpha* of fire ants. J. Invertebr. Pathol. 72: 154-159.
- SNOWDEN, K. F., K. LOGAN, AND S. B. VINSON. 2002. Simple, filter-based PCR detection of *Thelohania solenopsae* (Microspora) in fire ants (*Solenopsis invicta*). J. Eukaryot. Microbiol. 49: 447-448.
- TRAGER, J. C. 1991. A revision of the fire ants, *Solenopsis geminata* group (Hymenoptera: Formicidae: Myrmicinae). J. New York Entomol. Soc. 99: 141-198.
- VALLES, S. M., D. H. OI, O. P. PERERA, AND D. F. WILLIAMS. 2002. Detection of *Thelohania solenopsae* (Microsporidia: Thelohaniidae) in *Solenopsis invicta* (Hymenoptera: Formicidae) by multiplex PCR. J. Invertebr. Pathol. 81: 196-201.
- VANDER MEER, R. K., AND C. S. LOFGREN. 1990. Chemotaxonomy applied to fire ant systematics in the United States and South America. pp. 75-84. In R. Vander Meer, K. Jaffe and A. Cedeno [eds.]. Applied myrmecology, a world perspective. Westview Press, Boulder, CO. 741 pp.
- WILLIAMS, D. F., G. J. KNUE, AND J. J. BECNEL. 1999. Discovery of *Thelohania solenopsae* from the imported fire ant, *Solenopsis invicta*, in the United States. J. Invertebr. Pathol. 71: 175-176.