



NATIVE HYMENOPTERAN PARASITOIDS ASSOCIATED WITH FRUIT FLIES (DIPTERA: TEPHRITIDAE) IN SANTA CATARINA STATE, BRAZIL

Authors: Garcia, Flávio R. M., and Corseuil, Elio

Source: Florida Entomologist, 87(4) : 517-521

Published By: Florida Entomological Society

URL: [https://doi.org/10.1653/0015-4040\(2004\)087\[0517:NHPAWF\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2004)087[0517:NHPAWF]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

NATIVE HYMENOPTERAN PARASITIDS ASSOCIATED WITH FRUIT FLIES
(DIPTERA: TEPHRITIDAE) IN SANTA CATARINA STATE, BRAZILFLÁVIO R. M. GARCIA¹ AND ELIO CORSEUIL²¹Unochapecó, CCAA, Laboratório de Entomologia, Caixa Postal 747, CEP. 89809-000, Chapecó, SC, Brazil
E-mail: flaviog@unochapeco.edu.br²PUCRS, Faculdade de Biociências, PPG em Biociências, Laboratório de Entomologia
Av. Ipiranga, 6681, Prédio 12c, CEP. 90619-900, Porto Alegre, RS, Brazil
E-mail: corseuil@pucrs.br

ABSTRACT

A survey of tephritid fruit fly species and their parasitoids was conducted in the western portion of Santa Catarina state, Brazil. A total of 9,197 fruits belonging to 46 plant species in 24 families was collected. The parasitoids recovered were *Aganaspis pelleranoi* (Brèthes), *Lopheucoila anastrephae* (Rhower) (Figitidae), *Doryctobracon areolatus* (Szèpligeti), *Doryctobracon brasiliensis* (Szèpligeti), *Odontosema anastrephae* Borgmeier, *Opius bellus* Gahan, *Opius* sp., *Utetes (Bracanastrepha) anastrephae* (Viereck) (Braconidae), and *Trichopria anastrephae* Lima (Diapriidae). *Aganaspis pelleranoi* is the most frequent parasitoid species found in the west region of Santa Catarina. We recorded the first occurrence of *L. anastrephae* in Santa Catarina. Parasitism ranged from 1.2 to 46.9%.

Key Words: Insecta, *Anastrepha*, Braconidae, Figitidae, Diapriidae, parasitoids, fruit flies.

RESUMEN

Se realizo un estudio del reconocimiento de las especies tefrítidas de las moscas de las frutas y sus parasitoides en el área occidental del estado de Santa Catarina, Brasil. Un total de 9,197 frutas que pertenecen a 46 especies de plantas fueron recolectadas. Los parasitoides recolectados fueron: *Aganaspis pelleranoi* (Brèthes), *Lopheucoila anastrephae* (Rhower) (Figitidae), *Doryctobracon areolatus* (Szèpligeti); *Doryctobracon brasiliensis* (Szèpligeti), *Odontosema anastrephae* Borgmeier, *Opius bellus* Gahan, *Opius* sp., *Utetes (Bracanastrepha) anastrephae* (Viereck) (Braconidae) y *Trichopria anastrephae* Lima (Diapriidae). *Aganaspis pelleranoi* es el parasitoide encontrado mas frecuentemente en la región oeste de Santa Catarina. El primer registro de la ocurrencia de *L. anastrephae* en Santa Catarina fue obtenido. El rango de parasitismo fue de 1.2 a 46.9%.

Surveys of tephritid fruit flies and their parasitoids are a first step to a better understand of the ecology of these economically important taxa (Zucchi 2000a). The major natural enemies of the fruit flies in Brazil belong to the families Braconidae, Figitidae, and Pteromalidae (Hymenoptera). Pteromalidae, mainly *Pachycrepoideus vindemiae* Rondani, are sporadically collected and specimens of Figitidae are collected in small numbers throughout the country. Typically the most frequent parasitoids collected in Brazil are members of the Braconidae family (Canal & Zucchi 2000).

In Santa Catarina state, Nora et al. (2000) previously found, in order of abundance, *Doryctobracon areolatus* (Szèpligeti), *Doryctobracon brasiliensis* (Szèpligeti), *Opius bellus* Gahan, and *Opius tomoplagiae* Lima. In addition, they obtained unidentified Diapriidae, Eulophidae, Figitidae, and Pteromalidae. Guimarães et al. (2000) added to this list the figitids *Aganaspis pelleranoi* (Brèthes) and *Odontosema anastrephae* Borgmeier, and Leonel, Jr. et al. (1995) and Canal & Zucchi (2000) obtained braconids *Microcrasis lon-*

chaea (Lima) and *Utetes (Bracanastrepha) anastrephae* (Viereck).

The present paper describes the parasitoids of fruit flies from the western portion of Santa Catarina state, an area which is a growing producer of citrus in the state (Koller et al. 1999) and which has not been surveyed thoroughly in the past. We collected 9,197 mature fruit from trees or on the soil comprising 46 species belonging to 25 families in the six towns Anchieta (26°53'S and 53°33'W), Chapecó (27°06'S and 53°16'W), Cunha Porã (26°07'S and 53°16'W), Palmitos (27°06'S and 53°16'W), São Carlos (27°07'S and 53°00'W), and Xanxerê (26°87'S and 52°40'W), Santa Catarina. Each fruit was weighed and put in a plastic container with about seven centimeters of sterilized sand, and covered with a net. The containers were kept in the entomology laboratory of the Agricultural and Environmental Science Center at the Universidade Comunitária Regional de Chapecó at 25 ± 3°C, 70 ± 10% and a 12-h photoperiod. After five days, the sand with pupae was transferred to lab Petri dishes containing filter

TABLE 1. PARASITOID SPECIES (HYMENOPTERA) FROM FRUIT FLIES COLLECTED IN SIX DIFFERENT LOCALITIES FROM THE WEST OF SANTA CATARINA, BRAZIL, DURING 1998-2000.

Vegetal specie	S ¹	N ²	PaT ³	Hymenoptera parasitoid species and relative frequency (%)			
				Braconidae	Figitidae and Diapriidae	%P ⁴	TIP ⁵
Fabaceae							
<i>Inga sellowiana</i>	5	246	2		<i>L. anastrephae</i> (100.0)	4.1	4.7
Myrtaceae							
<i>Psidium cattleianum</i>	11	635	42	<i>D. areolatus</i> (19.0) <i>D. brasiliensis</i> (33.3) <i>Opius</i> sp. (9.7) <i>U. anastrephae</i> (19.0)	<i>A. pelleranoi</i> (19.0)	1.9	2.1
<i>Eugenia involucrata</i>	3	446	46	<i>D. areolatus</i> (69.6) <i>Opius bellus</i> (8.7) <i>Opius</i> sp. (4.3) <i>U. anastrephae</i> (17.4)		46.9	47.9
<i>Psidium guajava</i>	17	190	147	<i>D. brasiliensis</i> (2.0) <i>Opius bellus</i> (1.1) <i>Opius</i> sp. (2.0)	<i>A. pelleranoi</i> (49.0) <i>T. anastrephae</i> (45.9)	20.1	20.3
<i>Feijoa sellowiana</i>	2	80	58	<i>D. areolatus</i> (48.3) <i>D. brasiliensis</i> (48.3) <i>Opius bellus</i> (3.4)		11.6	14.3
<i>Myrcianthes pungens</i>	2	52	6	<i>D. brasiliensis</i> (100.0)		28.5	46.1
<i>Campomanesia xanthocarpa</i>	4	702	4	<i>D. brasiliensis</i> (100.0)		7.7	12.5
<i>Britoa guazumaefolia</i>	6	255	48	<i>D. areolatus</i> (16.7) <i>D. brasiliensis</i> (37.5)	<i>A. pelleranoi</i> (3.5) <i>O. anastrephae</i> (8.3)	7.1	8.8
<i>Eugenia pyriformis</i>	5	264	4		<i>A. pelleranoi</i> (100.0)	3.0	4.0
Rosaceae							
<i>Prunus domestica</i>	5	109	26	<i>D. brasiliensis</i> (23.1) <i>U. anastrephae</i> (76.1)		16.1	16.8
<i>Prunus avium</i>	2	18	8	<i>D. areolatus</i> (75.0) <i>U. anastrephae</i> (25.0)		24.2	50.0

¹S—sample, ²n—number of fruit, ³PaT—parasitoids total, ⁴%P—Parasitism percentage, ⁵TIP—total index Parasitism.

TABLE 1. (CONTINUED) PARASITOID SPECIES (HYMENOPTERA) FROM FRUIT FLIES COLLECTED IN SIX DIFFERENT LOCALITIES FROM THE WEST OF SANTA CATARINA, BRAZIL, DURING 1998-2000.

Vegetal specie	S ¹	N ²	PaT ³	Hymenoptera parasitoid species and relative frequency (%)			
				Braconidae	Figitidae and Diapriidae	%P ⁴	TIP ⁵
<i>Eriobotrya japonica</i>	9	1166	48	<i>D. areolatus</i> (8.3)		4.3	5.9
				<i>D. brasiliensis</i> (41.6)			
				<i>Opius bellus</i> (4.3)			
				<i>Opius</i> sp. (4.2)			
				<i>U. anastrephae</i> (41.6)			
<i>Pyrus communis</i>	2	62	4		<i>A. pelleranoi</i> (100.0)	14.8	33.3
<i>Prunes persica</i>	16	562	18	<i>D. brasiliensis</i> (22.2)	<i>A. pelleranoi</i> (66.7)	1.2	1.6
					<i>O. anastrephae</i> (11.1)		
Total	89	4787	461				

¹S—sample, ²n—number of fruit, ³PaT—parasitoids total, ⁴%P—Parasitism percentage, ⁵TIP—total index Parasitism.

paper dampened with distilled water. Flies and parasitoids were counted after seven days.

The relationship between a fly species and its parasitoids was determined only when a single species of fly was held in an emergence container (Canal et al. 1994).

The total index of parasitism (TIP) was calculated as the number of parasitoids emerged \times 100/ number of flies emerged + number of parasitoids emerged. The relative frequency of fly species and parasitoids (RF) was defined as number of samples of a given species collected \times 100/total number of collected species according to Matrangolo et al. (1998), and the parasitism percentage was calculated as %P = total parasitism \times 100/total pupae, which was modified from Silveira Neto et al. (1997).

Species of *Anastrepha* were identified with Steyskal's key (1997) and Zucchi's key (2000b), which includes only Brazil species. The Braconidae were identified according to the key of Canal & Zucchi (2000). The flies and parasitoids belonging to other families were sent to Prof. Dr. Manoel Araécio Uchôa Fernandes, Biologist Jorge Anderson Guimarães, Dr. Allen Norrbom, and Prof. Dr. Roberto Antonio Zucchi for identification.

Of the 46 fruit species collected, 35 were infested by fruit flies, but only 14 of these fruit species contained parasitoids (Table 1). A total of 682 samples of parasitoids belonging to nine species and three families were obtained, as follows: *D. areolatus*, *D. brasiliensis*, *O. bellus*, *Opius* sp., *U. anastrephae* (Braconidae), *A. pelleranoi*, *O. anastrephae* (Figitidae), and *Trichopria anastrephae* Lima (Diapriidae).

Of the 461 hymenopterans associated with a particular fruit fly, *A. pelleranoi* was the most common and represented 25.6% of the total, followed by *D. brasiliensis* (21.1%) and *D. areolatus* (18.6%). These figures differ from those obtained by Salles (1996) for the state of Rio Grande do Sul and Leonel Jr. (1995, 1996) for the state of São Paulo. However, the relative abundance of parasitoids collected in the valley of Rio do Peixe, state of Santa Catarina by Nora et al. (2000) are similar to ours.

All the parasitoids developed in *A. fraterculus*, except for *L. anastrephae*, which was associated only with *Neosilba* sp. Also a lonchaeid was obtained in the samples (Table 2). *D. areolatus* and *U. anastrephae* also parasitized *Neosilba* sp. De Santis (1980) catalogued eleven species of *Trichopria* for Brazil, and believed that only *T. anastrephae* parasitized the genus *Anastrepha*. Aguiar-Menezes et al. (2001) previously reported *A. fraterculus* as the host of *T. anastrephae*. *L. anastrephae* has been recorded previously only in the southern part of the country, in the state of São Paulo and central Mato Grosso do Sul (De Santis 1980).

Parasitoids attacked fruit flies in 14 host-plant species, one Fabaceae, five Rosaceae and eight Myrtaceae (Table 1). Myrtaceae contained 79.6%

TABLE 2. ASSOCIATION BETWEEN COLLECTED PARASITIDS AND FRUIT FLIES IN SIX DIFFERENT LOCALITIES IN THE WEST OF SANTA CATARINA, BRAZIL, DURING 1998-2000.

Parasitoids	Fruit flies	
	<i>A. fraterculus</i>	<i>Neosilba</i> sp.
Braconidae		
<i>D. areolatus</i>	X	X
<i>D. brasiliensis</i>	X	
<i>O. bellus</i>	X	
<i>Opius</i> sp.	X	
<i>U. anastrephae</i>	X	X
Diapriidae		
<i>T. anastrephae</i>	X	
Figitidae		
<i>A. pelleranoi</i>	X	
<i>L. anastrephae</i>		X
<i>O. anastrephae</i>	X	

of the parasitoids and Rosaceae had 21.2%. The total parasitism index and the parasitism percentage were greatest in *Eugenia involucrata*, followed by *Prunus avium* and *Myrcianthes pungens*. These indexes are higher than those obtained by Leonel, Jr. et al (1996), Salles (1996) and Matrangolo et al. (1998), but are similar to those of Guimarães et al. (1999).

The high percent parasitism in *E. involucrata* was previously found by Salles (1996) in the Rio Grande do Sul state, and may be due, as he suggested, to the thin peel and small size of the fruit. Guimarães et al. (2000) previously observed affinity of *A. pelleranoi* to the Myrtaceae fruit, which we confirmed. Sivinski (1991), Sivinski et al. (1997, 2000) and Hickel (2002) found that braconid parasitism was negatively correlated to fruit pulp thickness, and we showed that weight data correlated to parasitism.

ACKNOWLEDGMENTS

We thank the following researchers for taxonomic identification: Professor Dr. Manoel Araécio Uchôa Fernandes—UFMS (Lonchaeidae), Biologist Jorge Anderson Guimarães—ESALQ/USP (Figitidae), Dr. Allen Norrbom—Systematic Entomology Laboratory, and Professor Dr. Roberto Antonio Zucchi—ESALQ/USP for some *Anastrepha* species confirmation. We thank Professors Lúcia S. Verona and Rosiane B. Denardin from the Botanic Department of UNOCHAPECÓ for identification of the plants.

REFERENCES CITED

- AGUIAR-MENEZES, E., E. B. MENEZES, P. S. SILVA, A. C. BITTAR, AND P. C. CASSINO. Native hymenopteran parasitoids associated with *Anastrepha* spp.

- (Diptera: Tephritidae) in Seropedica city, Rio de Janeiro Brazil. *Florida Entomol.* 84: 706-711.
- CANAL, N. A., AND R. A. ZUCCHI. 2000. Parasitóides Braconidae, pp. 119-126 *In* A. Malavasi and R. A. Zucchi [eds.], Moscas-das-frutas de Importância Econômica no Brasil, Conhecimento Básico e Aplicado. Holos, Ribeirão Preto.
- CANAL, N. A., R. A. ZUCCHI, N. M. SILVA, AND F. L. LEONEL, JR. 1994. Reconocimiento de las especies de parasitoides (Hym.: Braconidae) de moscas de las frutas (Dip.: Tephritidae) en dos municipios del Estado de Amazonas, Brasil. *Bol. Mus. Entomol. Universidad del Valle* 2: 1-17
- DE SANTIS, L. 1980. Catalogo de los Himenopteros Brasileños de la Serie Parasítica Incluyendo Bethyloidea. UFPR, Curitiba. 395 pp.
- GUIMARÃES, J. A., N. B. DIAZ, AND R. A. ZUCCHI. 2000. Parasitóides—Figitidae (Eucoilinae), pp. 127-134 *In* A. Malavasi and R.A. Zucchi [eds.], Moscas-das-frutas de Importância Econômica no Brasil, Conhecimento Básico e Aplicado. Holos, Ribeirão Preto.
- GUIMARÃES, J. A., R. A. ZUCCHI, N. B. DIAZ, M. F. DE SOUZA FILHO, AND M. A. UCHÔA F. 1999. Espécies de Eucoilinae (Hymenoptera: Cynipoidea: Figitidae) parasitóides de larvas frugívoras (Diptera: Tephritidae e Lonchaeidae) no Brasil. *An. Soc. Entomol. Brasil* 28: 263-273.
- HICKEL, E. R. 2002. Espessura da polpa como condicionante do parasitismo de moscas-das-frutas (Diptera: Tephritidae) por Hymenoptera: Braconidae. *Ciência Rural* 32: 1005-1009
- KOLLER, O. L., E. SOPRANO, L. A. F. VERONA, L. A. CHIARADIA, A. BRANCHER, AND E. N. DE OLIVEIRA. 1999. Citros, pp. 49-52 *In* EPAGRI, Recomendações de Cultivares para o Estado de Santa Catarina 1999/2000. EPAGRI, Florianópolis.
- LEONEL JR., F. L., R. A. ZUCCHI, AND N. A. D. CANAL. 1996. Parasitismo de moscas-das-frutas por Braconidae (Hymenoptera) em duas localidades do Estado de São Paulo. *An. Soc. Entomol. Brasil* 25: 199-206.
- LEONEL JR., F.L., R. A. ZUCCHI, AND R. A. WHARTON. 1995. Distribution and tephritid hosts (Diptera) of braconid parasitoids (Hymenoptera) in Brazil. *Intl. J. Pest Management* 41: 208-213
- MATRANGOLO, W. J. R., A. S. NASCIMENTO, R. S. CARVALHO, E. D. MELO, AND M. DE JESUS. 1998. Parasitóides de moscas-das-frutas (Diptera: Tephritidae) associados a frutíferas tropicais. *An. Soc. Entomol. Brasil* 27: 593-603.
- NORA, I., E. R. HICKEL, AND H. F. PRANDO. 2000. Moscas-das-frutas nos estados brasileiros: Santa Catarina, pp. 271-275 *In* A. Malavasi and R. A. Zucchi [eds.], Moscas-das-frutas de Importância Econômica no Brasil, Conhecimento Básico e Aplicado. Holos, Ribeirão Preto.
- SALLES, L. A. B. 1996. Parasitismo de *Anastrepha fraterculus* (Wied.) (Diptera: Tephritidae) por Hymenoptera, na região de Pelotas, RS. *Pesq. Agrop. Brasileira* 31: 769-774.
- SILVEIRA NETO, S., NAKANO, O., BARBIN, D., AND N. A. VILLA NOVA. 1976. Manual de Ecologia dos Insetos. Agronômica Ceres, Piracicaba, 419 pp.
- SIVINSKI, J. 1991. The influence of host fruit morphology on parasitism rates in the Caribbean fruit fly (*Anastrepha suspensa* (Loew)). *Entomophaga* 36: 447-455
- SIVINSKI, J. M., M. ALUJA, AND M. LÓPEZ. 1997. Spatial and temporal distribution of Mexican *Anastrepha* species (Diptera: Tephritidae), within the canopies of fruit trees. *Ann. Entomol. Soc. Amer.* 90: 604-618.
- SIVINSKI, J. M., J. PIÑERO, AND M. ALUJA. 2000. The distributions of parasitoids (Hymenoptera) of *Anastrepha* fruit flies (Diptera: Tephritidae) along an altitudinal gradient in Veracruz, Mexico. *Biological control* 18: 258-269.
- STEYSKAL, G. C. 1977. Pictorial Key to Species of the Genus *Anastrepha* (Diptera: Tephritidae). *The Entomol. Soc. of Washington, Washington, D.C.* 35 pp.
- ZUCCHI, R. A. 2000a. Espécies de *Anastrepha*, sinônímias, plantas hospedeiras e parasitóides, pp. 41-48 *In* A. Malavasi and R.A. Zucchi [eds.], Moscas-das-frutas de Importância Econômica no Brasil, Conhecimento Básico e Aplicado. Holos, Ribeirão Preto.
- ZUCCHI, R. A. 2000b. Taxonomia, pp. 13-24 *In* A. Malavasi and R.A. Zucchi [eds.], Moscas-das-frutas de Importância Econômica no Brasil, Conhecimento Básico e Aplicado. Holos, Ribeirão Preto.