



**RED IMPORTED FIRE ANTS (HYMENOPTERA:
FORMICIDAE) AT GOPHER TORTOISE (TESTUDINES:
TESTUDINIDAE) BURROWS**

Authors: Wetterer, James K., and Moore, Jon A.

Source: Florida Entomologist, 88(4) : 349-354

Published By: Florida Entomological Society

URL: [https://doi.org/10.1653/0015-4040\(2005\)88\[349:RIFAHF\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2005)88[349:RIFAHF]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

RED IMPORTED FIRE ANTS (HYMENOPTERA: FORMICIDAE) AT GOPHER TORTOISE (TESTUDINES: TESTUDINIDAE) BURROWS

JAMES K. WETTERER AND JON A. MOORE

Wilkes Honors College, Florida Atlantic University, 5353 Parkside Drive, Jupiter, FL 33458

ABSTRACT

The gopher tortoise, *Gopherus polyphemus* Daudin, is endemic to the southeastern US, where its populations are declining primarily due to habitat destruction. Tortoises are preyed upon by many species, including the red imported fire ant, *Solenopsis invicta* Buren, a destructive exotic species now common throughout the tortoises' entire range. We surveyed ants using tuna bait at 154 *G. polyphemus* burrows in a greenway reserve established to protect the tortoises in a residential area of southeast Florida. We found *S. invicta* present, typically recruiting to the bait in very high numbers, on the aprons of 33% of the tortoise burrows. *Solenopsis invicta* occurred significantly more often at burrows within 30 m of the greenway's outer edge than at burrows in more interior parts of the greenway (57% versus 16%). Among the interior burrows, *S. invicta* occurred significantly more often at burrows directly on two narrow strips of disturbed habitat, along an old fence line and an old pipeline, than at burrows not on these two strips (46% versus 12%). The greenway interior appears to offer tortoises and other species some refuge from *S. invicta*. However, the long thin design typical of greenways, the inclusion of walking paths through the greenways, and the policies of prescribed burning and reduction mowing used to maintain open habitat for the tortoises all may increase the tortoises' exposure to *S. invicta*. *Solenopsis invicta* is also a grave threat to other native species in these reserves, including the many animals that obligately live inside gopher tortoise burrows.

Key Words: edge effect, exotic species, Florida, *Gopherus*, greenway, *Solenopsis invicta*.

RESUMEN

La especie tortuga gopher, *Gopherus polyphemus* Daudin, es endémica al sudeste de los EE.UU., donde sus poblaciones disminuyen principalmente debido a la destrucción de hábitat. Las tortugas son atacadas por muchas especies, inclusive la hormiga roja de fuego importada, la *Solenopsis invicta* Buren, una especie exótica destructiva ahora común a toda la región geográfica de las tortugas. Estudiamos las hormigas que atraemos con cebo de atún en 154 madrigueras de la *G. polyphemus* en una reserva ecológica que se estableció para proteger las tortugas en un área residencial del sudeste de la Florida. Encontramos a la *S. invicta* presente, alistando típicamente al cebo en números muy altos, en los delantales de 33% de las madrigueras de las tortugas. La presencia de la *S. invicta* ocurrió apreciablemente más a menudo en madrigueras dentro de un parámetro de 30 m de la orilla exterior de la reserva que en las madrigueras en partes más interiores de la reserva (57% contra 16%). Entre las madrigueras interiores, la *S. invicta* ocurrió apreciablemente más a menudo en madrigueras directamente en dos tiras estrechas del hábitat perturbado, por una línea vieja de la cerca y una tubería vieja, que en las madrigueras que no en estaban en estas dos tiras (46% contra 12%). El interior de la reserva ecológica aparece ofrecer las tortugas y otras animales algún refugio de *S. invicta*. El largo y estrecho diseño típico de las reservas ecológicas, la inclusión de senderos caminantes en las reservas ecológicas, y por la costumbre de incendios prescritos y la reducción por cortes utilizados para mantener el hábitat abierto para las tortugas, todo ello puede aumentar la exposición de las tortugas a la *S. invicta*. La *S. invicta* es también una amenaza grave a otras especies nativas en estas reservas, inclusive los muchos animales que se ven obligados a vivir en las madrigueras de las tortugas.

Translation provided by the authors.

Gopher tortoises, *Gopherus polyphemus* Daudin, are endemic to the coastal plains of the southeastern United States (Diemer 1992) and populations are declining through most of their range, primarily due to habitat destruction (Diemer 1986). The tortoises' decline has been particularly great in southeastern Florida, where

most of the tortoise's preferred upland habitat has been converted to, for example, citrus orchards, housing developments, and phosphate mines (Diemer 1992). The gopher tortoise is now protected in Florida as a "species of special concern" (Diemer 1992; Florida Game & Fresh Water Fish Commission 1994).

Gopher tortoises prefer habitats with sandy soils for burrows, herbaceous plants for food, and open sunny spots for nesting and basking (Diemer 1992). Fire is a normal element in gopher tortoise habitat, and when natural fires are suppressed, habitats may become overgrown. To maintain the open spaces necessary for herbaceous vegetation and tortoise nesting and basking sites, land managers often use prescribed burning or reduction mowing (Wade & Lunsford 1989; Main & Tanner 1999).

Female tortoises lay clutches of 3-15 eggs, usually in the sand apron in front of their burrow or in another nearby sunny spot (typically 16-67 cm from the burrow entrance; Butler and Hull 1996). Many predators attack gopher tortoise nests, including raccoons, foxes, skunks, armadillos, and snakes (Douglass & Winegarner 1977; Diemer 1992). Nest predation is often very high (Alford 1980). Landers et al. (1980) found that mammalian predators destroyed most gopher tortoise nests (87% within the first few weeks after laying) and the few surviving hatchlings were often attacked and killed by the red imported fire ant, *Solenopsis invicta* Buren. Epperson & Heise (2003) found that *S. invicta* predation accounted for 27% of post-hatching mortality of gopher tortoise hatchlings.

Solenopsis invicta is probably the most destructive exotic ant species in the southeastern US, negatively impacting both invertebrates and vertebrates (Wojcik 1994; Porter & Savignano 1990). This invasive species arrived in Alabama by ship from South America and has spread across the southeast US from Texas to North Carolina, now occupying the entire range of the gopher tortoise.

Solenopsis invicta is a well-known predator on bird hatchlings, particularly ground-nesting species (e.g., see Ridlehuber 1982; Sikes & Arnold 1986; Steigman 1993; Drees 1994; Lockley 1995; Powell 1995; Dickinson 1995; Giuliano et al 1996; Mueller et al. 1999; Kopachena et al. 2000; H. Smith, pers. comm.). *Solenopsis invicta* also attacks reptile hatchlings. Cintra (1985) found that *S. invicta* commonly attacked and killed hatchling caiman (*Caiman yacare* (Daudin)). Love (1997) noted records of *S. invicta* attacking captive tortoises and hatching turtles and alligators. Allen et al. (1997) found that hatchling American alligators (*Alligator mississippiensis* (Daudin)) stung by fire ants showed decreased weight gain and increased mortality. Reagan et al. (2000) found that *S. invicta* in alligator nests had a significant negative impact on hatching success. *Solenopsis invicta* also appears to be an important threat to hatching sea turtles (Wilmers et al. 1996; Moulis 1996; Allen et al. 1998, 2001; Krahe et al. 2003; Wetterer & Wood 2005).

Solenopsis invicta appears to be an important threat to some adult reptiles as well, and is considered important in the decline of numerous rep-

tile species in the southeastern US (Mount 1981). Montgomery (1996) observed six cases of *S. invicta* attacking adult three-toed box turtles (*Terrapene carolina triunguis* (Agassiz)) in Texas, with only one of the turtles surviving due to human intervention. "It seems that the box turtle's defensive reaction is to withdraw the plastron, which cannot keep the ants out completely even when tightly closed. This reaction also renders the turtle immobile, which enables the ants to swarm over it, seeking out small openings between the plastron and the carapace. The turtle must eventually relax the plastron, which allows more ants to gain further access to its body." Montgomery (1996) concluded, "it would be difficult to estimate just how much effect the ants alone are having. But my observations lead me to believe that fire ants pose a serious threat to three-toed box turtles."

In addition to *S. invicta*, two other ant species found in the gopher tortoises' range also are known to attack vertebrates and may pose a threat to gopher tortoises: *Solenopsis geminata* (Fab.) and *Wasmannia auropunctata* (Roger). Although not generally as virulent as its congener *S. invicta*, *S. geminata* is also known to attack the hatchlings of birds and reptiles (e.g., Stoddard 1931; Travis 1941; Mrazek 1974). Where *W. auropunctata* occurs at high densities, it also can have a great impact on vertebrates. For example, in Gabon, *W. auropunctata* has been implicated in blinding house cats (*Felis catus* L.) and native wildlife, including elephants (*Loxodonta africana* (Blumenbach)) (Wetterer et al. 1999). In the Solomon Islands, locals reported that *W. auropunctata* commonly stings the eyes of dogs that eventually become blind, and attacks hatchlings of the ground-nesting Melanesian Scrubfowl (*Megapodius eremita* Hartlaub) (Wetterer 1997).

The present study was motivated, in part, by the discovery in a local greenway reserve of a Florida box turtle (*Terrapene carolina bauri* Taylor) being attacked by *S. invicta* (M. Floyd, pers. comm.). To evaluate the potential threat of *S. invicta* to gopher tortoises in the reserve, we surveyed ants at gopher tortoise burrows in the greenway.

MATERIALS AND METHODS

Study Area

Our study site was one section ("Range VIa") of a greenway reserve set up through the Abacoa residential development in Jupiter, Florida (26.90°N, 80.11°W), to preserve gopher tortoise habitat. The entire Abacoa greenway system encompasses roughly 10% of the development's land area. Our study segment included a 9.16-ha wooded range and an adjacent sunken water retention basin (Fig. 1). A path encircles the range



Figure 1. Aerial photograph of "Range VIa" in the Abacoa greenway, Jupiter, Florida, a wooded range bounded in the north by Frederick Small Road and on other sides by a sunken water catchment basin. An old fence line and a recent pipeline cross the northern and eastern parts of the range. The black bar equals approximately 30 m.

and two linear areas of disturbance cross through the northern and eastern portions of the range; one is an old (>6 years-old) cleared cattle fence line (fence removed) and the other is a recently (<3 years-old) dug pipeline (Fig. 1). A chain-link fence, separating the reserve from Frederick Small Road, bounds the north side of the range. The range is bounded on the other three sides by the water catchment basin with channels for directing and holding rainwater runoff from storms. Except for the channels, the basin is dry through most of the year. A chain-link fence separates the catchment basin from Central Road to the east and several baseball fields to the south and west.

The relatively undisturbed portions of the range consist of typical flatwood scrub (Myers and Ewel 1990) with a sparse canopy of mature slash pines (*Pinus elliottii* Engelman), an understory of saw palmetto (*Serenoa repens* (Bartram)) thickets and scrubby oaks (*Quercus* spp.), and open spaces dominated by wiregrass (*Aristida beyrichiana* Trin. & Rupr.), with lesser amounts of runner oak (*Quercus minima* Small) and deer moss lichens (*Cladina* spp. and *Cladonia* spp.). The old fence line is dominated by bunches of wiregrass and chalky bluestem (*Andropogon virginicus* L.) with small stands of young slash pine saplings and gallberry (*Ilex glabra* (L.)). The pipeline area is largely open sand with low grasses and herbs

growing in patches. The path around the outer edge of the wooded range is primarily covered with bahiagrass (*Paspalum notatum* Flueggé) that is mowed about every two to three weeks.

We surveyed ants at gopher tortoise burrows on 26 February, 14 March, and 16 April 2002 (sampling ~1/3 of the burrows on each day) using a folded index card with ~1 g of canned water-packed tuna inside placed within 0.2 m of the burrow entrance between 1300 and 1500 h. We surveyed ants using tuna bait to assess ant species present at the burrows that recruit heavily to rich animal protein resources. We returned 2 h (± 10 min) later and collected the cards, putting each in a separate zip-lock bag. We found that two h was enough time to allow significant recruitment, but not complete bait removal. After killing the collected ants in a freezer, we counted them, then preserved them in alcohol for identification. Stefan Cover (Museum of Comparative Zoology) and Mark Deyrup (Archbold Biological Station) identified the ants.

As of 16 April 2002, there were 85 marked gopher tortoises and 164 marked burrows within our study site. This density of 9.3 tortoises/ha is extremely high, e.g., it is more than three times higher than the highest mean density (2.7/ha) at Kennedy Space Center in east-central Florida (Breining et al. 1994). This high density is due,

in part, to people releasing gopher tortoises found elsewhere into the greenway (Diemer 1986).

From a map of burrow locations in the study area, we calculated the distance to the nearest edge of the wooded range for each burrow. We classified burrows less than 30 m from the perimeter of the wooded range as "edge" burrows; all others we classified as "interior" burrows.

RESULTS

We surveyed ants at 154 of the 164 marked gopher tortoise burrows; we excluded ten burrows because other animals removed the bait cards. All surveyed burrows had at least one ant present (range = 1-637 ants; median = 62 ants). We found 19 ant species (1-3 species per bait; Table 1); the most common was *S. invicta*, which occurred at 51 burrows (33%; Table 1). The only other ant found that may pose a threat to tortoises was the little fire ant, *Wasmannia auropunctata*, which occurred at four burrows (3%). *Solenopsis invicta* tended to recruit more workers to baits they occupied (median = 160 ants) than did other ant species (median = 40 ants).

We found *S. invicta* significantly more often at edge burrows (36/63 = 57%) than at interior burrows (15/91 = 16%; $\chi^2 = 27.8$; $P < 0.001$). In contrast, we found one or more species of native ant significantly less often at edge burrows (23/63 = 37%) than at interior burrows (62/91 = 68%; $\chi^2 =$

15.0; $P < 0.001$). We found one or more species of exotic ant other than *S. invicta* equally often at edge burrows (8/63 = 13%) as at interior burrows (18/91 = 20%; $\chi^2 = 1.3$; ns). We found only one species significantly less often at edge burrows than at interior burrows, the native *Crematogaster atkinsoni* Wheeler ($\chi^2 = 6.4$; $P < 0.05$); several ant species showed a strong trend in this direction, but we lacked sufficient sample size to demonstrate statistical significance.

Among the 91 interior burrows, we found *S. invicta* significantly more often at burrows directly on the narrow disturbed strips along the old fence line and pipeline (6/13 = 46%) than at interior burrows not on these strips (9/78 = 12%; $\chi^2 = 9.7$; $P < 0.005$). Sample size for edge burrow on and off these disturbed strips was too small for statistical comparison.

DISCUSSION

We found *S. invicta* present at most gopher tortoise burrows around the outer perimeter of the greenway reserve, but at few burrows in the interior portions of the greenway. Conversely, native ants were significantly less common on the outer perimeter than in the interior of the greenway. We also found that *S. invicta* was present more often at burrows on two disturbed strips through the interior of the greenway than at burrows in other parts of the greenway interior. These corridors of

TABLE 1. ANTS ON TUNA BAITS AT TORTOISE BURROWS ON THE GREENWAY EDGE ($n = 63$) AND THE GREENWAY INTERIOR ($n = 91$). ‡ = POTENTIAL THREAT TO GOPHER TORTOISES.

Exotic ant species	Occurrence at burrows	
	# edge	# interior
‡ <i>Solenopsis invicta</i> Buren	36	15
<i>Pheidole moerens</i> Wheeler	3	10
<i>Technomyrmex albipes</i> (Smith)	1	4
‡ <i>Wasmannia auropunctata</i> (Roger)	1	3
<i>Brachymyrmex obscurior</i> Forel	1	1
<i>Cardiocondyla emeryi</i> Forel	1	1
<i>Tapinoma melanocephalum</i> (Fabr.)	1	1
<i>Paratrechina longicornis</i> (Latr.)	0	1
Native ant species		
<i>Pheidole floridana</i> Emery	13	32
<i>Crematogaster atkinsoni</i> Wheeler	4	21
<i>Crematogaster ashmeadi</i> Mayr	0	6
<i>Forelius pruinosus</i> (Roger)	3	3
<i>Pheidole dentata</i> Mayr	3	3
<i>Dorymyrmex bureni</i> (Trager)	3	2
<i>Pheidole morrisii</i> Forel	2	2
<i>Aphaenogaster miamiana</i> Wheeler	1	1
<i>Temnothorax pergandei</i> (Emery)	1	1
<i>Aphaenogaster flemingi</i> Smith	1	0
<i>Dorymyrmex bossutus</i> (Trager)	0	1

disturbance thus allowed *S. invicta* more access to the interior of the greenway. Footpaths through greenway areas may have a similar influence. The greenway interior appears to offer tortoises and other species some refuge from *S. invicta*.

Our findings have conservation and development policy implications. For example, the long, thin design of many segments of the greenway may increase the exposure of gopher tortoises and other native species to *S. invicta* predation by increasing the spatial extent of edges (Saunders et al. 1991). Our study site was one of the wider segments in the Abacoa greenway. Narrower portions of the greenway may not have as much intact interior fauna. The division of the Abacoa greenway into sections separated by roads further increases the amount of edge. Because *S. invicta* prefers disturbed habitats (Tschinkel 1988; Stiles & Jones 1998), prescribed burning and reduction mowing, which land managers use to maintain open habitat for the gopher tortoises, could lead to increased *S. invicta* infestation and allow these ants to penetrate interior sites.

Stings of *S. invicta* may have serious detrimental effects on both adult and hatchling gopher tortoises. Gopher tortoise hatchlings sometimes remain in their nest one or more days after hatching (Butler & Hull 1996), during which time they may be particularly at risk to *S. invicta* attack. Hatching sea turtles, which typically take from several hours to several days after pipping before they emerge from their nests, are similarly vulnerable to ant attack. During this time, ants invade the nests and attack trapped hatchlings, particularly their sensitive eyes (Krahe et al. 2003). Because even a single fire ant sting may seriously impair a hatchling, ants may be having a tremendous impact on sea turtle hatchlings that are stung while emerging, through increases in subsequent mortality, e.g., increased vulnerability to other predators (Krahe et al. 2003).

Solenopsis invicta also poses a threat to other animals that use gopher tortoise burrows for shelter, including some species that are known only from gopher tortoise burrows, e.g., several species of beetle (including *Copris gopheri* Hubbard, *Onthophagus polyphemi* Hubbard, and *Aphodius troglodytes* Hubbard), a robberfly (*Machimus polyphemi* Bullington & Beck), the gopher frog (*Rana capito* LeConte), and the Florida mouse (*Podomys floridanus* (Chapman)) (Lago 1991; Lips 1991; Witz et al. 1991; Diemer 1992; Humphrey 1992; Deyrup & Franz 1995). Many of these animals are likely to be poorly defended against attacks by this exotic ant.

ACKNOWLEDGMENTS

We thank M. Deyrup and S. Cover for ant identification; M. Wetterer, A. Wetterer, and H. Smith for comments on this manuscript; and Florida Atlantic University for financial support.

REFERENCES CITED

- ALFORD, R. A. 1980. Population structure of *Gopherus polyphemus* in northern Florida. *J. Herpetol.* 14: 177-182.
- ALLEN, C. R., E. A. FORYS, K. G. RICE, AND D. P. WOJCIK. 2001. Effects of fire ants (Hymenoptera: Formicidae) on hatching turtles and prevalence of fire ants on sea turtle nesting beaches in Florida. *Florida Entomol.* 84: 250-253.
- ALLEN, C. R., E. A. FORYS, K. G. RICE, AND D. P. WOJCIK. 1998. Are red imported fire ants a threat to hatching sea turtles? pp. 113 *In* Proc. 17th Ann. Symp. Sea Turtle Biol. Conserv. NOAA Tech. Mem. NMFS-SEFSC-415.
- ALLEN, C. R., K. G. RICE, D. P. WOJCIK, AND H. F. PERCIVAL. 1997. Effect of red imported fire ant envenomization on neonatal American alligators. *J. Herpetol.* 31: 318-321.
- BREININGER, D. R., P. A. SCHMALZER, AND C. R. HINKLE. 1994. Gopher tortoise (*Gopherus polyphemus*) densities in coastal scrub and slash pine flatwoods in Florida. *J. Herpetol.* 28: 60-65.
- BUTLER, J. A., AND T. W. HULL. 1996. Reproduction of the tortoise, *Gopherus polyphemus*, in northeastern Florida. *J. Herpetol.* 30: 14-18.
- CINTRA, R. 1985. Nascimento de filhotes de *Caiman yacare* (Daudin, 1802) (Crocodylia: Alligatoridae) em condicoes semi-naturais no pantanal matogrossense. *Pap. Avul. Zool., Sao Paulo* 36(10): 91-101.
- DEYRUP, M., AND R. FRANZ. 1995. The Rare and Endangered Biota of Florida. IV. Invertebrates. University Press of Florida, Gainesville.
- DICKINSON, V. M. 1995. Red imported fire ant predation on Crested Caracara nestlings in south Texas. *Wilson Bull.* 107: 761-762.
- DIEMER, J. E. 1986. The ecology and management of the gopher tortoise in the southeastern United States. *Herpetologia* 42: 125-133.
- DOUGLASS, J. F. AND C. E. WINEGARNER. 1977. Predators of eggs and young of Gopher Tortoise, *Gopherus polyphemus* (Reptilia, Testudines, Testudinidae) in southern Florida. *J. Herpetol.* 11: 236-238.
- DIEMER, J. E. 1992. Gopher Tortoise, pp. 117-127 *In* P. E. Moler [ed.], The Rare and Endangered Biota of Florida. III. Amphibians and Reptiles, Univ. Press of Florida, Gainesville.
- DREES, B. M. 1994. Red imported fire ant predation on nestlings of colonial water birds. *Southwestern Entomol.* 19: 355-359.
- EPPERSON, D. M., AND C. D. HEISE. 2003. Nesting and hatchling ecology of gopher tortoises (*Gopherus polyphemus*) in Southern Mississippi. *J. Herpetol.* 37: 315-324.
- FLORIDA GAME AND FRESH WATER FISH COMMISSION. 1994. Official lists of endangered and potentially endangered fauna and flora in Florida. Florida Game & Fresh Water Fish Commission, Tallahassee.
- GIULIANO, W. M., C. R. ALLEN, R. S. LUTZ, AND S. DEMARAIS. 1996. Effects of red imported fire ants on Northern Bobwhite chicks. *J. Wildl. Manag.* 60: 309-313.
- HUMPHREY, S. R. 1992. The Rare and Endangered Biota of Florida. I. Mammals. Univ. Press of Florida, Gainesville, FL.
- KOPACHENA, J. G., A. J. BUCKLEY, AND G. A. POTTS. 2000. Effects of the red imported fire ant (*Solenopsis invicta*) on reproductive success of barn swallows

- (*Hirundo rustica*) in northeast Texas. *Southwestern Nat.* 45: 477-482.
- KRAHE, H., L. D. WOOD, AND J. K. WETTERER. 2003. Impact of predatory ants on post-emergence sea turtle hatchlings, pp. 211-212 *In Proc. 22nd Ann. Symp. Sea Turtle Biol. Conserv. NOAA Tech. Mem. NMFS-SEFSC 503.*
- LAGO, P. K. 1991. A survey of arthropods associated with gopher tortoise burrows in Mississippi. *Entomol. News* 102: 1-13.
- LANDERS, J. L., J. A. GARNER, AND W. A. MCRAE. 1980. Reproduction of gopher tortoises (*Gopherus polyphemus*) in southwestern Georgia. *Herpetol.* 36: 353-361.
- LIPS, K. R. 1991. Vertebrates associated with Tortoise (*Gopherus polyphemus*) burrows in four habitats in south central Florida. *J. Herpetol.* 25: 477-481.
- LOCKLEY, T. C. 1995. Effect of imported fire ant predation on a population of the least tern—an endangered species. *Southwestern Entomol.* 20: 517-519.
- LOVE, B. 1997. Fire ants. *Reptiles* 5(7): 86-89.
- MAIN, M. B., AND G. W. TANNER. 1999. Effects of fire on Florida's wildlife and wildlife habitat. University of Florida, Inst. Food Agric. Sci., Florida Coop. Ext. Serv., Doc. WEC 137.
- MONTGOMERY, W. B. 1996. Predation by the fire ant, *Solenopsis invicta*, on the three-toed box turtle, *Terrapene carolina triunguis*. *Bull. Chicago Herpetol. Soc.* 31: 105-106.
- MOULIS, R. A. 1996. Predation by the imported fire ant (*Solenopsis invicta*) on loggerhead sea turtle (*Caretta caretta*) nests on Wassaw National Wildlife Refuge, Georgia. *Chelon. Conserv. Biol.* 2: 433-436.
- MOUNT, R. H. 1981. The red imported fire ant, *Solenopsis invicta* (Hymenoptera: Formicidae), as a possible serious predator of some native southeastern vertebrates: direct observations and subjective impressions. *J. Alabama Acad. Sci.* 52: 71-78.
- MRAZEK, R. W. 1974. The relationship of the fire ant (*Solenopsis geminata* Fab) to nestlings of birds nesting on two spoil islands in the Laguna Madre. *Texas J. Sci.* 25: 140.
- MUELLER, J. M., C. B. DABBERT, S. DEMARAI, AND A. R. FORBES. 1999. Northern Bobwhite chick mortality caused by red imported fire ants. *J. Wildl. Manag.* 63: 1291-1298.
- MYERS, R. L., AND J. J. EWEL. 1990. Ecosystems of Florida. Univ. Press of Florida, Tallahassee, FL.
- PORTER, S. D., AND D. A. SAVIGNANO. 1990. Invasion of polygyne fire ants decimates native ants and disrupts the arthropod community. *Ecology* 71: 2095-2106.
- POWELL, J. 1995. Nesting ecology, diet, and the relationship of habitat and prey to site use by Mississippi barn owls (*Tyto alba*). Thesis. Mississippi State Univ., Mississippi State, MS.
- REAGAN, S. R., J. M. ERTEL, AND V. L. WRIGHT. 2000. David and Goliath retold: fire ants and alligators. *J. Herpetol.* 34: 475-478.
- RIDLEHUBER, K. T. 1982. Fire ant predation on Wood Duck ducklings and pipped eggs. *Southwestern Natur.* 27: 222.
- SAUNDERS, D. A., R. J. HOBBS, AND C. R. MARGULES. 1991. Biological consequences of ecosystem fragmentation. *Conserv. Biol.* 5: 18-32.
- SIKES, P. J., AND K. A. ARNOLD. 1986. Red imported fire ant (*Solenopsis invicta*) predation on Cliff Swallow (*Hirundo pyrrhonota*) nestlings in East-central Texas. *Southwestern Natur.* 31: 105-106.
- STEIGMAN, K. L. 1993. Nesting ecology of the Dickcissel (*Spiza americana*) on a tallgrass prairie relict in North Central Texas. Dissertation. Univ. North Texas, Denton.
- STILES, J. H. AND R. H. JONES. 1998. Distribution of the red imported fire ant, shape *Solenopsis invicta*, in road and powerline habitats. *Landsc. Ecol.* 13: 335-346.
- STODDARD, H. L. 1931. The bobwhite quail. Charles Scribner's Sons, New York, NY.
- TRAVIS, B. V. 1941. Notes on the biology of the fire ant *Solenopsis geminata* (F.) in Florida & Georgia. *Florida Entomol.* 24: 15-22.
- TSCHINKEL, W. R. 1988. Distribution of fire ants *Solenopsis invicta* and *S. geminata* in north Florida in relation to habitat and disturbance. *Ann. Entomol. Soc. Amer.* 81: 76-81.
- WADE, D. D. AND J. D. LUMSFORD. 1989. A guide for prescribed fire in southern forests. USDA, Forest Service Southern Region, Tech. Pub. R8-TP 11.
- WETTERER, J. K. 1997. Alien ants of the Pacific islands. *Aliens* 6: 3-4.
- WETTERER, J. K., P. D. WALSH, AND L. J. T. WHITE. 1999. *Wasmannia auropunctata* (Roger) (Hymenoptera: Formicidae), a destructive tramp ant, in wildlife refuges of Gabon. *Afr. Entomol.* 7: 292-294.
- WETTERER, J. K., AND L. D. WOOD. 2005. Distribution and impact of ants on a sea turtle nesting beach in Palm Beach County, Florida, pp. 351-353 *In Proc. 23rd Ann. Symp. Sea Turtle Biol. Conserv. NOAA Technical Memorandum NMFS-SEFSC 528.*
- WILMERS, T. J., E. S. WILMERS, M. MILLER, AND P. WELLS. 1996. Imported fire ants (*Solenopsis invicta*): a growing menace to sea turtle nests in Key West National Wildlife Refuge, pp. 341-343 *In Proc. 15th Ann. Symp. Sea Turtle Biol. Conserv.. NOAA Tech. Mem. NMFS-SEFSC-387.*
- WITZ, B. W., D. S. WILSON, AND M. D. PALMER. 1991. Distribution of *Gopherus polyphemus* and its vertebrate symbionts in 3 burrow categories. *Amer. Midl. Natur.* 126: 152-158.
- WOJCIK, D. P. 1994. Impact of the red imported fire ant on native ant species in Florida, pp. 269-281 *In D. F. Williams [ed.], Exotic Ants. Biology, Impact, and Control of Introduced Species.* Westview Press, Boulder, CO.