



**A New Genus *Pectinimura* (Lepidoptera, Gelechioidea, Lecithoceridae), with Four New Species from Thailand and the Philippines**

Authors: Park, Kyu-Tek, and Byun, Bong-Kyu

Source: Florida Entomologist, 91(1) : 110-115

Published By: Florida Entomological Society

URL: [https://doi.org/10.1653/0015-4040\(2008\)091\[0110:ANGPLG\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2008)091[0110:ANGPLG]2.0.CO;2)

---

BioOne Complete ([complete.BioOne.org](http://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A NEW GENUS *PECTINIMURA* (LEPIDOPTERA, GELECHIOIDEA, LECITHOCERIDAE), WITH FOUR NEW SPECIES FROM THAILAND AND THE PHILIPPINES

KYU-TEK PARK<sup>1</sup> AND BONG-KYU BYUN<sup>2</sup>

<sup>1</sup>McGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History, University of Florida, Gainesville, FL 32611 USA

<sup>2</sup>Division of Forestry Biodiversity, Korea National Arboretum, Pocheon, 487-821 Korea

ABSTRACT

A new genus *Pectinimura* Park (Lepidoptera, Lecithoceridae) is described, based on the type species, *Pectinimura montiatilis* sp. nov. Three additional species, *P. crassipalpis*, *P. batubatuensis*, and *P. crinalis* are described from Thailand or the Philippine Islands. A key to species of the genus is given, with illustrations of wing venation, and the male genitalia and female genitalia when available.

Key Words: Taxonomy, Lepidoptera, Lecithoceridae, *Pectinimura*, new genus, new species

RESUMEN

Se describe un nuevo género *Pectinimura* Park en la familia Lecithoceridae, basado sobre la especie tipo *Pectinimura montiatilis*, de Tailandia y las Islas Filipinas. Tres especies adicionales, *P. crassipalpis*, *P. batubatuensi* y *P. crinalis* son descritas e incluidas en este género de Tailandia o de las Islas Filipinas. Se incluye una clave de las especies de este género con ilustraciones de la genitalia de los machos.

The new genus *Pectinimura* (Lepidoptera: Lecithoceridae) is established, based upon the type species, *Pectinimura montiatilis* Park and Byun. *Pectinimura* contains four species described herein that are distributed in a region from Thailand to the Philippine Islands. The new genus is related to *Lecithocera* Herrich-Schäffer in the venation of both wings and the structure of the male genitalia. It is considered to be a monophyletic group having a very specialized comb-like plate on the valva of the male genitalia medially. Of these four new species, *P. crinalis* has some differences in the venation of both wings with the absence of  $R_2$  and  $M_2$  in the forewing and  $CuA_1$  and  $CuA_2$  coincident in the hindwing. However, there is no doubt that this species belongs to *Pectinimura* due to the similar structure of the comb-like plate in the valva of the male genitalia. No host plants are known for these species, however, most species of Lecithoceridae feed on debris and presumably *Pectinimura* may feed similarly.

MATERIALS AND METHODS

Descriptions are based on loaned material from the Zoological Museum, Copenhagen, collected by O. Karsholt and his colleagues in Philippines in 1961 and in Thailand in 1984. Along with

examination of external characters, all male and female genitalia of the species were dissected and examined. All types are deposited in the Zoological Museum, Copenhagen, Denmark. The color standard for the description of adults was based on "Methuen Handbook of Colour" by Kornerup and Wanscher (1978).

RESULTS

Genus *Pectinimura* Park, **gen. nov.**

The new genus *Pectinimura* is closely related to the genus *Lecithocera* Herrich-Schäffer in the external appearance, wing venation, and male genitalia. It can be separated from the latter by the shape of the labial palpus and male genitalia. Autapomorphies for the genus *Pectinimura* include the second segment of labial palpus thickened with rough scales or a long scale-tuft dorsally, and the male genitalia with deeply emarginate basal plate of uncus medially dividing into two lateral lobes caudally and the valva with a specialized comb-like plate medially.

Known distribution. Thailand and the Philippine Islands.

Etymology. The generic name is derived from Latin, *Pecten* (= comb) + *murus* (= wall).

KEY TO SPECIES OF THE GENUS *PECTINIMURA* PARK

1. Forewing with  $R_5$  and  $M_2$  present, without long hair-pencils on under surface . . . . . 2  
—Forewing with  $R_5$  and  $M_2$  absent, with a tuft of hair-pencils in a longitudinal furrow on under surface  
. . . . . *P. crinalis* **sp. nov.**
2. Second segment of labial palpus brownish, thickened with roughly erect scales above; mid tibia brownish ventrally. . . . . 3  
—Second segment of labial palpus golden yellow, thickened with fewer rough scales above; mid tibia pale orange ventrally. . . . . *P. batubatuensis* **sp. nov.**
3. Third segment of labial palpus about 1/2 length of 2<sup>nd</sup>; 2<sup>nd</sup> segment with long hair-like scales above; forewing with  $R_3$  approximated with  $R_{4+5}$  at base . . . . . *P. montiatilis* **sp. nov.**  
—Third segment of labial palpus as long as 2<sup>nd</sup>; 2<sup>nd</sup> segment with short erect scales above; forewing with  $R_3$  separated from  $R_{4+5}$  at base. . . . . *P. crassipalpis* **sp. nov.**

*Pectinimura montiatilis* Park and Byun, **new species**  
(Figs. 1, 6, 6a, 11)

Diagnosis. *Pectinimura montiatilis* is very similar to *P. crassipalpis* by sharing the characteristic comb-like plate in the valva of the male genitalia, but can be externally distinguished from the latter by the 2<sup>nd</sup> segment of labial palpus more roughly scaled with dark fuscous hair-like scale-tufts dorsally, and 3<sup>rd</sup> segment shorter, about 1/2 length of 2<sup>nd</sup>, and  $R_3$  in the forewing approximated with  $R_{4+5}$  at base.

Description. Male. Wingspan, 15.0-16.0 mm. Head mustard brown, with appressed scales dorsally. Scape of antenna slender, brownish above; flagellum orange white, with pale brownish flagellomeres, not ciliate ventrally. Second segment of labial palpus roughly scaled with dark fuscous hair-like scale-tufts above, brownish on outer surface, paler on inner surface; 3<sup>rd</sup> segment more or less stout, about 1/2 length of 2<sup>nd</sup>, dark fuscous beyond ventral half. Hind tibia brownish above. Tegula and thorax dark brown. Forewing mustard brown, speckled with dark fuscous scales sparsely throughout; costa slightly expanded anteriorly beyond 2/3; apex round; termen slightly oblique, but not concave or sinuate; fringe mustard brown with paler basal line;  $R_3$  approximated with  $R_{4+5}$  at base;  $R_4$  and  $R_5$  stalked near middle;  $R_6$  extending to termen;  $M_1$  far from  $R_{4+5}$  at base, nearly parallel to  $M_2$ ;  $CuA_1$  and  $CuA_2$  stalked at basal 1/5 of  $CuA_1$ . Hindwing grayish brown; apex more or less acute; fringe concolorous;  $M_3$  and  $CuA_1$  stalked basally.

Male genitalia (Figs. 6, 6a): Basal plate of uncus deeply incised at middle, with ovate lateral lobes; lateral margin slightly emarginate medially. Valva elongate, with round apex; costa with basal expansion anteriorly at basal one-fourth, angled on anterior corner, deeply concave; comb-like plate arising from middle of ventral margin above and extended to 3/5 length of valva; sacculus sclerotized, crescent-shaped, extending beyond middle of ventral margin; apex round. Juxta

subtriangular, about 1/3 length of vinculum. Aedeagus stout, slightly shorter than valva; cornutus weakly sclerotized, spatulate with a small, round subapical process. Seventh sternite convex on caudal margin, with long lateral arms, as long as the length of 8<sup>th</sup> segment.

Female. Unknown.

Holotype: Male, Philippines, Palawan, Manta-lingajan, Pinigisan 600 m, 24 IX 1961, Noona dan Exp. 61-62, gen. prep. no. CIS-5448/Park. Paratypes: 6 males, same locality as the holotype, 10, 12 & 13 IX 1961, gen. prep. no. CIS-5455/Park; 3 males, same locality, 19, 23 & 24 IX 1961, gen. prep. No. CIS-5489/Park.


Distribution. Philippine Islands (Palawan).

Etymology. The species name is derived from Latin, *montanus*, meaning mountain.

*Pectinimura crassipalpis* Park and Byun, **new species**  
(Figs. 2, 7, 7a)

Diagnosis. This new species is externally similar to *Pectinimura montiatilis*, but is differentiated from the latter by the 2<sup>nd</sup> labial palpus with fewer short scales dorsally, and 3<sup>rd</sup> segment longer.

Description. Wingspan, 17.0 mm. Head, tegula, and thorax mustard brown. Scape of antenna slender, dark brown above. Second segment of labial palpus thickened, roughly scaled dorsally, brownish yellow on outer surface speckled with dark-brown scales, and orange white on inner surface; 3<sup>rd</sup> segment nearly as long as 2<sup>nd</sup>. Hind tibia brownish with dense setae dorsally and ventrally. Forewing mustard brown; a dark fuscous discal spot at upper corner of cell, distinct or inconspicuous; costa nearly straight; apex round; termen slightly oblique, not sinuate; fringe yellowish brown, with narrow pale gray basal line;  $R_3$  divergent from  $R_{4+5}$  at base;  $R_4$  and  $R_5$  stalked near 3/5;  $R_6$  extending to or below the wing apex;  $M_1$ ,  $M_2$  and  $M_3$  near parallel;  $CuA_1$  and  $CuA_2$  stalked at basal one-fourth. Hindwing grayish brown; apex rather obtuse;  $R_s$  and  $M_1$  divergent from beyond cell;  $M_3$



Figs. 1-5. Adults, labial palpus, forewing venation of *Pectinimura* spp. (1) *P. montiatilis*, (2) *P. crassipalpis*, (3) *P. batubatuensis*, (4) *P. crinalis*, (5) Lateral aspect of head with labial palpus: (a) *P. montiatilis*, (b) *crassipalpis*, (c) *P. batubatuensis*, (d) *P. crinalis*.

and  $CuA_1$  stalked at basal  $2/5$ ; fringe dark brown, with pale orange gray basal line.

Male genitalia (Figs. 7, 7a): Basal plate of uncus deeply emarginate at middle, divided into two triangular lateral lobes, slightly concave on lateral margin. Valva with costa deeply concave medially; comb-like plate arising from middle of ventral margin, extended to about  $3/5$  length of valva towards termen; saccus heavily sclerotized, extended to  $2/3$  of ventral margin; apex round. Juxta subtriangular, about  $1/2$  length of vinculum; caudal margin slightly concave. Aedeagus stout, slightly shorter than valva; cornutus spatulate, weakly sclerotized, truncate on caudal margin, with a small subapical protrusion; and with a short, broad plate medially and a large patch of spinules internally.

Female. Unknown.

Holotype: Male, Loei, Phu Luang Wildlife Sanc. 8-14 X 1984, 700-900 m (Karsholt, Lam-

holdt, & Nielsen), gen. prep. no. CIS-4773/Park. Paratype: 1 male, same data as the holotype, gen. prep. no. CIS-5500/Park.


Distribution. Thailand (Loei Province).

Etymology. The species name is derived from Latin, *crassus*, meaning thick, and referring to abnormally thickened 2<sup>nd</sup> segment of the labial palpus.

*Pectinimura batubatuensis* Park and Byun, **new species** (Figs. 3, 8, 8a, 10, 12)

Diagnosis. *Pectinimura batubatuensis* differs from the 2 preceding species by the normally thickened, golden yellow 2<sup>nd</sup> segment of labial palpus, with less rough scales above.


Description. Male and female. Wingspan, 15.0-16.0 mm. Head, tegula, and thorax dark brown. Antenna with dark brown scape dorsally; flagellum yellowish white with pale brownish flagello-



Figs. 6-9. Male genitalia of *Pectinimura* spp: (a) Magnification of the basal plate of uncus, (b) Aedeagus, (6) *P. montiatilis*, (7) *P. crassipalpis*, (8) *P. batubatuensis*, (9) *P. crinalis* (arrows indicate the comb-like pectinate plate). Scale bar: 1 mm.

meres. Second segment of labial palpus somewhat slender, normally thickened, smooth scaled above, golden yellow outer surface, paler on inner surface; 3<sup>rd</sup> segment slender, slightly shorter than 2<sup>nd</sup>, dark fuscous ventrally. Mid tibia grayish orange ventrally; hind tibia dark brown on ventral surface. Forewing dark brown; apex obtuse; termen

oblique, slightly sinuate; fringe dark brown with pale brownish gray basal line;  $R_{4+5}$  divergent from  $R_4$  and  $R_5$  stalked before middle;  $R_5$  extending to termen;  $M_1$  far from  $R_3$  at base, nearly parallel to  $M_2$ ;  $CuA_1$  and  $CuA_2$  stalked at basal 1/5 of  $CuA_1$ ; Hindwing dark brown, with silvery white scales between costa and Sc vein; apex more or



Figs. 10-13. Female genitalia and wing venations of *Pectinimura* spp. (10) Female genitalia of *P. batubatuensis*, (10a) Magnification of signum, (11) Forewing venation of *P. montiatilis* sp. nov., (12) Forewing venation of *P. batubatuensis*, (13) Forewing venation of *P. crinalis*. Scale bar: 1 mm.

less acute; termen oblique; fringe brown;  $M_3$  and  $CuA_1$  stalked before middle.

Male genitalia (Figs. 8, 8a): Basal plate of uncus roundly convex on caudal margin. Valva slender; costa concave medially, then almost straight; a comb-like plate, nearly extending to costa. Juxta subtriangular, about 1/2 length of vinculum, with nearly straight caudal margin. Aedeagus slender, narrower than width of distal part of valva, with a narrow plate.

Female genitalia (Fig. 10): Eighth sternite weakly sclerotized, slightly emarginate on caudal margin medially. Antrum cup-shaped, about 2/5 length of ductus bursae; ductus bursae narrow; ductus seminalis arising from conjunction of corpus bursae. Corpus bursae ovate; signum elliptical, and denticulate on surface.

Holotype: Male, Philippine, Tawi Tawi, Tarawakan, north of Batu Batu, 16 X 1961, Noona Dan Exp. 61-62, gen. prep. No. CIS-5493/Park. Paratype: 11 males and females, same locality, 13-16 IX 1961, gen. prep. No. CIS-5452/Park (male); -5454/Park (female); 25 males and females, same locality, 22-30 X 1961, gen. prep. no. CIS-5453/Park; 12 males and females, same locality, 7 XI 1961.

Distribution. Philippine Islands (Batu Batu).

**Etymology.** The species name is derived from the type locality.

*Pectinimura crinalis* Park and Byun, **new species**  
(Figs. 4, 9, 9a, 13)

**Diagnosis.** *Pectinimura crinalis* differs from the 3 preceding species by the forewing with  $R_3$  and  $M_2$  absent, with specialized long hair-pencils in a longitudinal furrow in the cell and cubital pectins on under surface, hindwing with  $CuA_2$  absent, and 3<sup>rd</sup> segment about 2/3 as long as 2<sup>nd</sup>.

**Description.** Male. Wingspan, 14.5-15.5 mm. Head yellowish brown, with more or less erect scales laterally. Antenna with slender, yellowish white scape; flagellum yellowish white with pale brown flagellomeres. Second segment less roughly scaled than *P. montiatilis* or *P. crassipalpis*, light brown on outer surface, speckled with dark-brown scales; 3<sup>rd</sup> segment of labial palpus about 2/3 length of 2<sup>nd</sup>. Mid tibia creamy white on ventral surface; Hind tibia black. Forewing yellowish brown, speckled with dark-fuscous scales; with a tuft of long hair-pencils in a furrow along central line from basal 1/3 to end of cell; with cubital pectin on under surface;  $R_2$  absent;  $R_4$  and  $R_5$

stalked near middle;  $R_5$  extending to or below the wing apex;  $M_2$  absent;  $CuA_1$  and  $CuA_2$  stalked near 1/5. Hindwing grayish brown; cell closed with strong cross veins;  $M_2$  and  $M_3$  close at base;  $CuA_1$  and  $CuA_2$  absent.

Male genitalia (Figs. 9, 9a): Basal plate of uncus small, with some irregular caudal margin. Valva short, roundly expanded in distal part; costa concave medially; ventral margin emarginate at middle; comb-like plate well-developed. Juxta subtriangular, about 3/5 length of vinculum. Aedeagus slender, narrower than 1/2 width of valva, with 2 subtriangular process preapically and a weakly sclerotized narrow plate before basal half.

Female. Unknown.

Holotype: Male, Philippine, Palawan, Manta-lingajan, Pinigisan 600 m, 24 IX 1961, Noona Dan Exp. 61-62, gen prep. no. CIS-5450/Park. Paratypes: 4 males, same locality as the holotype, gen prep. no. CIS-5451/Park; 3 males, same locality, 9-11 IX 1961; same locality, gen. prep. no. CIS-5487/Park; 2 males, same locality, 13 IX 1961, gen. prep. no. CIS-5488/Park.

Distribution. Philippine Islands (Palawan).

Etymology. The species name is derived from Latin, *crini*, meaning hair.

Remarks. Although this species has different venation than the preceding species, it shares the specialized genital structure of a comb-like plate in the valva, and is considered to be congeneric with them.

#### DISCUSSION

*Pectinimura* Park, **n. gen.** has a very specialized comb-like plate in the valva of the male genitalia, which is not known in any other genera, and is considered to be an autapomorphic character. The external appearance and the venation of both wings of the species are similar to those of

the genus *Lecithocera*, but  $R_5$  in the forewing of these species is not stalked with  $R_{4+5}$ . Of the 4 described species, the first 3 new species have similar wing venations, but *P. crinalis* differs from them in wing venation as noted in the introduction. However, there is no doubt that *P. crinalis* belongs to *Pectinimura* due to the similar structure of the male genitalia with a specialized comb-like plate on the valva. Because some infraspecific variations of the venation within certain genera of the family Lecithoceridae were previously noted by Park (1999), emphasizing that the venation can not always be a main reliable character for a genus in the family, with some different opinions from previous workers (Gozmány 1978; Wu 1997).

#### ACKNOWLEDGMENTS

We are grateful to O. Karsholt, Zoological Museum, Copenhagen, Denmark, for the loan of valuable specimens for this study; M. Thomas, Florida State Collection of Arthropods, Division of Plant Industry, Gainesville, FL, and D. Matthews Lott, McGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History, Gainesville, FL, for helpful comments of the manuscript, and Eon-Mi Ji, Kangwon National University, Korea, for assistance in the preparation of plates.

#### REFERENCES CITED

- GOZMÁNY, L. 1978. Lecithoceridae, In H. G. Amsel, F. Gregor, and H. Reisser [eds.], *Microlepidoptera Palaearctica*. 5. Georg Fromme & Co., Wien. 306 pp.
- KORNERUP, A., AND J. H. WANSCHER. 1978. *Methuen Handbook of Colour*, 3rd ed. Methuen, London. 252 pp.
- PARK, K. T. 1999. Lecithoceridae of Taiwan (I): Subfamily Lecithocerinae: Genera *Homaloxestis* Meyrick and *Lecithocera* Herrich-Schäffer. *Zoological Studies* 38(2): 238-256.
- WU, C. 1997. *Lepidoptera Lecithoceridae*. Fauna Sinica, Insecta. Vol. 7. Science Press, Beijing. 302 pp.