



Establishment of *Amblyomma maculatum* (Gulf Coast Tick) in Arkansas, U.S.A.

Authors: Trout, R. T., Steelman, C. D., Szalanski, A. L., and Loftin, K.

Source: Florida Entomologist, 93(1) : 120-122

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.093.0117>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ESTABLISHMENT OF *AMBLYOMMA MACULATUM* (GULF COAST TICK) IN ARKANSAS, U.S.A.

R. T. TROUT, C. D. STEELMAN, A. L. SZALANSKI AND K. LOFTIN
AGRI 319 Entomology, University of Arkansas, Fayetteville Arkansas

Amblyomma maculatum Koch (Gulf Coast tick) has been reported throughout the southeastern United States, California, and Arizona (Goddard & Norment 1983) and on various animals including cattle, deer, mesocarnivores, and ground dwelling birds (Tugwell & Lancaster 1962). The tick is a pest of economic and pathogenic concern since it readily infests cattle (Semtner & Hair 1973), and is the vector of *Cowdria ruminantium* Cowdry (the causative agent of Heartwater disease) (Uilenburg 1982) and *Rickettsia parkeri* (the causative agent of newly described spotted fever) (Paddock et al. 2004). Previous confirmations of *R. parkeri* in *A. maculatum* included specimens from Georgia, Florida, Kentucky, Mississippi, and South Carolina (Sumner et al. 2007). *Amblyomma maculatum* was initially thought to reside within 160 km of the Atlantic and Gulf Coasts (Bishop & Hixson 1936), but it has been found in 18 eastern counties of Oklahoma (Semtner & Hair 1973) and in Washington County and Ashley County in northwest Arkansas (Lancaster 1973). It may have become redistributed with livestock movement (Goddard & Norment 1983) and cattle egrets migration (USDA APHIS 2008).

As a part of a larger study, investigating ticks on canines and deer (Trout & Steelman *in press*), veterinarians and the Arkansas Game and Fish Commission removed ticks from hosts and stored them in 100% ethanol. Collection date, location, and host were recorded, and collections were submitted to the Veterinary Entomology laboratory at the University of Arkansas where they were identified to species, sex, life stage, and engorgement with keys from Arthur (1961), Lancaster (1973), Goddard & Norment (1985).

To confirm the species identity, 9 identified *A. maculatum* were subjected to phylogenetic analyses based on DNA sequences of a portion of the mitochondrial 16S mt rDNA gene. Extraction and PCR amplification procedures were similar to Trout et al. (2009). Briefly, individual specimens were cut longitudinally with a sterile razor blade and subjected to the Qiagen Dneasy Insect Protocol (Qiagen Inc. Rohm and Haas Company, Valencia, CA). The DNA extractions were assessed by PCR with mitochondrial primers 16S+2 and 16S-1 per the methods previously described (Black & Piesman 1994). Reaction products were analyzed with gel electrophoresis. Positive reaction products were purified and concentrated with minicolumns according to the manufacturer's instructions (Wizard PCR preps, Promega) and sent to the University of Arkansas Medical Sciences

DNA Sequencing Facility (Little Rock, AR) for direct sequencing in both directions. Sequences were aligned with Clustal in Bioedit 5.09 (Hall 1999). Bayesian Evolutionary Analysis Sampling Trees (BEAST) version 1.4.2 software (Drummond & Rambaut 2003) was used to obtain phylogenetic trees with the GTR + G model. To create the consensus tree, 4 Markov chains ran for 10⁶ generations with a burn-in of 2 x 10⁴. Phylogenetic relationships were compared to other *Amblyomma* ticks and 1 *Argas* tick obtained from GenBank, as follows: *Amblyomma americanum* Say (L34314), *A. americanum* (L34313), *A. aureolatum* (Pallas) (AF541254), *A. cajennense* Fabricius (L34317), *A. dubitatum* Neumann (DQ858954), *A. glauerti* Keirans, King & Sharrad (U95853), *A. hebraeum* Koch (L34316), *A. maculatum* Koch (AY375442), *A. ovale* Koch (AF541255), *A. parvum* Arag (EU306154), *A. triste* Koch (AY498563), *A. tuberculatum* Marx (U95856), *A. variegatum* Fabricius (L34315), and *Argas brevipes* Banks (U95863). All 9 *A. maculatum* specimens were confirmed with Bayesian analysis and GenBank comparisons. Seven haplotypes of the 16S mt rDNA were identified which were 100% homologous with NCBI GenBank accession numbers DQ076315-DQ76321 collected from Kansas, Texas, and Oklahoma (Ketchum et al. 2009). All of the haplotypes were placed into 1 clade suggesting that the group is monophyletic.

A total of 232 *A. maculatum* were collected in Arkansas between 2006 and 2009 (Table 1, Fig. 1). We identified *A. maculatum* collected from 22 canines (*Canis lupus familiaris* L.), 50 white-tailed deer (*Odocoileus virginianus* (Boddaert)), 1 feline (*Felis catus* L.), and 1 cow (*Bos taurus* L.) from Arkansas between I-III-2006 and I-VI-2009. The collection sites were then georeferenced in ArcMap 9.3.1 (ESRI Redlands CA) to map the distribution. Previous reports of *A. maculatum* included a specimen in a 1950s drag sample from Washington County and in a 1960s cattle sample from Ashley County (Lancaster 1973). Eads (2001) found 13 Gulf Coast ticks between May and Jul of 2000 at the Arkansas National Post, located in Arkansas County in the southeast region of the state. We believe this tick's distribution across the state suggests the tick has established itself compared to the few numbers and limited distribution mentioned by Koch (1982). The calculated Moran's I for *A. maculatum* on the canines was 0.01 (0.47 standard deviation) and 0.46 (1.19 standard deviation) on deer, indicating the patterns were neither clustered nor dispersed, but random across

TABLE 1. IDENTIFIED *AMBLIOMMA MACULATUM* TICKS COLLECTED FROM ARKANSAS MAMMALS.

Host	No. Hosts Infested/ No. Hosts Sampled (% infestation)	Life Stage				Total (e)
		Larva (e ¹)	Nymph (e)	Male (e)	Female ² (e)	
Canine	23/156 (15%)	0	8 (2)	95 (2)	9 (0)	113 (4)
White-tailed deer	50/250 (20%)	0	8 (0)	46 (10)	41 (27)	95 (37)
Cow	1/1 (100%)	0	0	12 (8)	11 (5)	23 (13)
Feline	1/16 (6%)	0	0	0	1 (1)	1 (1)
Total	74/422 (17.5%)	0	16 (2)	153 (20)	62 (33)	232 (55)

¹Number of engorged specimens.

²One adult specimen was damaged and could not be properly sexed.

Distribution of Amblyomma maculatum




Fig. 1. Geographical distribution of *Amblyomma maculatum* from canines (triangles), white-tailed deer (circles), feline (square), and cow (pentagon) in Arkansas. The larger the symbol the greater the number of ticks found on that animal.

the state. Previous reports considered this tick's distribution as "sporadic" since they are dispersed with livestock shipments (Goddard & Paddock 2005; Goddard 2007).

SUMMARY

The Gulf Coast tick was found throughout Arkansas on wild and domesticated animals; conse-

quently, this tick has established itself in Arkansas. Bayesian analyses confirmed the presence of this species. Because *Amblyomma maculatum* is the primary vector of *Rickettsia parkeri* and the secondary vector of *Cowdria ruminantium*, additional studies on the distribution of this tick should be investigated as well as its relation to human diseases.

REFERENCES CITED

ARTHUR, D. R. 1961. Ticks and Disease. Row, Peterson and Company. Evanston, Ill.

BISHOP, F. C., AND HIXSON, H. 1936. Biology and economic importance of the Gulf Coast tick. J. Econ. Entomol. 29: 1068-1076.

BLACK, W. C., AND PIESMAN, J. 1994. Phylogeny of hard- and soft- tick taxa (Acari: Ixodida) based on mitochondrial 16S rDNA sequences. Proc. Nat. Acad. Sci. 91: 10034-10038.

DRUMMOND, A. J., AND RAMBAUT, A. 2003. BEAST v1.0 (<http://evolve.zoo.ox.ac.uk/beast/>).

EADS, G. K. 2001. An Assessment of Tick Density and Tick-borne Disease Frequency at Arkansas Post National Memorial. [Thesis] University of Arkansas-Monticello.

GODDARD, J., AND NORMENT, B. R. 1983. Notes on the geographical distribution of the Gulf Coast tick, *Amblyomma maculatum* (Koch) [Acari: Ixodidae]. Entomol. News. 94: 103-104.

GODDARD, J., AND NORMENT, B. R. 1985. A guide to the ticks of Mississippi. Mississippi State University, MS Agricultural and Forestry Exp. Sta. Bull. No. 935.

GODDARD, J., AND PADDOCK, C. D. 2005. Observations on distribution and seasonal activity of the Gulf Coast Tick in Mississippi. J. Med. Entomol. 42: 176-179.

GODDARD, J. 2007. Seasonal activity of *Amblyomma* spp. in Mississippi. J. Vector Ecol. 32: 157-158.

HALL, T. A. 1999. BioEdit: a user-friendly biological sequence alignment and analysis program for Windows 95/98/NT. Nucleic Acids Symp. Ser. 41: 95-98.

KETCHUM, H. R., TEEL, P. D., COATES, C. J., STREY, O. F., AND LONGNECKER, M. T. 2009. Genetic variation in 12S and 16S mitochondrial rDNA genes of four geographically isolated populations of Gulf Coast ticks (Acari: Ixodidae). J. Med. Entomol. 46: 482-489

- KOCH, H. G. 1982. Seasonal incidence and attachment sites of ticks (Acari: Ixodidae) on domestic dogs in southeastern Oklahoma and northwestern Arkansas. *J. Med. Entomol.* 19: 293-298.
- LANCASTER, J. L. 1973. A Guide to Ticks of Arkansas. Agricultural Experiment Station. Division of Agriculture. University of Arkansas, Fayetteville. Bulletin 779.
- PADDOCK, C. D., SUMNER, J. W., COMER, J. A., ZAKI, S. R., GOLDSMITH, C. S., GODDARD, J., MCLELLAN, S. L. F., TAMMINGA, C. L., AND OHL, C. A. 2004. *Rickettsia parkeri*: a newly recognized cause of spotted fever Rickettsiosis in the United States. *Clin. Infect. Dis.* 38: 805-11.
- SEMTNER, P. J., AND HAIR, J. A. 1973. Distribution, seasonal abundance, and hosts of the Gulf Coast tick in Oklahoma. *Ann. Entomol. Soc. Amer.* 66: 1264-1268.
- SUMNER, J. W., DURDEN, L. A., GODDARD, J., STROMDAHL, E. Y., CLARK, K. L., REEVES, W. K., AND PADDOCK, C. D. 2007. Gulf Coast ticks (*Amblyomma maculatum*) and *Rickettsia parkeri*, United States. *Emerg. Infect. Dis.* 13: 751-753.
- TROUT, R. T., AND STEELMAN, C. D. 2009. Ticks parasitizing canines and deer in Arkansas. *J. Entomol. Sci.* *In Press.*
- TROUT, R. T., STEELMAN, C. D., AND SZALANSKI, A. L. 2009. Population genetics and phylogeography of *Ixodes scapularis* Say (Acari: Ixodidae) collected from Arkansas canines and deer. *Southwest. Entomol.* 34 (3): 273-287.
- TUGWELL, P., AND LANCASTER, JR., J. L. 1962. Results of a tick-host study in Northwest Arkansas. *J. Kansas Entomol. Soc.* 35: 202-211.
- UILENBERG, G. 1982. Experimental transmission of *Cowdria ruminantium* by the Gulf Coast tick *Amblyomma maculatum*: Danger of introducing heartwater and benign African theileriasis onto the American mainland. *American J. Vet. Res.* 43: 1279-1282.
- USDA APHIS (UNITED STATES DEPARTMENT OF AGRICULTURE ANIMAL AND PLANT HEALTH INSPECTION SERVICE). 2008. Assessment of change in risk for release of *Ehrlichia ruminantium* (Heartwater) into the continental United States following discontinuation of the Tropical Bont Tick (*Amblyomma variegatum*) eradication program on St. Croix, U.S. Virgin Islands. Centers for Epidemiology and Animal Health. Ft. Collins CO.