



New Records on the Geographical Distribution of South American Sharpshooters (Cicadellidae: Cicadellinae: Proconiini) and their Potential as Vectors of *Xylella fastidiosa*

Authors: Dellapé, Gimena, Logarzo, Guillermo A., Virla, Eduardo G., and Paradell, Susana L.

Source: Florida Entomologist, 94(2) : 364-366

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.094.0239>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

NEW RECORDS ON THE GEOGRAPHICAL DISTRIBUTION OF SOUTH AMERICAN SHARPSHOOTERS (CICADELLIDAE: CICADELLINAE: PROCONIINI) AND THEIR POTENTIAL AS VECTORS OF *XYLELLA FASTIDIOSA*

GIMENA DELLAPÉ¹, GUILLERMO A. LOGARZO², EDUARDO G. VIRLA³ AND SUSANA L. PARADELL¹
¹División Entomología, Facultad de Ciencias Naturales y Museo, UNLP, Paseo del Bosque s/n^o
 (B1900FWA), La Plata, Buenos Aires, Argentina
 E-mail: gimenadellape@fcnym.unlp.edu.ar

²USDA-ARS South American Biological Control Laboratory, Bolivar 1559 (1686), Hurlingham,
 Buenos Aires, Argentina

³PROIMI-Biotecnología, División de Control Biológico, Av. Belgrano y Pje Caseros, (4000),
 San Miguel de Tucumán, Tucumán, Argentina

The Proconiini comprises 422 species distributed in the continental Americas, the highest biodiversity is found in the Neotropical Region (Wilson et al. 2009). Members of the tribe Proconiini have been identified as vectors of many diseases caused by the bacteria *Xylella fastidiosa* Wells et al. 1978, which occurs only in the xylem of plants (Marucci et al. 2002).

Strains of *X. fastidiosa* cause diseases such as "Pierce's Disease" (PD) in grape (*Vitis vinifera* L.), "Phony Peach Disease" (PPD), "Coffee Leaf Scorch" (CLS), "Oleander Leaf Scorch" (OLS), and "Citrus Variegated Chlorosis" (CVC) among others. These incurable maladies produce substantial economic losses in a diverse variety of crops (Hernandez-Martinez et al. 2006).

In South America the major threat is CVC which has spread rapidly throughout Brazil (Lopes 1996). *X. fastidiosa* is also present in United States, México, Venezuela, Brazil, Paraguay, Uruguay, Argentina (Redak et al. 2004), and Costa Rica (Aguilar et al. 2005). However, CVC is not yet reported from the USA although it has the potential to threaten orange (*Citrus × sinensis* (L.) Osbeck) production in the Americas if a suitable vector is available (Damsteegt et al. 2006).

Diseases caused by *X. fastidiosa* have attained great importance worldwide as insect vectors of this pathogen have demonstrated an ability to spread, as happened with *Homalodisca vitripennis* (Germar), which invaded many islands in the Pacific Ocean (Pilkington et al. 2005). Pathogen acquisition and transmission by sharpshooters occurs because these insects feed exclusively on xylem fluids (Young 1968).

Despite this obvious importance, there are few studies from South America that have identified Proconiini species that can transmit *X. fastidiosa*. Moreover, there is no basic information on biology, geographic distributions, phenology, natural enemies or host plant associations for many South American Proconiini species. To address this shortcoming, work presented here provides new distributional records for thirteen South Ameri-

can Proconiini sharpshooters that may be potential vectors of *X. fastidiosa*.

The examined material is deposited in the following entomological collections of Argentina: Instituto Miguel Lillo (IMLA); Museo de Ciencias Naturales de La Plata (MLP) and Museo Argentino de Ciencias Naturales 'Bernardino Rivadavia' (MACN). Specific identification and distributional data were compiled from Young (1968), Marucci et al. (2002) and Wilson et al. (2009).

Examined Material

Acrogonia citrina Marucci & Cavichioli, PARAGUAY: Carumbé, III-1965, Golbach Leg., 1♂, 1♀, without information on host plant and collecting method (IMLA).

Acrogonia flavoscutellata (Signoret), ECUADOR: Santo Domingo, IV-1958, Weyrauch Leg., 1♀, without information on host plant and collecting method (IMLA).

Dechacona missionum (Berg), URUGUAY: 1♂, without date, no information on host plant and collecting method (MACN).

Diestostemma huallagana Young, BOLIVIA: 1♂ without date and locality (MACN).

Molomea consolidata Schröder, ECUADOR: El Puyo, IV-1958, 1♂. PERÚ: Tingo María, 1♂; Chanchamayo, II-1939, 1♂ (IMLA). BOLIVIA: Santa Cruz, I-1958, Wygodzinsky Leg.; 1♂ (IMLA), 1♀ (MACN), without information on host plant and collecting method.

Molomea personata (Signoret), PERÚ: Fundo Génova, IV-2002, Logarzo-Varone Legs., 8♀♀, on papaw; V-2002, Logarzo Leg., 2♀♀, on grasses, by sweeping (IMLA).

Ochrostacta diadema (Burmeister), PARAGUAY: Itagua, I-1957, Montes Leg., 3♀♀, without information on host plant and collecting method (MLP).

Oncometopia rubescens Fowler, PERÚ: Fundo Génova, IV-2002, Logarzo Leg., 14♂♂, 16♀♀, on grasses and papaw; V-2002, Logarzo Leg., 16♂♂,

4♀♀, by sweeping (IMLA). PARAGUAY: Carumbé, San Pedro, I-1971, Golbach Leg., 1♀, without collecting method (IMLA).

Proconia fusca Melichar, BOLIVIA: 9♂♂, 4♀♀, without information on host plant and collecting method (MACN).

Tapajosa doeringi (Berg), PERÚ: Cuzco, Machu Picchu, II-1952, Monrós Leg., 1♂, without information on host plant and collecting method (IMLA).

Tapajosa rubromarginata (Signoret), PARAGUAY: Caaguazú, XII-2000, Logarzo Leg., 1♂, on weeds (MLP).

Tapajosa similis (Melichar), BRAZIL: I-1948, Cuezco Leg., 1♂, without information on host plant and collecting method (IMLA).

Tretogonia callifera Melichar, PARAGUAY: Carumbé, 3♂♂, 2♀♀; Caaguazú, I-1965, 2♂♂, 3♀♀, without information on host plant and collecting method (IMLA).

In Table 1, we summarize the species recorded for Central and South America, the presence of *X. fastidiosa* and the new distribution data for the species listed here. Sharpshooters are known to occur in 24 of the 37 Central and S. America countries. No data are available for some islands associated with Central America and the Caribbean. This lack of information about the Proconiini in

these countries is probably due to a deficiency in surveys and collections and not because of the absence of representatives of these insects in those territories.

Most studies investigating the transmission of *Xylella* have been conducted in the USA. In the Neotropics, the majority of studies have been made in Brazil (Redak et al. 2004). Some South America countries, such as Perú, Bolivia, Colombia, and Ecuador have more than 50 sharpshooter species capable of vectoring *X. fastidiosa*, but no reference to occurrence of this bacterium in those countries is available.

Most South American countries are at risk from *X. fastidiosa* because the bacterium has a wide host range and may be transported accidentally to new areas via infected plant species. There are strong epidemiological relationships between the presence of Proconiini sharpshooters and incidence of the bacterium. Resulting diseases can take months or years to develop significant symptoms, or infections may remain asymptomatic and undetected while acting as reservoirs from which continued bacterial transmission can occur (Hopkins 1989).

We thank the museum curators who provided access to the specimens used from their entomological collections.

TABLE 1. KNOWN AND NEW DISTRIBUTION RECORDS FOR 13 SPECIES OF PROCONIINI COLLECTED IN CENTRAL AND SOUTH AMERICA.

	<i>Acrogonia citrina</i>	<i>Acrogonia flavoscutellata</i>	<i>Dechaona missionum</i>	<i>Diestostemma huallagana</i>	<i>Molomea consolidata</i>	<i>Molomea personata</i>	<i>Ochrostacta diadema</i>	<i>Oncometopia rubescens</i>	<i>Proconia fusca</i>	<i>Tapajosa doeringi</i>	<i>Tapajosa rubromarginata</i>	<i>Tapajosa similis</i>	<i>Tretogonia callifera</i>	Presence of <i>X. fastidiosa</i>	Total specimens
Argentina			*		*					*	*	*	*	*	38
Bolivia			*	X	X				X				*	*	58
Brazil	*	*	*	*	*	*	*	*			*	X	*	*	142
British Guiana		*													18
Colombia		*						*					*	*	65
Costa Rica							*	*						*	49
Ecuador		X			X		*	*							68
El Salvador		*													12
French Guiana		*											*		25
México														*	80
Panama		*						*							30
Paraguay	X		*		*		X	X			X		X	*	24
Perú			*	*	X	X	X	X	*	X			*	*	95
Suriname													*	*	14
Uruguay			X				*	*						*	8
Venezuela		*						*					*	*	47

*Data from Young 1968, Takiya 2008, and Wilson et al. 2009.
X—Present work.

SUMMARY

Xylella fastidiosa is endemic to the Americas, it causes economically important diseases in a variety of different crops, and is transmitted by xylem-feeding sharpshooters. This paper provides new geographic records for Proconiini sharpshooters in South America which helps to better understand their distribution. To develop these new records, we examined material from 3 of the main entomological collections held in Argentina. As a result, 5 species are cited for the first time from Paraguay; 4 for Perú; 3 for Bolivia; 2 for Ecuador; and 1 each for Uruguay and Brazil. Some of the species could be vectors of *X. fastidiosa* because congeners of the species studied here are known to transmit this bacterium.

REFERENCES CITED

- AGUILAR, E., VILLALOBOS, W., MOREIRA, L., RODRIGUEZ, C. M., KITAJIMA, E. W., AND RIVERA, C. 2005. First report of *Xylella fastidiosa* infecting citrus in Costa Rica. *Plant Dis.* 89: 687.
- DAMSTEEGT, V. D., BRLANSKY, R. H., PHILLIPS, P. A., AND ROY, A. 2006. Transmission of *Xylella fastidiosa*, causal agent of Citrus Variegated Chlorosis, by the glassy-winged sharpshooter, *Homalodisca coagulata*. *Plant Dis.* 90: 567-570.
- HERNANDEZ-MARTINEZ, R., PINCKARD, T. R., COSTA, H. S., COOKSEY, D. A., AND WONG, F. P. 2006. Discovery and characterization of *Xylella fastidiosa* strains in southern California causing mulberry leaf scorch. *Plant Dis.* 90:1143-1149.
- HOPKINS, D. L. 1989. *Xylella fastidiosa*: Xylem-limited bacterial pathogen of plants. *Annu. Rev. Phytopathol.* 27: 271-290.
- LOPES, J. R. S. 1996. Mecanismo de transmissão de *Xylella fastidiosa* por cigarrinhas. *Laranja* 17: 79-92.
- MARUCCI, R. C., CAVICHIOLI, R. R., AND ZUCCHI, R. A. 2002. Espécies de cigarrinhas (Hemiptera, Cicadellidae, Cicadellinae) em pomares de citros da região de Bebedouro, SP, com descrição de uma nova espécie de *Acrogonia* Stål. *Rev. Brasileira Entomol.* 46 (2): 149-164.
- PILKINGTON, L. J., IRVIN, N. A., BOYD, E. A., HODDLE, M. S., TRIAPITSYN, S. V., CAREY, B. G., JONES, W. A., AND MORGAN, D. J. W. 2005. Introduced parasitic wasps could control glassy-winged sharpshooter. *California Agric.* 59: 223-228.
- REDAK, R., PURCELL, A., LOPES, J. R. S., BLUA, M., MIZELL, R. F., AND ANDERSEN, P. C. 2004. The biology of Xylem Fluid-Feeding Insect Vectors of *Xylella fastidiosa* and their relation to disease epidemiology. *Ann. Rev. Entomol.* 49: 243-270.
- TAKIYA, D. 2008. Sharpshooter (Cicadellinae) database. <http://ctap.inhs.uiuc.edu/takiya/index.asp> (accessed online Feb 2011).
- WELLS, J. M., RAJU, B. C., HUNG, H. Y., WEISBURG, W. G., MANDELCO-PAUL, L., AND BRENNER, D. J. 1987. *Xylella fastidiosa*: Gram-negative, xylem-limited, fastidious plant bacteria related to *Xanthomonas*. *Intl. J. Syst. Bacteriol.* 37: 136-143.
- WILSON, M. R., TURNER, J. A., AND MCKAMEY, S. H. 2009. Sharpshooter Leafhoppers of the World (Hemiptera: Cicadellidae: Cicadellinae). <http://naturalhistory.museumwales.ac.uk/Sharpshooters> (accessed online Feb 2011).
- YOUNG, D. A. 1968. Taxonomic Study of the Cicadellinae (Homoptera: Cicadellidae). Part 1. Proconiini. *Bull. United States Nat. Mus.* 261: 1-287.