

## **Lasioderma serricorne (Coleoptera: Anobiidae): First Record in Dehydrated Bee Pollen in Sergipe State, Brazil**

Authors: Poderoso, Julio César M., Correia-Oliveira, Maria E., Vieira, Jaci Mendes, Ribeiro, Genésio T., Ribeiro, Rafael C., et al.

Source: Florida Entomologist, 96(2) : 682-685

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.096.0247>

---

BioOne Complete ([complete.BioOne.org](http://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## LASIODERMA SERRICORNE (COLEOPTERA: ANOBIIDAE): FIRST RECORD IN DEHYDRATED BEE POLLEN IN SERGIPE STATE, BRAZIL

JÚLIO CÉSAR M. PODEROSO<sup>1</sup>, MARIA E. CORREIA-OLIVEIRA<sup>2</sup>, JACI MENDES VIEIRA<sup>2</sup>, GENÉSIO T. RIBEIRO<sup>3</sup>, RAFAEL C. RIBEIRO<sup>1</sup>  
AND JOSÉ C. ZANUNCIO<sup>1,\*</sup>

<sup>1</sup>Departamento de Entomologia, Universidade Federal de Viçosa. Viçosa, Minas Gerais State, Brazil

<sup>2</sup>Departamento de Entomologia e Acarologia, Escola Superior de Agricultura Luiz de Queiroz/Universidade de São Paulo, Piracicaba, São Paulo State, Brazil

<sup>3</sup>Departamento de Engenharia Florestal, Universidade Federal de Sergipe, São Cristóvão. Sergipe State, Brazil

\*Corresponding author; E-mail: zanuncio@ufv.br

Coleopterans have global importance in grain storage and industrial food production and can cause damage to the appearance of products as well as weight loss and reduced nutrient levels, rendering products unfit for human consumption (Nansen et al. 2004). *Lasioderma serricorne* (Fabricius, 1792) (Coleoptera: Anobiidae), variously known as the cigarette beetle or the tobacco beetle, is cosmopolitan and occurs frequently in tropical and subtropical areas. This insect is usually found in perishable food products such as dry fruit, grain, crumbs, cereal flour and animal food (Papadopoulou & Buchelos 2002).

Pollen, which adheres to honey bees (*Apis mellifera* L.) as they collect nectar from flowers and which they agglutinate into balls with nectar and bee saliva is important for survival of these insects. Beekeepers remove pollen for processing, commercialization and human consumption as a protein source (Mattila et al. 2007; Bastos et al. 2000).

Damage to bee pollen stored at 10% RH was found in samples collected in Sergipe State, Brazil. The purpose of this study was to characterize the damaged pollen and to ascertain its cause.

Pollen samples were obtained from the producers of the municipalities of Pacatuba, Brejo Grande, Neópolis, Estância and Tobias Barreto in Sergipe State, Brazil in Feb (summer) and Jun (winter) 2007. A bee returning to the hive after collecting nectar from flowers passes through a retention grill at the entrance that allows her passage but removes the pollen ball, which then falls into a box. This material is collected by the beekeeper, stored at 4 °C for 48 h and is air-dried at 40 °C for 48 h. Any visible impurities such as insect fragments, fibers, propolis, balls, etc., are removed from the dry material.

The pollen samples were supplied by producers in commercial packaging of plastic or polyethylene terephthalate (PET) bags. The pollen was stored in 50 mL plastic packages at an average temperature of 28 °C and 80% RH in the Agricultural and Forest Pests Laboratory of the Department of Forest Engineering of the Federal University of Sergipe.

The beetles were removed from contaminated samples and stored in 70% (v/v) alcohol. Adults were pinned in the laboratory and sent to the Taxonomy Laboratory of the Entomology and Acarology Department of ESALQ, University of São Paulo, for identification.

Palynological analysis was done in the Agricultural and Forest Pests Laboratory of the Federal University of Sergipe. This analysis compared pollen grains with those of an image bank to identify the botanic plant families that bees have visited to collect nectar, i.e., the images of the pollen grains were compared to those of known plant families. Microscope slides of the pollen samples were made with the acetolysis method (Erdtman 1960; Barth & Luz 1998; Terrab et al. 2003) and photomicrographed by a digital system of photodocumentation connected to an optical microscope with objectives of 40' and 100', and a quantitative assessment was made with 300 pollen grains per slide. The occurrence of each type of pollen (botanical family to which it belonged) was classified as: dominant ( $\geq 45\%$ ), accessorial (16-45%), or occasional ( $\leq 15\%$ ). The water content of 2 g samples of pollen was measured gravimetrically with an infrared drying system (Mettler LP16).

Insects found in pollen samples were identified as the tobacco beetle *Lasioderma serricorne* (Fabricius, 1792) (Coleoptera: Anobiidae) (Fig. 1). The hard plastic packages containing the pollen samples were undamaged (Fig. 2), so their infestation must have occurred in the field or during processing and sample handling. However, this beetle can cut into plastic bags and cause problems for the food industry (Arbogast et al. 2005; Begum et al. 2007). This beetle reproduces and the larvae grow and develop into adults when feeding on stored bee pollen.

Damage to pollen balls was similar to that observed in grains, with the consumption of the internal part, leaving only a thin husk. *Lasioderma serricorne* larvae also feed on pollen with galleries inside the packages of these products in which they leave excretions and feces. As a result, the pollen grains become lumpy and unfit for human or animal consumption (Fig. 1).




Fig. 1. *Lasioderma serricorne* (Coleoptera: Anobiidae), the cigarette beetle, found in dehydrated pollen. a. side view of the insect; b. dorsal view; c. damage by the cigarette beetle, *Lasioderma serricorne* (Coleoptera: Anobiidae) in dehydrated pollen; d. appearance of the infested product.

Insect contamination was observed in dehydrated bee pollen from Neópolis, Pacatuba, Brejo Grande and Estância municipalities which had a water content > 8% (Table 1).



Fig. 2. Container with dehydrated pollen with a heavy infestation by the cigarette beetle, *Lasioderma serricorne* (Coleoptera: Anobiidae).

TABLE 1. RESULTS OF THE PERCENTAGE OF MOISTURE IN DEHYDRATED SAMPLES OF POLLEN DURING THE SUMMER (FEB) AND WINTER (JUN) 2007 PRODUCED IN VARIOUS MUNICIPALITIES OF SERGIPE, BRAZIL.

Municipalities	February/2007	June/2007
Brejo Grande	10.6	9.9
Estância	10.5	12.5
Neópolis	10.3	10.93
Pacatuba	9.83	8.96
Tobias Barreto	11.66	11.16

The dehydrated bee pollen from Tobias Barreto was the only product not infested with *L. serricorne* beetles. Bee pollen from this municipality had a moisture level > 8%. Pollen from Areaceae species was < 10% of the total in bee pollen from Tobias Barreto, whereas Areaceae pollen occurred more frequently in samples from the other 4 municipalities, suggesting that *L. serricorne* might have a preference for this type of pollen (Table 2).

*Lasioderma serricorne* has been found in packaged foods destined for birds, cats and dogs in the

TABLE 2. BOTANICAL FAMILIES FOUND IN SAMPLES OF POLLEN DEHYDRATED IN SERGIPE. D: DOMINANT POLLEN GRAIN,  $\geq 45\%$  OF REPRESENTATIVENESS; A: ACCESSORIAL, 16 -45%; O: OCCASIONAL,  $\leq 15$

Cities	Botanical Family	
	February	June
Brejo Grande	Compositae (D); Arecaceae (A); Fabaceae, Mimo-Arecaceae (D); Compositae(A); Mimosaceae, Rubiaceae and Rubiaceae (O)	biaceae and Myrtaceae (O)
Estância	Arecaceae, Amaranthaceae, Compositae (A); Mi-Arecaceae, Compositae (A); Mimosaceae, Myrtaceae, Fabaceae and Labiatae (O)	ceae, Fabaceae and Rubiaceae (O)
Neópolis	Mimosaceae (D); Arecaceae (A); Compositae, An-Mimosaceae, Arecaceae, Compositae (A); Rubiacardiaceae and Myrtaceae (O)	ceae (O)
Pacatuba	Arecaceae (D); Compositae (A)	Arecaceae (D); Mimosaceae (A); Compositae (O)
Tobias Barreto	Myrtaceae, Solanaceae (A), Arecaceae, Composi-Compositae, Labiatae (A); Amaranthaceae, Fatae and Mimosaceae (O)	biaceae, Mimosaceae and Arecaceae (O)

USA (Roesli et al. 2003) and in wheat flour, maize and oat in Portugal, and it is common in factories using these cereals as raw material (Carvalho et al. 2006). This beetle infested tobacco and processed foods in Greece (Papadopoulou & Buchelos 2002). Despite its wide range of damaged products, *L. serricorne* has not been reported previously to infest stored pollen.

The water content of the dehydrated bee pollen samples was  $> 8\%$ , which might be important for *L. serricorne* reproduction; however, this beetle was not found in pollen collected from Tobias Barreto. Dehydrated bee pollen is supposed to have a maximum water content of 4% but it can change during storage (Almeida-Muradian et al. 2005).

Dehydrated bee pollen contains high levels of proteins, amino acids, lipids and secondary nutrients such as vitamins, which makes it a desirable food source for *L. serricorne* (Shinoda & Fujiasaki 2001; Patt et al. 2003; Lundgren & Wiedenmann 2004); however, its nutrient content varies among main botanic families and species of the same family (Richards & Schmidt 1996). The pollen from Tobias Barreto without *L. serricorne* had lower percentage from species of the botanic family Arecaceae compared to all other samples.

This is the first report of *L. serricorne* in dehydrated bee pollen. This beetle damages the dehydrated pollen balls, making them unfit for human or animal consumption.

#### SUMMARY

Bee pollen, which is the material that adheres to honeybees as they collect nectar from flowers, is agglutinated into balls with nectar and bee saliva substances and is important to the survival of colonies of these insects. Damage to bee pollen stored at 10% RH was found in samples collected in Sergipe State, Brazil. Insects found in the pollen were identified as *Lasioderma ser-*

*ricorne* (Fabricius, 1792) (Coleoptera: Anobiidae), a cosmopolitan insect and one of the main insects infesting tobacco and many stored food products. This is the first record of damage by this insect in dehydrated pollen balls, which makes this product unfit for human consumption.

Key Words: apiarian products, stored products, tobacco beetle

#### RESUMO

O pólen de abelha, o material levado por abelhas quando coletam o néctar das flores, é aglutinado em bolas com substâncias da saliva e importante para a sobrevivência das colônias desses insetos. Danos ao pólen de abelha armazenado a 10% de umidade foram encontrados em amostras coletadas no Estado de Sergipe, Brasil. Insetos encontrados no pólen foram identificados como *Lasioderma serricorne* (Fabricius, 1792) (Coleoptera: Anobiidae), um inseto cosmopolita e um dos principais insetos que infestam tabaco, assim como muitos outros produtos alimentares armazenados. Este é o primeiro registro de danos causados por este inseto e tornando o pólen desidratado impróprio para consumo humano.

Palavras Chave: Besouro do tabaco, Produtos Apícolas, produtos armazenados

#### ACKNOWLEDGMENTS

We express thanks to “Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)”, “Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES)” and “Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG)” for financial support.

#### REFERENCES CITED

ALMEIDA-MURADIAN, L. B., PAMPLONA, L. C., COIMBRA, S., AND BARTH, O. M. 2005. Chemical composition and

- botanical evaluation of dried bee pollen pellets. *J. Food Comp. Analysis* 18: 105-111.
- ARBOGAST, R. T., CHINI, S. R., AND MCGOVERN, J. E. 2005. Use of contour analysis in monitoring stored-product insects. *J. Econ. Entomol.* 99: 601-603.
- BARTH, O. M., AND LUZ, C. F. P. 1998. Melissopalynological data obtained from a mangrove area near to Rio de Janeiro, Brazil. *J. Apicult. Res.* 37: 155-163.
- BASTOS, E. M., OLIVEIRA, V. D. C., AND SOARES, A. E. E. 2000. Microscopic characterization of the green própolis, produced in Minas Gerais State, Brazil. *Honeybee Sci.* 21: 179-180.
- BEGUM, K., REDDY, P. V., LEEAJA, B. C., RAJASHEKAR, Y., AND RAJENDRAN, S. 2007. Studies on insect infestation in chocolates. *J. Stored Prod. Res.* 43: 118-122.
- CARVALHO, M. O., PASSOS DE CARVALHO, J., TORRES, L. M., AND MEXIA, A. 2006. Developing sequential plans for classifying *Lasioderma serricornis* (F) (Coleoptera, Anobiidae) status in a cigarette factory. *J. Stored Prod. Res.* 42: 42-50.
- ERDTMAN, G. 1960. The acetolysis method, a revised description. *Svensk Bot Tidskr* 54: 561-564.
- LUNDGREN, J. G., AND WIEDENMANN, R. N. 2004. Nutritional suitability of corn pollen for the predator *Coleomegilla maculata* (Coleoptera: Coccinellidae). *J. Insect Physiol.* 50: 567-575.
- MATTILA, H. R., AND OTIS, G. W. 2007. Dwindling pollen resources trigger the transition to broodless populations of long-lived honeybees each autumn. *Ecol. Entomol.* 32: 496-505.
- NANSEN, C., SUBRAMANYAM, B., AND ROESLI, R. 2004. Characterizing spatial distribution of trap captures of beetles in retail pet stores using SADIE® software. *J. Stored Prod. Res.* 40: 471-483.
- PAPADOPOULOU, S. C., AND BUCHELOS, C. T. 2002. Identification of female adult *Lasioderma serricornis* (F.) by simple external observation of the abdomen. *J. Stored Prod. Res.* 38: 315-318.
- PATT, J. M., WAINRIGHT S. C., HAMILTON, G. C., WHITTINGHILL, D., BOSLEY, K., DIETRICK, J., AND LASHOMB, J. H. 2003. Assimilation of carbon and nitrogen from pollen and nectar by a predaceous larva and its effects on growth and development. *Ecol. Entomol.* 28: 717-728.
- RICHARDS, P. C., AND SCHMIDT, J. M. 1996. The effect of selected dietary supplements on survival and reproduction of *Orius insidiosus* (Say) (Hemiptera: Anthicoridae). *Can. Entomol.* 128: 171-176.
- ROESLI, R., SUBRAMANYAM, J. F. C., AND KEMP, K. 2003. Stored-product insects associated with a retail pet store chain in Kansas. *J. Econ. Entomol.* 96: 1958-1966.
- SHINODA, K., AND FUJISAKI, K. 2001. Effect of adult feeding on longevity and fecundity of the cigarette beetle, *Lasioderma serricornis* F. (Coleoptera: Anobiidae). *Appl. Entomol. Zool.* 36: 219-223.
- TERRAB, A., DÍEZ, M. J., AND HEREDIA, F. J. 2003. Palynological, physicochemical and colour characterisation of Moroccan honeys: I. River red gum (*Eucalyptus camaldulensis* Dehnh) honey. *Int. J. Food Sci. Tech.* 38: 379-386.