



## **Manduca rustica (Lepidoptera: Sphingidae) Damage on Olive (*Olea europaea*; Lamiales: Oleaceae) trees in Florida**

Authors: Gillett-Kaufman, Jennifer L., Allan, Sandra A., and Buss, Lyle J.

Source: Florida Entomologist, 98(4) : 1260-1261

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.098.0443>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# *Manduca rustica* (Lepidoptera: Sphingidae) damage on olive (*Olea europaea*; Lamiales: Oleaceae) trees in Florida

Jennifer L. Gillett-Kaufman<sup>1,\*</sup>, Sandra A. Allan<sup>2</sup>, and Lyle J. Buss<sup>1</sup>

There is a growing interest in cultivating olives *Olea europaea* L. (Lamiales: Oleaceae) in Florida, USA. Although some parts of Florida seem well suited for olive production, the pests and diseases associated with cultivated olives in Florida are generally unknown. During a 2014 survey of olives being grown at the commercial and homeowner level, a novel pest was identified. A hornworm (Fig. 1), now known to be the larval stage of the rustic sphinx *Manduca rustica* (F.) (Lepidoptera: Sphingidae) (Wagner 2010; D. L. Wagner, personal communication), was reported as a pest of concern by growers from the Florida Panhandle in the north to Polk County in the south. Tuttle (2007) described *M. rustica* as a broadly distributed Neotropical species that breeds in the mid-Atlantic states as far north as Delaware, USA. In Florida, Heppner (2003) reported *M. rustica* ranges as far south as Miami-Dade County, but the species has not been reported from Monroe County (the Florida Keys). Worldwide, there are 49 recorded lepidopteran pests of *O. europaea*, 9 of which are in the family Sphingidae, but none is within the genus *Manduca* (Robinson et al. 2010). The primary concern of

growers was the defoliation caused by larval feeding. One larva can quickly denude a 2-yr-old olive tree, causing problems for nursery producers and for growers with new plantings. Although trees can recover from this damage, olive trees are slow growing, and feeding damage can remain apparent for several months depending on the time of year.

*Manduca rustica* is listed as having a host range that includes plants in the family Oleaceae (Kitching & Cadiou 2000; Tuttle 2007; Robinson et al. 2010); however, we were unable to find any report that it has been recorded as feeding on *O. europaea*. The rustic sphinx moth is recognized as a pollinator in the southeastern United States (Graham 2010). The only notable report of the rustic sphinx as a pest comes from Clavijo & Chacín (1992), who reported *M. rustica* as a pest of sesame in Venezuela. This is important to note for growers who are interested in using sesame as a cover crop or an attractant for beneficial insects near olive production areas in Florida.

Initial reports from growers included photos of specimens collected from orchards and nurseries. Many growers elected to control mechani-



Fig. 1. The rustic sphinx *Manduca rustica* found feeding on olive *Olea europaea* in Suwannee County, Florida. Photograph by Lyle J. Buss.

<sup>1</sup>University of Florida, Entomology and Nematology Department, Gainesville, Florida 32611, USA

<sup>2</sup>United States Department of Agriculture, Agriculture Research Service, Center for Medical, Agricultural, and Veterinary Entomology, 1700 SW 23rd Dr., Gainesville, Florida 32608, USA

\*Corresponding author; E-mail: gillett@ufl.edu



**Fig. 2.** A parasitized rustic sphinx *Manduca rustica* found on olive *Olea europaea* in Alachua County, Florida. Protruding larva (indicated by the red arrow), cocoons, and an adult *Cotesia congregata* can be seen in the photograph. Photograph by Lyle J. Buss.

cally (hand pick) the larvae when possible, but some resorted to using insecticide applications. It is interesting to note that one grower reported finding several larvae with white bumps. We collected a specimen and were able to identify the bumps as cocoons of *Cotesia congregata* (Say) (Hymenoptera: Braconidae) (Fig. 2). This initial collection of a parasitized larva was from Alachua County, Florida. To date, this is the first finding of *C. congregata* from *M. rustica*. However, a photograph of a larva with cocoons and exit wounds was submitted from a commercial olive producer in Suwannee County, Florida. As more olive groves are planted in Florida, we anticipate that more feeding damage will be caused by *M. rustica*. It remains to be determined if *C. congregata* will provide some level of control of *M. rustica* for Florida olive growers.

All photographs were taken by Lyle J. Buss. We acknowledge identification assistance from David Wagner (*M. rustica*), University of Connecticut, and Christopher Crockett (*C. congregata*), University of Florida. We would like to thank the president of the Florida Olive Council, Michael Garcia, for assistance locating producers and samples. We also would like to thank the following Florida producers for supplying samples and information on olive pests: Kent Cannon, Alachua County; Richard Williams and family, Florida Olive Systems, Volusia County; Rik Davis, Certified Crop Consultant, Volusia County; and Jonathan Carter, Florida Olive Farms, Suwannee County.

## Summary

The rustic sphinx *Manduca rustica* (F.) (Lepidoptera: Sphingidae) has been identified as an occasional pest for olive *Olea europaea* L. (Lamiales: Oleaceae) in Florida, USA. Reports of this pest's distribution on olive trees in Florida range from the northern Panhandle as far south as Polk County.

Key Words: rustic sphinx; feeding damage; *Cotesia congregata*

## Sumario

El esfingido rústico *Manduca rustica* (F.) (Lepidoptera: Sphingidae) ha sido identificado como una plaga ocasional del olivo, *Olea europaea* L. (Lamiales: Oleaceae), en la Florida, EE.UU. Los informes de la distribución de esta plaga en los árboles de olivo en la Florida son desde el norte de Panhandle hasta el sur del condado de Polk.

Palabras Clave: esfinge rústico; daño por alimentación; *Cotesia congregata*

## References Cited

- Clavijo JA, Chacín ME. 1992. Sphingidae (Insecta: Lepidoptera) reported as pests of Venezuelan crops: key to the species. *Boletín de Entomología Venezolana* 7: 119-125.
- Graham SP. 2010. Visitors to southeastern hawkmoth flowers. *Southeastern Naturalist* 9: 413-426.
- Heppner JB [ed.]. 2003. *Arthropods of Florida and Neighboring Land Areas, Volume 17: Lepidoptera of Florida*. Florida Department of Agriculture and Consumer Services, Division of Plant Industry. Gainesville, Florida, USA.
- Kitching IJ, Cadiou J-M. 2000. *Hawkmoths of the World: An Annotated and Illustrated Revisionary Checklist (Lepidoptera: Sphingidae)*. Cornell University Press, Ithaca, New York, USA.
- Robinson GS, Ackery PR, Kitching IJ, Beccaloni GW, Hernández LM. 2010. *HOSTS - A Database of the World's Lepidopteran Hostplants*. Natural History Museum, London, United Kingdom, <http://www.nhm.ac.uk/hosts> (last accessed: 27 Oct 2014).
- Tuttle JP. 2007. *The Hawk Moths of North America: A Natural History Study of the Sphingidae of the United States and Canada*. Wedge Entomological Research Foundation, Washington, District of Columbia, USA.
- Wagner DL. 2010. *Caterpillars of Eastern North America: A Guide to Identification and Natural History*. Princeton University Press, Princeton, New Jersey, USA.