

Delayed Emergence of *Trachykele blondeli blondeli* (Coleoptera: Buprestidae)

Authors: Basham, Joshua P., Oliver, Jason B., and Herzog, Claire E.

Source: Florida Entomologist, 99(3) : 578-579

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.099.0344>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Delayed emergence of *Trachykele blondeli blondeli* (Coleoptera: Buprestidae)

Joshua P. Basham¹, Jason B. Oliver^{1,*}, and Claire E. Herzog²

Delayed and prolonged emergence of Buprestidae occurs in several genera. Huguenin (1915) was the first to report the phenomenon of delayed emergence of *Buprestis (Cypriacis) aurulenta* L. (Coleoptera: Buprestidae), emerging after 26 yr from a door sill that had been varnished and painted twice. Smith (1962) reported 32 instances of *B. aurulenta* emerging after 26 yr and the longest of 51 yr, setting a new record for the longest insect life cycle.

On 1 Mar 2015, C. E. Herzog witnessed a buprestid emerging from a door frame of a house in Sarasota, Florida. She photographed the emerging adult and placed the images on BugGuide (Herzog 2015) for identification. A tentative identification of *Trachykele blondeli* Marseul (Coleoptera: Buprestidae) was made by Vassili Belov. The specimen identification was subsequently confirmed by the first author as *Trachykele blondeli blondeli* Marseul (Coleoptera: Buprestidae), the western cedar borer. The specimen emerged from a 2.5 × 10 × 20 cm kiln-dried cedar board purchased at a Sarasota, Florida, lumber supply store during Mar 2013 and subsequently installed and painted shortly afterward. The specimen collection data is FLORIDA, Sarasota Co., Sarasota, 1-III-2015, C. Herzog. The specimen currently is housed in TSRS (Evenhuis & Samuelson 2007). The report is the first known collection of *T. blondeli blondeli* in Florida. The western cedar borer is endemic from British Columbia south to Arizona, New Mexico, and Mexico (Nelson et al. 2008).

The initial genus and species description of western cedar borer was based on a single specimen found at a customs warehouse in Paris, France (Marseul 1865). The first host record description was described as “. . . sortant d'un cèdre du Liban” or out of a cedar of Lebanon (*Cedrus libani* A. Rich.; Pinales: Pinaceae), and the type locality was deemed to be Liban (i.e., Lebanon) (Marseul 1865). Burke (1928) stated that the wood's origin was never accurately determined, but must be from an unknown North American cedar species (Pinales: Cupressaceae), because all other known collections of *T. blondeli blondeli* are from [western] North America. Eight species and subspecies of *Trachykele* are known to occur in the United States. Seven *Trachykele* species utilize hosts in the family Cupressaceae, whereas the 8th, *Trachykele nimbose* Fall (Coleoptera: Buprestidae), uses *Abies* spp. (Pinales: Pinaceae) and *Tsuga mertensiana* (Bong.) Carrière (Pinales: Pinaceae) (Nelson et al. 2008).

This report highlights the potential for Buprestidae to remain undetected in lumber products and to be redistributed to new areas. The report also provides evidence for *T. blondeli blondeli* survival during the lumber kiln-drying process, but it is unknown whether the specimen was in the larval, pupal, or adult stage during kiln drying.

We thank Nadeer Youssef for his input on this report and Ted MacRae, Hans Mühle, Mark Volkovitch, and Rick Westcott for their assistance with finding the original description of *T. blondeli* Marseul and for constructive comments to improve the manuscript.

Summary

Prolonged emergence of *Trachykele blondeli blondeli* Marseul (Coleoptera: Buprestidae) is reported from kiln-dried and subsequently painted wood in this report. The specimen emergence is a new record for Florida. Because the specimen did not emerge from host plant material grown in Florida, it is likely that its origin was exterior to Florida. Because of delayed and prolonged emergence capability in Buprestidae, as well as specimen survival of wood preservation processes like kiln drying, the current report does indicate a concern for redistribution of non-indigenous buprestid species to new areas.

Key Words: BugGuide; metallic wood-boring beetle; flatheaded borer; western cedar borer; Florida; kiln; lumber

Sumario

Se informa de la emergencia prolongada de *Trachykele blondeli blondeli* Marseul (Coleoptera: Buprestidae) demadera secada en el horno y pintada posteriormente. La emergencia del espécimen es un nuevo registro para la Florida. Dado que el espécimen no emergió de material de una planta hospedera cultivada en Florida y el origen del producto, la madera es desconocida, es probable el origen espécimen era exterior a la Florida. Debido a la capacidad de aparición retardada y prolongada en Buprestidae, así como la supervivencia de muestras de los procesos de conservación de la madera, como el secado en estufa, el informe actual no indica una preocupación por la redistribución de las especies buprestid no indígenas a nuevas áreas.

Palabras Clave: BugGuide; escarabajo barrenador de madera metálico; barrenador de cabeza plana; barrenador del cedro occidental; Florida; horno; madera

References Cited

Burke HE. 1928. The western cedar pole borer or powder worm. United States Department of Agriculture Technical Bulletin 48: 1–15.

¹Tennessee State University, College of Agriculture, Human, and Natural Sciences, Otis L. Floyd Nursery Research Center, 472 Cadillac Lane, McMinnville, Tennessee 37110, USA; E-mail: Joshua.basham@tn.gov (J. P. B.), joliver@tnstate.edu (J. B. O.)

²5531 Calle del Verano, Sarasota, Florida 34242, USA; E-mail: ceherzog@gmail.com (C. E. H.)

*Corresponding author; E-mail: joliver@tnstate.edu (J. B. O.)

- Evenhuis NL, Samuelson GA. 2007. The insect and spider collections of the world website. <http://hbs.bishopmuseum.org/codens/codens-r-us.html> (last accessed 21 Dec 2015).
- Herzog CE. 2015. Western cedar borer (*Trachykele blondeli*). BugGuide.net. Mar 1, 2015. Website. <http://bugguide.net/node/view/1043490> (last accessed 3 May 2016).
- Huguenin JC. 1915. An observation on a buprestid. Entomological News 26: 364–365.
- Marseul S-A de. 1865. Monographie des buprestides famille des sternoxes de Latreille. L' Abeille, Mémoires d'Entomologie 2: 1–396. [English summary including tribal key in Zoological Record for 1865]
- Nelson GH, Walters Jr GC, Haines RD, Bellamy CL. 2008. A Catalog and Bibliography of the Buprestoidea of America North of Mexico. The Coleopterists Society Special Publication Number 4. The Coleopterists Society, North Potomac, Maryland.
- Smith DN. 1962. Prolonged larval development in *Buprestis aurulenta* L. (Coleoptera: Buprestidae). A review with new cases. The Canadian Entomologist 94: 586–593.