

## **First Record of *Heliothrips haemorrhoidalis* (Thysanoptera: Thripidae) Causing Damage on Greenhouse Strawberries**

Authors: de Souza, Mireli Trombin, de Souza, Michele Trombin, Dybas Ivankio, Launa Bianca, Durau, Bruna Caroline, Maleski, Letícia Tamara, et al.

Source: Florida Entomologist, 102(3) : 651-653

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.102.0326>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# First record of *Heliothrips haemorrhoidalis* (Thysanoptera: Thripidae) causing damage on greenhouse strawberries

Mireli Trombin de Souza<sup>1,2,\*</sup>, Michele Trombin de Souza<sup>1,2</sup>, Launa Bianca Dybas Ivankio<sup>1</sup>, Bruna Caroline Durau<sup>1</sup>, Leticia Tamara Maleski<sup>1</sup>, and Maria Aparecida Cassilha Zawadneak<sup>1,2</sup>

Characteristic silvery patches made by thrips (Thysanoptera) were observed on leaves of semi-hydroponic greenhouse strawberries (*Fragaria × ananassa* Duch.; Rosaceae) 'Camino Real' cultivar, in Curitiba, Paraná, Brazil (25.428888°S; 49.267500°W), in Sep 2017. Forty-nine adult insects were collected and preserved in AGA solution (1:1:10 parts of glycerin, glacial acetic acid, and ethyl alcohol 60%, respectively). Specimens were slide-mounted (Mound & Kibby 1998) and identified as the greenhouse thrips, *Heliothrips haemorrhoidalis* (Bouché) (Thysanoptera: Thripidae), based on specialized taxonomic keys (Mound & Marulo 1998; Nakahara et al. 2015).

All specimens obtained were females (Fig. 1a), with strongly reticulated head sculpture (Fig. 1b); 8-segmented antennae, antennal segments III–V and VII–VIII yellow, VI brown in apical half (Fig. 1c); pronotum polygonally reticulate, without long setae; metanotum with a bold and reticulate triangle, median setae short (Fig. 1d); tergites III–IV strongly reticulated, the median setae long and close to each other (Fig. 1e); wings hyaline, with round apex; and yellowish legs (Fig. 1f). The body color presented variation according to the life stage: recently emerged individuals had the head and thorax dark brown and the abdomen light brown (Fig. 2a), whereas in mature females the abdomen was dark brown, except by the paler segments IX and X (Fig. 2b). According to Bernardo et al. (2005), males are rare in this species, and reproduction is predominately by thelytokous parthenogenesis, where unfertilized eggs produce females (Fig. 2c).

*Heliothrips haemorrhoidalis* is native from South America, and it is known as highly polyphagous on the leaves of several unrelated plants. It has been reported on several crops, including coffee (*Coffea arabica* L.; Rubiaceae), orange (*Citrus aurantium* L.; Rutaceae), passionfruit (*Passiflora caerulea* Auct.; Passifloraceae) (Suris & González 2008), lime (*Citrus aurantiifolia* (Christm.) Swingle; Rutaceae), rose (*Rosa* sp. L.; Rosaceae), wild strawberry (*Fragaria vesca* L.; Rosaceae) (Etienne et al. 2015), and avocado (*Persea americana* Mill.; Lauraceae) (Larral et al. 2018). A greenhouse bioassay was conducted to confirm the damage caused by *H. haemorrhoidalis* in strawberries. Damage characterization was performed with cage-confined individuals following the methodology described by Nondillo et al. (2010). Each cage was set up with 1 leaf or 1 semi-ripe

strawberry fruit, free from arthropod-induced damage, resulting in a total of 20 replicates for each plant part. Larvae ( $n = 10$ ) and adults ( $n = 10$ ) were placed in each cage using a fine paintbrush and kept confined for 72 h. A control was set up with an insect-free cage. Feeding damage was found on leaves and sepals. Thrips are piercing-sucking feeders, piercing the plant surface to feed on chlorophyll and other cell contents from plant tissue (Chhagan & Stevens 2007). As a consequence, silver-colored areas and black spots, due to deposition of excrement, appeared around feeding areas (Fig. 2c). Feces may act as a protective barrier against natural enemies (Denmark & Fasulo 2010) since, after oviposition, *H. haemorrhoidalis* were observed depositing excrement over the eggs.

*Hyaliodescoris insignis* (Stål) (Heteroptera: Miridae) and *Orius insidiosus* (Say) (Hemiptera: Anthracoridae) were observed preying on *H. haemorrhoidalis* adults and immatures, including larvae, prepupae, and pupae (Fig. 2a, b, f, g, and h). This is the first record of *H. insignis* preying *H. haemorrhoidalis*. So far, biological control studies have indicated approximately 10 natural enemies of the greenhouse thrips (CABI 2018). However, a parasitoid, *Thripobius semiluteus* Bouček (Hymenoptera: Eulophidae), has been considered the most efficient natural enemy for the control of *H. haemorrhoidalis* (McMurtry 1992; Bernardo et al. 2005; CABI 2018).

Vouchers specimens were deposited in the Padre Jesus Moure Museum, Federal University of Paraná, Curitiba, Brazil: *H. haemorrhoidalis* (DZUP 519163), *H. insignis* (DZUP 490170), and *O. insidiosus* (DZUP 490169).

This is the first report of *H. haemorrhoidalis* causing damage on strawberry crops in the world. Observations of beneficial agents naturally controlling *H. haemorrhoidalis* populations may provide new perspectives for future biocontrol programs for managing this species.

We thank the Coordination for the Improvement of the Higher Level Personnel (CAPES), National Council for Scientific and Technological Development (CNPq), for the financial support; we are grateful to Paulo Fiuza Ferreira (Federal University of Viçosa) and the Germano Rosado Neto (Federal University of Paraná) for identification of *Hyaliodescoris insignis* and *Orius insidiosus*, respectively. We also thank BioAgro for providing the seedlings used in experiment.

<sup>1</sup>Federal University of Paraná, Department of Basic Pathology, Federal University of Paraná, Caixa Postal 19031, 81531-980, Curitiba, Paraná, Brazil; E-mail: mirelitrombin@gmail.com (M. T. S.); mictrombin@gmail.com (M. T. S.); laudybas@gmail.com (L. B. D. I.); brudurau535@gmail.com (B. C. D.); leticia19.maleski@gmail.com (L. T. M.); mazawa@ufpr.br (M. A. C. Z.)

<sup>2</sup>Postgraduate Program in Agronomy, Department of Phytotechnology and Plant Health, Federal University of Paraná, Curitiba, Paraná, Brazil.

\*Corresponding author; E-mail: mirelitrombin@gmail.com




Fig. 1. Morphological characteristics of *Heliothrips haemorrhoidalis*: (a) female, (b) head and pronotum reticulated, (c) antennae, (d) mesonotum, metanotum, and tergite I–II reticulated, (e) tergites III–VI, and (f) forewing and leg.

## Summary


*Heliothrips haemorrhoidalis* (Bouché, 1833) (Thysanoptera: Thripidae), the greenhouse thrips, is reported here for the first time damaging strawberry leaves and sepals (*Fragaria × ananassa* Duch.; Rosaceae) in the world. Two natural enemies were observed preying on the species: *Hyaliodocoris insignis* (Stål, 1860) (Heteroptera: Miridae) and *Orius insidiosus* (Say, 1832) (Hemiptera: Anthracoridae).

Key Words: *Fragaria × ananassa*; greenhouse thrips; leaf damage; natural enemies

## Sumário

*Heliothrips haemorrhoidalis* (Bouché, 1833) (Thysanoptera: Thripidae), o tripes-das-casas-de-vegetação, é relatado pela primeira vez danificando folhas e sépalas de morangueiro (*Fragaria × ananassa*).





**Fig. 2.** *Heliiothrips haemorrhoidalis* and its damage in strawberry plants: (a) teneral female, (b) mature female, (c) egg, (d) damage on leaves, (e) leaf chlorosis and fecal specks, (f) larva, (g) pre-pupa, and (h) pupa.

Duch.; Rosaceae) no mundo. Dois inimigos naturais foram observados predando a espécie: *Hyaliodocoris insignis* (Stål, 1860) (Heteroptera: Miridae) e *Orius insidiosus* (Say, 1832) (Hemiptera: Anthorcoridae).

Palavras Chave: *Fragaria × ananassa*; tripes-das-casas de-vegetação; dano foliar; inimigos naturais

## References Cited

- Bernardo U, Viggiani G, Sasso R. 2005. Biological parameters of *Thripobius semiluteus* Bouček (Hym., Eulophidae), a larval endoparasitoid of *Heliiothrips haemorrhoidalis* (Bouché) (Thysan., Thripidae). *Journal of Applied Entomology* 129: 250–257.
- CABI (Centre for Agriculture and Biosciences International). 2018. *Heliiothrips haemorrhoidalis* (black tea thrips). Wallingford, Oxfordshire, United Kingdom. <https://www.cabi.org/isc/datasheet/26818> (last accessed 18 May 2019).
- Chhagan A, Stevens PS. 2007. Effect of temperature on the development, longevity and oviposition of greenhouse thrips (*Heliiothrips haemorrhoidalis*) on lemon fruit. *New Zealand Plant Protection* 60: 50–55.
- Denmark HA, Fasulo TR. 2010. Greenhouse thrips, *Heliiothrips haemorrhoidalis* (Bouché) Insecta: Thysanoptera: Thripidae). UF/IFAS Publication EENY-75. University of Florida, Gainesville, Florida, USA. [https://entnemdept.ifas.ufl.edu/creatures/orn/thrips/greenhouse\\_thrips.htm](https://entnemdept.ifas.ufl.edu/creatures/orn/thrips/greenhouse_thrips.htm) (last accessed 18 May 2019).
- Etienne J, Ryckewaert P, Michel B. 2015. Thrips (Insecta: Thysanoptera) of Guadeloupe and Martinique: updated check-list with new information on their ecology and natural enemies. *Florida Entomologist* 98: 298–304.
- Larral P, Ripa R, Funderburk J, Lopez E. 2018. Population abundance, phenology, spatial distribution and a binominal sampling plan for *Heliiothrips haemorrhoidalis* (Thysanoptera: Thripidae) in avocado. *Florida Entomologist* 101: 166–171.
- McMurtry JA. 1992. The role of exotic natural enemies in the biological control of insect and mite pests of avocado in California, pp. 247–252, *In Proceedings of the Second World Avocado Congress*, 21–26 Apr 1991, Orange, California, USA.
- Mound LA, Kibby G. 1998. *Thysanoptera: An Identification Guide*, 2nd edition. CAB International, Wallingford, Oxfordshire, United Kingdom.
- Mound LA, Marullo R. 1998. *Thysanoptera: An Identification Guide*. 2nd edition. CAB International, Wallingford, Oxfordshire, United Kingdom.
- Nakahara S, O'donnell CA, Mound LA. 2015. *Heliiothrips haemorrhoidalis* and its relatives, with one new species and one new genus (Thysanoptera: Thripidae). *Zootaxa* 4021: 578–584.
- Nondillo A, Redaelli LR, Pinet SMJ, Botton M. 2010. Caracterização das injúrias causadas por *Frankliniella occidentalis* no morangueiro. *Ciência Rural* 40: 820–826.
- Suris M, González C. 2008. Especies de trips asociadas a hospedantes de interés en las provincias habaneras: II. Plantas frutales. *Revista de Protección Vegetal* 23: 85–89.