

First Report of *Gymnandrosoma aurantianum* (Lepidoptera: Tortricidae) in Pecan in Brazil

Authors: Nava, Dori Edson, Sturza, Vinícius Soares, and Martins, Carlos Roberto

Source: Florida Entomologist, 103(1) : 130-131

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.103.0422>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First report of *Gymnandrosoma aurantianum* (Lepidoptera: Tortricidae) in pecan in Brazil

Dori Edson Nava¹, Vinícius Soares Sturza^{2,*}, and Carlos Roberto Martins¹

The cultivation of hardy pecan, *Carya illinoensis* (Wangenh.) K. Koch (Juglandaceae), in Brazil started in the 1870s, but significant expansion occurred only during the 1960s and 1970s due to governmental stimuli aiming to bring in new investments for forestation (Backes & Irgang 2004; Ortiz & Camargo 2005). Economic exploration of pecan cultivation garnered attention in Brazil in the 21st century due to its high revenue, crop nutritional value, use of lands with irregular (sloping) surfaces, and the development of integrated systems with pastures for animal production, or the cultivation of pecan along with other crops (Fronza et al. 2013, 2018; Martins et al. 2017). In 2017, harvested areas were concentrated in the southeastern (21%) and southern (79%) regions of Brazil. The state of Rio Grande do Sul increased its production area to 86% between 2007 and 2017, and is currently the largest producing state, accounting for two-thirds of the total harvest area in Brazil (IBGE 2017). Simultaneously with the increase of pecan orchards in the country, pests seem to be damaging plants; however, information on this problem continues to be limited. This is the first report of the occurrence of *Gymnandrosoma aurantianum* (Lima) (Lepidoptera: Tortricidae) in pecan in Brazil.

Pecan samples were taken from 2 areas in May 2018: 1 in the municipality of Uraí in the state of Paraná (23.315200°S, 50.501339°W) from a pecan orchard of about 80 ha, which was planted in 1965. The second was taken in the municipality of Chapada in the state of Rio Grande do Sul (28.070686°S, 53.155628°W) from an orchard of about 5.2 ha, planted in 2009, which was attached to another area of about 27.6 ha of pecan trees planted between 2013 and 2014. In these 2 areas, plants were sampled randomly. Three nuts infested with lepidopteran caterpillars were found in Uraí and 5 were found in Chapada, respectively. Insects were fed in order to obtain adult moths (Fig. 1), which were identified according to Adamski & Brown (2001).

The damage on fruits occurred by excavation and perforation of the light green with darkish green outer pericarp layer (husk) that encloses the endocarp. The larval feeding resulted in deep galleries, even promoting some grooves in the outer pecan pericarp. Excrement (frass) of the larvae were found between the outer pericarp layer and the endocarp as shown in Fig. 2. On these infested pecans, the damage did not reach the endocarp, but galleries did compromise the pericarp, which might imply risk of contamination by microorganisms, and trouble for transportation and storage because of shorter shelf life.

Twenty phytophagous insects have been identified that feed on pecan plants, 3 of which are common in North America where this species is indigenous: *Phylloxera devastatrix* Pergande and *Phylloxera notabilis* Pergande (both Hemiptera: Phylloxeridae), and *Monellia caryella* (Fitch) (Hemiptera: Aphididae) (Paulsen et al. 2013;

Boscardin & Costa 2018). No species recorded in Brazil belongs to the Tortricidae family, although 2 species have been found in pecan nuts and trees in North America: *Cydia caryana* (Fitch) and *Gretchena bolliana* (Slingerland) (both Lepidoptera: Tortricidae) (Ree & Knutson 1997; Thompson & Conner 2012; Boscardin & Costa 2018).




Fig. 1. *Gymnandrosoma aurantianum*: dorsal (A) and lateral (B) view of fifth instar larvae, and adult moth (C).

¹Embrapa Clima Temperado, Pelotas, 96010-971, Brazil; E-mail: dori.edson-nava@embrapa.br (D. E. N.), carlos.r.martins@embrapa.br (C. R. M.)

²Universidade Federal de Pelotas, Programa de Pós-graduação em Fitossanidade, Capão do Leão, 96010-610, Brazil; E-mail: vsturza27@yahoo.com.br (V. S. S.)

*Corresponding author; E-mail: vsturza27@yahoo.com.br



Fig. 2. Larva of *Gymnandrosoma aurantianum* and injury to pecan pericarp.

Gymnandrosoma aurantiana is known in Brazil as the citrus fruit borer, and became a key pest by the end of the 1980s, when larval injuries on fruits became serious damage, and compromised the use of fruits for industry or for consumption in natura, resulting in loss of production up to 50% (Prates 1992; Parra et al. 2004; Carvalho et al. 2015). This led to research efforts aimed at lowering losses, and basic studies on biology, rearing techniques, monitoring, and use of egg parasitoids, which resulted in parameters for *G. aurantiana* pest management, particularly in the state of São Paulo, the largest citrus producer in Brazil (García & Parra 1999; Vilela et al. 2001; Parra et al. 2004; Bento et al. 2006; Molina & Parra 2006). In Mexico, the egg parasitoid *Trichogramma platneri* Nagarkatti (Hymenoptera: Trichogrammatidae) was effective in pecan orchards when the abundance of tortricidae *C. caryana* was low or moderate (García-Nevárez & Tarango-Rivero 2013).

In the southern states, particularly in Rio Grande do Sul, pest management is not common in small orchards with citrus plants, which is why the pest status of *G. aurantiana* is unknown. Moreover, the increase of pecan harvest areas, as well as the existence of abandoned orchards from the last century, may pose a threat to pecan production (Ortiz & Camargo 2005). These zones may serve as additional reproduction sites, and become sources of infestation for commercial pecan areas. Considering that other tortricid species already are recorded in pecan, and that *G. aurantianum* is a key pest in citrus plants and is hosted by other plant species (Silva et al. 1968; Molina & Parra 2006), it is reasonable to presume that the incidence of this pest in pecan could increase and eventually reach economic injury levels.

The authors thank Vitor O. Becker, the director of Scientific Research Institute “Uiraçu” for identification of *G. aurantiana*, Leyla Koc for the English review, and F. G. Gonçalves for providing pictures of *G. aurantiana* larvae.

Summary

The cultivation of hardy pecan, *Carya illinoensis* (Wangenh.) K. Koch (Juglandaceae), has increased in Brazil in the current century; however, the knowledge about insect pests is limited. This report deals with the occurrence of *Gymnandrosoma aurantianum* (Lima) (Lepidoptera: Tortricidae) in pecan trees for the first time in Brazil, discussing symptoms and possible risks.

Key Words: *Carya illinoensis*; Juglandaceae; Olethreutinae; Grapholitini; citrus fruit borer

Sumario

O cultivo de nogueira-pecã *Carya illinoensis* (Wangenh.) K. Koch (Juglandaceae), aumentou no Brasil no presente século, apesar disso informações sobre insetos praga são limitadas. Esse registro trata da ocorrência de *Gymnandrosoma aurantianum* (Lima) (Lepidoptera: Tortricidae) em nogueira-pecã no Brasil, relatando sintomas e possíveis riscos à cultura.

Palavras Chave: *Carya illinoensis*; Juglandaceae; Olethreutinae; Grapholitini; bicho furão dos citros.

References Cited

- Adamski D, Brown JW. 2001. Systematic revision of the Ecdytoplopha group of genera (Lepidoptera: Tortricidae: Grapholitini) in the New World. *Entomologica Scandinavica Supplements* 58.
- Backes P, Irgang B. 2004. Árvores do Sul. Guia de identificação & interesse Ecológico. As principais espécies nativas Sul-Brasileiras. Instituto Souza Cruz, Santa Cruz do Sul, Brasil.
- Bento JMS, Murata Y, Ono M, Parra JRP, Vilela EF. 2006. Identification, synthesis, and field evaluation of the sex pheromone from the citrus. *Journal of Chemical Ecology* 32: 155–168.
- Boscardin J, Costa EC. 2018. A nogueira-pecã no Brasil: uma revisão entomológica. *Ciência Florestal* 28: 456–468.
- Carvalho JHS, Barbosa JC, Yamamoto PT, Bicalho IB. 2015. Distribuição espacial do bicho-furão, *Gymnandrosoma aurantiana* (Lima, 1927) (Lepidoptera: Tortricidae), em citros utilizando geoestatística. *Revista Brasileira de Fruticultura* 37: 600–609.
- Fronza D, Poletto T, Hamann JJ. 2013. O cultivo de nogueira-pecã. Universidade Federal de Santa Maria, Santa Maria, Brazil.
- Fronza D, Hamann JJ, Both V, Anese R de O, Meyer EA. 2018. Pecan cultivation: general aspects. *Ciência Rural* 48: 1–9.
- García-Nevárez G, Tarango-Rivero SH. 2013. Efficacy of *Trichogramma platneri* (Hymenoptera: Trichogrammatidae) for the biological control of *Acrobasis nuxvorella* (Lepidoptera: Pyralidae) and *Cydia caryana* (Lepidoptera: Tortricidae). *Southwestern Entomologist* 38: 523–530.
- Garcia MS, Parra JRP. 1999. Comparação de dietas artificiais, com fontes protéicas variáveis, para criação de *Ecdytoplopha aurantiana* (Lima) (Lepidoptera: Tortricidae). *Anais da Sociedade Entomológica do Brasil* 28: 219–232.
- IBGE – Instituto Brasileiro de Geografia e Estatística. 2017. Produção Agrícola Municipal. <https://sidra.ibge.gov.br/tabela/1613#resultado> (last accessed 10 Oct 2019).
- Martins CR, Fronza D, Malgarim MB, Bilharva MG, De Marco R, Hamann JJ. 2017. Cultura da noz-pecã para a agricultura familiar, pp. 65–81 *In* Wolff LF, Medeiros CAB [eds.], Alternativas para a diversificação da agricultura familiar de base ecológica. Embrapa Clima Temperado, Pelotas, Brazil.
- Molina RMDS, Parra JRP. 2006. Seleção de linhagens de *Trichogramma* (Hymenoptera, Trichogrammatidae) e determinação do número de parasitóides a ser liberado para o controle de *Gymnandrosoma aurantianum* Lima (Lepidoptera, Tortricidae). *Revista Brasileira de Entomologia* 50: 534–539.
- Ortiz ERN, Camargo LEA. 2005. Doenças da nogueira pecan (*Carya illioensis*), pp. 501–506 *In* Kimati H, Amorim L, Resende JAM, Bergamin Filho A, Camargo LEA [eds.], Manual de fitopatologia. Agronômica Ceres, São Paulo, Brazil.
- Parra JRP, Bento JMS, Garcia MS, Yamamoto PT, Vilela EF, Leal WS. 2004. Development of a control alternative for the citrus fruit borer, *Ecdytoplopha aurantiana* (Lepidoptera, Tortricidae): from basic research to the grower. *Revista Brasileira de Entomologia* 48: 561–567.
- Paulsen CM, Cottrell TE, Ruberson JR. 2013. Distribution of the black pecan aphid, *Melanocallis caryaefoliae*, on the upper and lower surface of pecan foliage. *Entomologia Experimentalis et Applicata* 146: 252–260.
- Prates HS. 1992. Resultados recentes do controle do bicho-furão - lagarta da mariposa das laranjas - *Gymnandrosoma aurantianum* (Lima, 1927) em citros 1992. *Informativo Coopercitrus* 71: 20–21.
- Ree B, Knutson A. 1997. Field guide to the insects and mites associated with pecan. Texas A&M University Libraries, College Station, Texas, USA. <http://hdl.handle.net/1969.1/87786> (last accessed 10 Oct 2019).
- Silva AGA, Gonçalves CR, Galvão DM, Gonçalves AJL, Gomes J, Silva MN, De Simoni L. 1968. Quarto catálogo dos insetos que vivem nas plantas do Brasil: seus parasitos e predadores: insetos do Brasil: hospedeiros e inimigos naturais. Ministério da Agricultura, Rio de Janeiro, Brazil.
- Thompson TE, Conner P. 2012. Pecan, pp. 771–801 *In* Badenes ML, Byrne DH [eds.], Handbook of Plant Breeding. Fruit Breeding, Part 4. Springer, New York, USA.
- Vilela EF, Bento IS, Walder JM, Leal WS. 2001. Sexual behavior and diel activity of citrus fruit borer. *Journal of Chemical Ecology* 27: 2053–2065.