

Stable Isotopes and Radiocarbon Assess Variable Importance of Plants and Fungi in Diets of Arctic Ground Squirrels

Authors: Hobbie, Erik A., Shamhart, Julee, Sheriff, Michael, Ouimette, Andrew P., Trappe, Matt, et al.

Source: Arctic, Antarctic, and Alpine Research, 49(3) : 487-500

Published By: Institute of Arctic and Alpine Research (INSTAAR), University of Colorado

URL: <https://doi.org/10.1657/AAAR0016-062>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Stable isotopes and radiocarbon assess variable importance of plants and fungi in diets of arctic ground squirrels

Erik A. Hobbie^{1,*}, Julee Shamhart¹, Michael Sheriff², Andrew P. Ouimette¹, Matt Trappe³, Edward A. G. Schuur⁴, John E. Hobbie⁵, Rudy Boonstra⁶, and Brian M. Barnes⁷

¹Earth Systems Research Center, 8 College Road, Morse Hall, University of New Hampshire, Durham, New Hampshire 03824, U.S.A.

²Department of Ecosystem Science and Management, 405 Forest Resources Building, Pennsylvania State University, University Park, Pennsylvania 16802, U.S.A.

³College of Forestry, 140 Peavy Hall, 3100 SW Jefferson Way, Oregon State University, Corvallis, Oregon 97331, U.S.A.

⁴Center for Ecosystem Sciences and Society, Department of Biological Sciences, Box 5640, Northern Arizona University, Flagstaff, Arizona 86011, U.S.A.

⁵Ecosystems Center, Marine Biological Laboratory, 7 MBL Street, Woods Hole, Massachusetts 02543, U.S.A.

⁶Department of Biological Sciences, University of Toronto Scarborough, 1265 Military Trail, Toronto, Ontario, M1C 1A4, Canada

⁷Institute of Arctic Biology, 311 Irving 1 Building, University of Alaska Fairbanks, Fairbanks, Alaska 99775, U.S.A.

*Corresponding author's email: erik.hobbie@unh.edu

ABSTRACT

Arctic ground squirrels (*Urocitellus parryii*) rely primarily on dietary protein derived from plants to fuel gluconeogenesis during hibernation, yet fungal sporocarps may be an important, yet overlooked, protein source. Fungivory levels depend on sporocarp productivity, which varies with the dominant plant species and is higher on acidic than on non-acidic soils. To test whether these factors altered fungal consumption, we used stable isotopes to investigate arctic ground squirrel diets at two sites in northern Alaska, Toolik (primarily moist acidic tundra) and Atigun (primarily moist non-acidic tundra). Radiocarbon estimates can also indicate fungivory levels because ectomycorrhizal fungi assimilate soil-derived organic nitrogen whose ¹⁴C levels are higher than current photosynthesis. We measured radiocarbon in hair and $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in hair, feces, ectomycorrhizal sporocarps, graminoids, and dicots. Feces were higher in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ at Toolik than at Atigun, and fecal $\delta^{15}\text{N}$ increased in August at Toolik, coincident with sporocarp production and fungal spores in feces. Mixing models indicated that graminoids contributed 64%, dicots 35%, and sporocarps 1% to Atigun hair protein, whereas graminoids contributed 37%, dicots 16%, and sporocarps 47% to Toolik hair protein. Acidic soils appeared to correlate with higher sporocarp production and fungivory at Toolik than at Atigun. Atigun hair resembled atmospheric CO_2 in ¹⁴C, whereas Toolik hair had higher ¹⁴C, consistent with greater fungal consumption at Toolik. Late-season sporocarps may be a key protein source for some squirrels and may provide an integrated signal of the soil organic nitrogen assimilated by ectomycorrhizal fungi.

INTRODUCTION

Arctic ground squirrels (*Urocitellus parryii*), the northernmost mammalian hibernator in North America, have an annual season of aboveground ac-

tivity of up to 20 weeks (Sheriff et al., 2011), which includes the 6 to 10 week growing season for plants (Chapin et al., 1995). During this period, they add both fat and muscle in anticipation of overwintering (Boonstra et al., 2011; Sheriff et al., 2013). Arctic

ground squirrels are omnivorous and consume dicots, graminoids, fungal sporocarps, carrion, insects (Holowaychuk et al., 1966; Batzli and Sobaski, 1980; McLean, 1985; Hubbs and Boonstra, 1997; Karels and Boonstra, 2000), and even small mammals such as lemmings (Boonstra et al., 1990) and juvenile snowshoe hares (O'Donoghue, 1994) (Fig. 1). Fungal sporocarp consumption by arctic ground squirrels has not been quantitatively assessed, but appears to be primarily comprised of ectomycorrhizal fungi that rely on plants such as *Betula nana*, *Salix* spp., and *Dryas* spp. for plant photosynthate. Ectomycorrhizal sporocarps are common in the Low Arctic where most ground squirrels live. Techniques to assess fungivory have included direct observations, gut content analyses, and assessment of fungal spores in fecal pellets (Pastor et al., 1996). A fourth technique, measurements of stable isotope ratios in animals, has only been used twice (McIlwee and Johnson, 1998; Flaherty et al., 2010), although it has an advantage over other techniques of directly quantifying assimilation.

Stable isotope signatures are used extensively in dietary analyses of mammals with the most common being isotope ratios of carbon ($^{13}\text{C}/^{12}\text{C}$, expressed as $\delta^{13}\text{C}$) and nitrogen ($^{15}\text{N}/^{14}\text{N}$, expressed as $\delta^{15}\text{N}$). The general principles have been well-characterized (Gannes et al., 1997) and rely on comparing the isotopic relationship among specific tissues (e.g., hair, blood, bone collagen, bone apatite) and potential dietary sources. In the Arctic, $\delta^{15}\text{N}$ values are highest in ectomycorrhizal sporocarps, intermediate in monocots (graminoids), and lowest in dicots (Hobbie et al., 2009). Although these three

groups have not been compared for $\delta^{13}\text{C}$ in the Arctic, ectomycorrhizal sporocarps in temperate and boreal systems are generally higher in $\delta^{13}\text{C}$ than co-occurring plants (Hobbie, 2005). Evaluation of fecal stable isotopes can provide seasonal information on dietary assessment because fecal $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures reflect diet over a short time period (Salvarina et al., 2013). In contrast, hair $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures reflect diet during the period of hair growth and primarily reflect dietary protein (Sponheimer et al., 2003a; Froehle et al., 2010).

Another isotopic technique to assess diet is radiocarbon analysis (Beavan and Sparks, 1998; Hobbie et al., 2013). Many fungi assimilate organic nitrogen from the soil into their protein, and this organic nitrogen should have a $^{14}\text{C}:^{12}\text{C}$ ratio (expressed as $\Delta^{14}\text{C}$) that is older than the current-year photosynthate incorporated by plants during photosynthesis (Hobbie et al., 2013). The bomb spike signal from thermonuclear testing in atmospheric $^{14}\text{CO}_2$ levels peaked in 1963 and has declined rapidly since (Fig. 2). If fungal amino acids are incorporated into animal protein, the $\Delta^{14}\text{C}$ of hair from these fungivores should be older (higher) than that of atmospheric CO_2 , and $\Delta^{14}\text{C}$ should correlate with stable isotope indicators of fungivory. Uptake of ^{15}N -labeled amino acids by nonmycorrhizal sedges has been demonstrated at the Arctic Long Term Ecological Research site at Toolik Lake (McKane et al., 2002). We will therefore also assess whether the dominant sedge *Eriophorum* could be a second source for soil-derived organic nitrogen in diet by testing whether sedge protein has a radiocarbon signature indicating uptake of soil-derived amino acids.



FIGURE 1. Arctic ground squirrel consuming (left) a *Leccinum* sporocarp or (right) grasses. In the left picture, woody dicots (*Betula* at upper left, *Dryas* immediately below the sporocarp) are visible along with graminoids such as *Eriophorum* and *Carex*. Left photo from http://en.wikipedia.org/wiki/Arctic_ground_squirrel; right photo by Art Scheiber.




FIGURE 2. Atmospheric $\Delta^{14}\text{C}$ curve for carbon dioxide from 1974 to 2006, illustrating the steady decline in the bomb spike ^{14}C peak over that period. Data from Hua et al. (2013).

We investigated the relative importance of fungivory versus herbivory using several different techniques in two populations (Toolik and Atigun) of arctic ground squirrels located about 22 km apart in the foothills of the Brooks Range in northern Alaska. We measured $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ signatures of plants, sporocarps, and hair to determine consumption of graminoids, dicots, and fungal sporocarps, and then used mixing models on hair isotopes to determine consumption of graminoids, dicots, and fungal sporocarps in these two populations. Other factors that could influence diet or isotopic signatures were addressed using multiple regressions on hair and fecal isotopes. The short turnover time of fecal isotopes could allow a seasonal signal of fungivory to be detected because the fruiting of the sporocarps is restricted to the second half of the growing season. In addition, we measured radiocarbon in hair for possible signals of soil-derived organic nitrogen, and examined fecal samples from the two sites for the presence of fungal spores and other dietary items.

We hypothesized that these sporocarps may be a key protein-rich food source for arctic ground squirrels based on evidence of fungivory in other rodents (Johnson and Packer, 1967) and on one record of fungal consumption by arctic ground squirrels (Mayer, 1953). We also hypothesized that arctic ground squirrels from Toolik would have higher levels of fungivory than Atigun because of positive relationships between sporocarp production and low pH (Rao et al., 1997; Högberg et al., 2007; Soudzilovskaia et al., 2015). We addressed this hypothesis through the multiple methods mentioned above. Specifically, our overarching hypothesis predicted

that fungivory would be higher at Toolik than Atigun as indicated by (1) mixing models, (2) radiocarbon estimates, (3) hair and fecal $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ levels, and (4) fecal fungal spores presence.

MATERIALS AND METHODS

Site Description

Tissues of arctic ground squirrels and of potential dietary items (plants and fungal sporocarps) were collected at two sites in the northern foothills of the Brooks Range in Alaska. The Toolik site is located near the Toolik Field Station along the Dalton Highway, 254 km north of the Arctic Circle ($68^{\circ}37'40''\text{N}$, $149^{\circ}35'41''\text{W}$) at an elevation of 720 m. The Atigun site is located 22 km south of Toolik Field Station along the Dalton Highway in the Atigun river drainage ($68^{\circ}26'58''\text{N}$, $149^{\circ}21'43''\text{W}$) at an elevation of 804 m. Both sites are underlain by continuous permafrost with a depth of summer thaw that is generally 1–2 m where squirrels are present. However, the origin of the soils at the sites differ, with the acidic Toolik soil derived from glacial till, whereas the alkaline soil at the Atigun site was originally a wind-blown sand dune (Hamilton, 1979; Brown and Kreig, 1983). Vegetation at Atigun generally corresponds to non-acidic tundra, whereas vegetation at Toolik consists of moist-acidic or dry-acidic shrub tundra (Walker et al., 1981).

At the Toolik site, described in Walker and Maier (2008) and Hobbie et al. (2005), ground squirrel burrows are in well-drained soils along a 2 km strip of the east side of Toolik Lake, with some burrows in the field station area. The vegetation is dry-acidic and moist-acidic shrub tundra. Most burrows are in prostrate-shrub tundra with some burrows in erect-shrub tundra and barren areas around the Toolik station. Dominant species in the prostrate-shrub tundra include *Arctostaphylos alpina*, *Betula nana*, *Cassiope tetragona*, *Dryas integrifolia*, *Dryas octopetala*, *Empetrum nigrum*, *Ledum palustre* ssp. *decumbens*, *Salix phlebophylla*, *Salix rotundifolia*, *Salix pulchra*, *Vaccinium uliginosum*, *Vaccinium vitis-idaea*, graminoids, and mosses. Erect-shrub tundra is dominated by *Betula nana*, *Rubus chamaemorus*, *Salix pulchra*, graminoids, and mosses.

At the Atigun site, previously described in Walker et al. (1981) and Sheriff et al. (2011), ground squirrel burrows are in well-drained soil, primarily on the east side of the Dalton Highway. Vegetation cover data were collected during August 2008 at the Atigun study site (Shamhart, 2010). The non-acidic tundra is dominated by medium-height willows, *Dryas octopetala*, *Rhododendron lapponicum*, *Arctostaphylos alpina*, and *Vaccinium uliginosum*.

Collection of Plants and Fungi

Isotopic data on plants at Toolik were taken from previously archived values collected in 2000 (<<http://ecosystems.mbl.edu/ARC/terrest/biomass/index.shtml>>, file lgshttcn). We also compared these data against archived 2012 harvest data from Toolik (Hobbie and Moore, 2017) and observed that 2000 and 2012 data for graminoids were essentially identical (differing by 0.0‰ for $\delta^{13}\text{C}$ and by 0.2‰ for $\delta^{15}\text{N}$). Dicots were 0.9‰ lower in $\delta^{13}\text{C}$ and 0.5‰ higher in $\delta^{15}\text{N}$ in 2012 than in 2000. At Atigun, roots, leaves, and other plant components were collected in 2007 and 2008 from known or likely food items of arctic ground squirrels. These consisted of shrub leaves of *Arctostaphylos rubra*, *Dryas octopetala*, *Betula nana*, *Salix* spp., and *Vaccinium uliginosum*; forb leaves of *Oxytropis* sp., *Hedysarum* sp., and *Artemisia* sp.; berries of *Arctostaphylos rubra*, *Empetrum nigrans*, *Shepherdia canadensis*, and *Vaccinium uliginosum*; and roots of *Dryas*, *Epilobium*, *Hedysarum*, and *Oxytropis*. The graminoids *Carex bigelowii* and *Eriophorum angustifolium* were collected at both Toolik and Atigun, with the addition of *Eriophorum vaginatum* at Toolik and unidentified graminoids at Atigun. The dicots at Toolik consisted of *Andromeda polifolia*, *Betula nana*, *Cassiope tetragona*, *Dryas integrifolia*, *Ledum palustre*, *Rubus chamaemorus*, *Vaccinium uliginosum*, and *Vaccinium vitis-idaea*. Fungal sporocarps were opportunistically collected in 2008 in both the Atigun River drainage and the Toolik Lake study areas but were not necessarily located within burrow areas. Plants and sporocarps were dried in a food dehydrator at Toolik Field Station and ground at the University of New Hampshire.

Hair and Feces Collection

Arctic ground squirrels were trapped using Tomahawk live traps baited with carrots during the spring, summer, and fall of 2007, 2008, 2011, and 2012 at the Toolik and Atigun study sites. Traps were set between 8:00 and 10:00 a.m. and checked within 2 h. Captured squirrels were brought to Toolik Field Station and anesthetized using isoflurane mixed with oxygen. The animals were then weighed, sexed, measured, and tagged with Monel #1 tags (National Band and Tag Co., Newport, Kentucky, U.S.A.) in both ears (first capture only). Hair was collected from 7 June to 8 September 2007, from 18 April to 30 August 2008, and from 5 April to 8 August 2011. Hair samples were collected by shaving a 4 cm² patch on the abdomen while under anesthesia. In 2008 and 2012, a fresh fecal sample was collected either in the lab prior to anesthetization or in the field from below the trap, which had been previously cleared. Fecal samples were placed on ice for

4–6 h prior to being stored at –20 °C. Squirrels were trapped from 24 to 30 August 2008 and from 9 May to 8 September 2012. The animals were held overnight and returned to the burrows where they were captured. All live animal procedures were approved by the Institutional Animal Care and Use Committees of the University of Fairbanks and the University of New Hampshire (UAF IACUC #06-25 and #08-59, UNH IACUC #080904A).

Microscopic Analysis of Feces

Freeze-dried fecal samples were weighed, and deionized water added at 110 $\mu\text{L mg}^{-1}$ fecal material to normalize dilution. Samples were ground to a slurry by micropestle and frozen at –20 °C for 24 h. For microscopic examination, samples were thawed and 50 μL dropped onto a slide. A drop of Melzer's reagent was added to the slurry to detect amyloid reactions, and a 24 \times 24 mm cover slip was placed on the slide.

Each sample was completely scanned at 100 \times magnification in approximately 16 objective passes across the slide, progressing from the top to the bottom. To identify potential dietary items or spores, magnification was increased to 400 \times or 1000 \times . Spores or spore-like structures were categorized into morphological groups, based on their size, shape, and surface features (Castellano et al., 1989; Colgan et al., 1997). The quantity of each morphological type and other features of interest were documented for each sample, and for most samples the relative dominance of plant, fungal, and insect remains was recorded. Spore counts were used as a measure of relative abundance on a given slide (not between slides) because, although weight-appropriate amounts of distilled water were added to each sample, the feces varied widely in their consistency, and some were quite concentrated in fecal material whereas others appeared much more dilute.

Stable Isotope Analysis

At the University of New Hampshire Stable Isotope Lab, foliage, sporocarps, and fecal samples were ground to a powder, and hair was clipped into small pieces. Foliage, sporocarps, hair, and fecal samples were weighed to 1 μg precision into tin cups. Hair samples were 1.0–1.5 mg, and other samples were 4–5 mg. The $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values were determined using a Costech 4010 Elemental Analyzer and a Delta XP Mass Spectrometer with a precision on duplicate samples of 0.2‰.

Lab standards of sporocarps, tuna muscle, apple leaves (NIST1515), and pine needles (NIST1575a) were included with each run. At least 15 standards and two sample duplicates were included with each run of 28

samples. All data were normalized and corrected for drift and linearity.

Radiocarbon Analysis

Five hair samples of individual adults at Toolik and two at Atigun were analyzed for ^{14}C content in animals collected in 2008. To provide a well-integrated site average, an additional six samples from Toolik and five samples from Atigun that consisted of 3–4 individuals each were also analyzed. These samples were from either 2008 or 2011. Hair was converted to graphite targets at the University of Florida and then analyzed at the Keck–Carbon Cycle AMS facility at the University of California at Irvine.

Isolating *Eriophorum* Protein for ^{14}C Analyses

We isolated *Eriophorum* protein from archived samples and compared its $\Delta^{14}\text{C}$ values against atmospheric CO_2 , because ^{14}C differences from atmospheric CO_2 indicate some assimilation of soil-derived organic nitrogen. Archived foliage collected at Toolik Field Station in 1986, 1995, and 2000 from the Ecosystems Center, Marine Biological Laboratory (U.S.A.) was used. Isolation of protein and structural carbon followed Hobbie et al. (2013). Ground foliage and the barley standard FIRI-G were initially extracted three times with hexanes (30 min at 70 °C) to remove non-polar compounds, followed by three extractions with 80% ethanol (30 min at 100 °C) to remove soluble polar compounds, mainly carbohydrates. The remaining solid was hydrolyzed once with 6 M hydrochloric acid at 110 °C for 24 h. Hydrolysate was filtered, and soluble compounds were purified to amino acids using cation-exchange resin. The amino acids were eluted and then dried down as the protein fraction. The remaining, insoluble material following acid hydrolysis was extracted at 100 °C for 2 h with 0.2 M sodium hydroxide to remove residual proteins and the remaining solid material was considered the structural fraction. The protein and structural fractions were then sent to the University of Florida, converted to graphite targets, and analyzed for ^{14}C at the Keck–Carbon Cycle AMS facility at the University of California at Irvine in 2013.

Estimating Diet Composition from Isotopic Signatures

We used two statistical approaches to examine the controls over dietary patterns of arctic ground squirrels. In one approach, isotopic mixing models developed over the past 15 years were used to partition the three sources of graminoids, dicots, and sporocarps, using isotopic information on hair (Simmens et al., 2009).

Potential food sources were classified into three discrete categories: graminoids, dicots, and fungi. Isotopic signatures for each category were then estimated by averaging potential food sources within each category. The Bayesian mixing model MixSIAR, as implemented within MixSIAR GUI 1.0 (Stock and Semmens, 2013), was used to identify the probability distributions of the contributions of these three food resources in the diet of Toolik and Atigun squirrels. MixSIAR estimates the probability distributions of the contributions of each source to a mixture and includes the uncertainty associated with multiple sources, isotope signatures, and tissue-diet discrimination (Simmens et al., 2009). The estimated median contribution is given for comparative purposes. The input parameters included in the mixing model were the isotopic values of consumers (hair), isotopic values of potential food sources (measured in this study), and tissue-enrichment factors (change in isotope signature from diet to tissue) and their associated standard deviations. Rat feeding trials in Kurlle et al. (2014) provided a hair-diet ^{13}C enrichment factor of $3.3 \pm 0.4\text{‰}$. This value was nearly identical to the average ^{13}C enrichment of $3.2 \pm 0.3\text{‰}$ for hair relative to diets of five mammalian herbivores (Sponheimer et al., 2003a) and ^{13}C enrichment of $3.6 \pm 0.3\text{‰}$ in bats (Salvarina et al., 2013). We used a value of $3.3 \pm 1.0\text{‰}$ for ^{13}C enrichment in our MixSIAR modeling. A ^{15}N enrichment for hair relative to a diet of $1.9 \pm 0.1\text{‰}$ was calculated for rats from five different diets in Caut et al. (2008) and a value of $3.3 \pm 1.0\text{‰}$ estimated from Sponheimer et al. (2003b), from five mammalian herbivores. We used a value for ^{15}N enrichment of $1.9 \pm 1.0\text{‰}$, because the digestive systems of arctic ground squirrels are presumably more similar to those of rats than to the herbivores of the Sponheimer study. The Gelman–Rubin, Heidelberger–Welch, and Geweke diagnostic tests were used to confirm that the model had converged (Hata et al., 2015). Markov Chain Monte Carlo parameters were set as follows: number of chains = 3, chain length = 100,000, burn in = 50,000, thin = 50.

Statistical Analyses on Hair and Feces

In another approach, we used mixed multiple regression models to examine how isotopic patterns in hair or feces correlated with year, season, sex, age class, or chemical composition. Patterns in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ signatures in hair and feces were analyzed using stepwise multiple regression in JMP (SAS Institute, Middleton, Massachusetts, U.S.A.). The $\delta^{15}\text{N}_{\text{hair}}$ and $\delta^{13}\text{C}_{\text{hair}}$ signatures were analyzed as dependent variables in separate regression models, and each model included sex, age class, year, and site as independent variables. Categories for sex were male and female; castrated males and animals of unknown sex were excluded

from regression analyses. Age classes were adults, yearlings, juveniles, and animals of unknown age. Years were 2007, 2008, or 2011, and sites were Toolik and Atigun.

The carbon and nitrogen isotope signatures of fecal samples were analyzed together for 2008 and 2012 using stepwise multiple regressions for $\delta^{15}\text{N}_{\text{feces}}$ and $\delta^{13}\text{C}_{\text{feces}}$. Independent variables for fecal analyses were site, year, date (day of year divided by 100), $\log_e \text{C/N}$, the interaction of date and site, and either $\delta^{15}\text{N}_{\text{feces}}$ (if $\delta^{13}\text{C}_{\text{feces}}$ was the dependent variable) or $\delta^{13}\text{C}_{\text{feces}}$ (if $\delta^{15}\text{N}_{\text{feces}}$ was the dependent variable). To account for possible non-linear relationships with the date, date was included as up to a third-order polynomial. Because other compound classes such as lipids and carbohydrates are higher in C/N and lower in $\delta^{13}\text{C}$ than protein, their possible influence on $\delta^{13}\text{C}_{\text{feces}}$ was accounted for by including C/N as an explanatory variable in feces. We examined the potential for seasonality in fecal isotopes by including the date of fecal collection as a continuous variable. In stepwise regression analyses, the model formulations that minimized the values of AICc (Akaike Information Criteria with a correction for sample size) were selected (Carleton et al., 2008). In addition, we also compared $\delta^{15}\text{N}$, $\delta^{13}\text{C}$, and C/N of fecal samples between Atigun and Toolik using *t*-tests.

RESULTS

Estimating Fungivory and Herbivory from Stable Isotope Patterns in Hair and Food Sources

Potential food sources at Toolik and at Atigun were separated into the three categories of fungi, graminoids, and dicots. For each site, stable isotope values within each category were averaged and used in subsequent analyses

(Table 1). Isotopic data by taxa on edible fungi are given in Table 1, and they averaged $-25.2 \pm 0.1\text{‰}$ for $\delta^{13}\text{C}$ and $4.0 \pm 0.1\text{‰}$ for $\delta^{15}\text{N}$ at Toolik and $-25.3 \pm 0.1\text{‰}$ for $\delta^{13}\text{C}$ and $6.4 \pm 1.0\text{‰}$ for $\delta^{15}\text{N}$ at Atigun. Isotopic data for plants by taxa at Toolik separated clearly into graminoid and dicot groupings (Fig. 3); average $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for graminoids were $-25.6 \pm 0.3\text{‰}$ and $1.7 \pm 0.4\text{‰}$, respectively, whereas average $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for dicots were $-28.2 \pm 0.4\text{‰}$ and $-4.3 \pm 1.0\text{‰}$, respectively. Dicots at Atigun included *Arctostaphylos rubra*, *Artemisia* sp., *Betula nana*, *Dryas octopetala*, *Hedysarum* sp., *Oxytropis* sp., *Salix* spp., *Shepherdia canadensis*, and *Vaccinium uliginosum*. Isotopic data for plants at Atigun are in online Appendix Table A1.

At Toolik, the hair $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures averaged $-23.52 \pm 0.07\text{‰}$ and $4.53 \pm 0.13\text{‰}$ ($n = 53$), respectively, whereas at Atigun the average signatures were $-24.66 \pm 0.05\text{‰}$ and $0.95 \pm 0.08\text{‰}$ ($n = 108$), respectively. Results for individuals are shown in Figure 4 by site and year. The C:N ratio of hair averaged 2.906 ± 0.007 at Toolik and 2.899 ± 0.006 at Atigun (*t*-test, $P = 0.494$).

In regression models, site was the dominant factor controlling both $\delta^{13}\text{C}$ (88% of variance) and $\delta^{15}\text{N}$ (96% of variance), with Toolik higher than Atigun in both parameters. Age class explained 10% of variance in $\delta^{13}\text{C}$, with adults ($n = 77$) averaging 0.35‰ higher than yearlings ($n = 15$), juveniles ($n = 48$), or animals of unknown age ($n = 21$). Sex explained 1% of variance in $\delta^{15}\text{N}$, with males ($n = 84$) 0.35‰ higher than females ($n = 77$). Year explained 2% of $\delta^{15}\text{N}$ variance, with 2011 ($n = 68$) 0.84‰ higher than 2007 ($n = 7$) and 0.4‰ higher than 2008 ($n = 84$) (Table 2). Given the primary importance of site in controlling stable isotope patterns, we grouped all age classes from both sexes when using isotope mixing models to explore diet.

TABLE 1

Isotopic values of graminoids, dicots, and taxa of edible sporocarps by taxa at Atigun and Toolik. Data are means \pm se.

Taxon	Atigun			Toolik		
	$\delta^{13}\text{C} \pm \text{se} (\text{‰})$	$\delta^{15}\text{N} \pm \text{se} (\text{‰})$	<i>n</i>	$\delta^{13}\text{C} \pm \text{se} (\text{‰})$	$\delta^{15}\text{N} \pm \text{se} (\text{‰})$	<i>n</i>
Boletaceae	-25.3 ± 0.1	8.7 ± 0.3	4	-25.5 ± 0.0	6.3 ± 0.1	2
<i>Boletus</i>	-24.9 ± 0.1	9.8 ± 0.1	2	-22.7	2.0	1
<i>Lecinum</i>	-25.0 ± 0.2	6.6 ± 2.4	2	-25.4 ± 0.3	6.5 ± 0.8	6
<i>Laccaria</i>	-25.2 ± 0.1	2.8 ± 1.9	2	-25.3 ± 0.6	1.3 ± 0.7	5
<i>Lactarius</i>	-25.7 ± 0.1	4.9 ± 0.5	7	-25.4 ± 0.2	3.0 ± 0.8	3
<i>Russula</i>	-25.6 ± 0.3	5.8 ± 1.3	6	-26.8	4.9	1
average	-25.3 ± 0.1	6.4 ± 1.0	—	-25.2 ± 0.1	4.0 ± 0.1	
Graminoids	-25.8 ± 0.7	1.2 ± 0.4	3	-25.6 ± 0.3	1.7 ± 0.4	3
Dicots	-28.2 ± 0.2	-3.8 ± 0.6	11	-28.0 ± 0.4	-4.3 ± 1.0	11




FIGURE 3. Carbon and nitrogen isotopes (\pm se) in photosynthetic tissues of Toolik plants sampled in 2000 from moist acidic and moist non-acidic tussock tundra. Three letter abbreviations indicate genus and species, for the dicots *Andromeda polifolia* (And), *Betula nana* (Bet), *Cassiope tetragona* (Cas), *Dryas integrifolia* (Dry), *Ledum palustre* (Led), *Polygonum bistorta* (Pol), *Rubus chamaemorus* (Rub), *Salix reticulata* (Sal), *Vaccinium uliginosum* (Vcu), and *Vaccinium vitis-idaea* (Vcv); and the graminoids *Carex bigelowii* (Car), *Equisetum* spp. (Equ), *Eriophorum angustifolium* (Era), and *Eriophorum vaginatum* (Erv). Dicots are in black, graminoids are in gray. Data are at <http://arc-lter.ecosystems.mbl.edu/2000lgshttcn>.




FIGURE 4. Carbon and nitrogen isotope signatures in arctic ground squirrel hair by year at Toolik (s, 2007; w, 2008; o, 2011) and Atigun (+, 2007; <, 2008; >, 2011).

Dietary patterns were estimated using the MixSIAR model. Dietary sources were aggregated into three categories of graminoids, dicots, and fungi because the different taxa within these sources clustered in similar isotopic space.

Average (\pm sd) contribution to hair from dicots, graminoids, and fungi was 0.134 ± 0.043 , 0.366 ± 0.085 , and 0.496 ± 0.065 at Toolik and 0.366 ± 0.046 , 0.593 ± 0.056 , and 0.041 ± 0.025 at Atigun, respectively (Table 3, complete MixSIAR results are in online Appendix Table A3).

Radiocarbon in Hair and *Eriophorum*

Toolik hair averaged 27‰ higher in $\Delta^{14}\text{C}$ than Atigun hair ($66.9 \pm 6.3\text{‰}$, $n = 12$ versus $40.2 \pm 3.2\text{‰}$, $n = 7$, t -test assuming unequal variance, $p = 0.002$). Toolik hair averaged somewhat higher than atmospheric CO_2 in $\Delta^{14}\text{C}$ and was more variable (F -test on $\Delta^{14}\text{C}$, $p = 0.027$), whereas Atigun hair was slightly below atmospheric CO_2 (Fig. 5). *Eriophorum* protein was slightly higher in $\Delta^{14}\text{C}$ than structural carbon, whereas *Eriophorum* structural carbon resembled atmospheric CO_2 in $\Delta^{14}\text{C}$ (Table 4). Extracts of the barley standard resembled the known bulk values for the barley of 102.5‰, with barley protein at 95.9‰ and 98.4‰ and barley structural carbon at 95.9‰ and 101.0‰.

Stable Isotope Patterns in Feces

Means of $\delta^{15}\text{N}$, $\delta^{13}\text{C}$, and $\log_e \text{C/N}$ at the two sites were compared for both 2008 and 2012 in Table 5, and regression models for samples are given in Table 6. In 2008, feces were higher in $\delta^{15}\text{N}$ at Toolik ($n = 3$) than at Atigun ($n = 31$) and were similar in $\delta^{13}\text{C}$ and in C/N at the two sites. In regression models, location and the cube of the date were significant factors for $\delta^{13}\text{C}$, whereas site, year, date, the square of the date, the $\log_e \text{C/N}$, and the interaction of location and date were significant factors for $\delta^{15}\text{N}$ (Table 6). The positive correlation of date and the square of date, and the interaction of date and the Toolik site reflected increased fecal $\delta^{15}\text{N}$ toward the end of the growing season at Toolik but not at Atigun (Fig. 6).

The $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ increased in Toolik in the fall. This paralleled a decrease in C/N for Toolik samples but not Atigun samples collected later in the growing season, presumably reflecting a shift in diet at Toolik to lower C/N sources such as fungi.

In a stepwise regression model, site, the cube of date, and the interaction of site and date significantly affected fecal C/N (Table 7; complete regression results in online Appendix Table A5). The cube of date accounted for 72% of explained variance, and location times date accounted for 21%. Atigun C/N was about 2 lower than Toolik C/N .

Microscopic Analysis of Feces

We examined 21 fecal samples at Toolik and 22 at Atigun, all from 2012. Collections were equally

TABLE 2

Stepwise multiple regression analyses of arctic ground squirrel hair $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ as correlated with site, sex, year, and age class. The % variance explained (%Var) for each of the parameters is also given. Adjusted r^2 for $\delta^{13}\text{C}$ was 0.570 and adjusted r^2 for $\delta^{15}\text{N}$ was 0.824 ($P < 0.001$ for both, $n = 161$). Data are means \pm se. For age classes, Y = yearling, J = juvenile, A = adult, U = unknown age; for years, 07 = 2007, 08 = 2008, and 11 = 2011. Full regression analyses are given in online Appendix Table A2.

Parameter	%Var	$\delta^{13}\text{C}$ regression		%Var	$\delta^{15}\text{N}$ regression	
		Value \pm se	P		Value \pm se	P
Intercept	—	-24.10 ± 0.04	<0.001	—	2.59 ± 0.08	<0.001
Site (Atigun)	88.4	-0.55 ± 0.04	<0.001	96.4	-1.88 ± 0.07	<0.001
Age Class (Y&U&J-A)	10.2	-0.17 ± 0.04	<0.001	—	—	—
Age Class (U&J-Y&A)	—	—	—	0.4	0.11 ± 0.07	0.099
Year (11-08&07)	1.4	-0.07 ± 0.04	0.096	—	—	—
Year (07-11)	—	—	—	1.9	-0.42 ± 0.11	<0.001
Sex (Female)	—	—	—	1.1	-0.19 ± 0.06	0.004

TABLE 3

Median assimilation by arctic ground squirrels at Toolik versus Atigun with 95% Bayesian credible intervals for dicots, graminoids, and fungi. This was estimated from MixSIAR analyses on $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of these three sources. Complete MixSIAR results are in online Appendix Table A3.

Diet source	Toolik	Atigun
Dicots	0.134 (0.056–0.226)	0.368 (0.275–0.453)
Graminoids	0.366 (0.207–0.540)	0.593 (0.486–0.701)
Fungi	0.496 (0.361–0.620)	0.040 (0.001–0.092)




FIGURE 5. Radiocarbon values for arctic ground squirrel hair, with individual Toolik and Atigun samples indicated with a lowercase “t” (6 samples) or “a” (2 samples). Pooled samples from 3 to 4 individuals each are indicated with an uppercase “T” (6 samples) or “A” (5 samples). Northern hemisphere (NH) atmospheric $^{14}\text{CO}_2$ levels are also plotted by year, data from Hua et al. (2013).

divided among early May, mid-July, and early September. Complete results are given in online Appendix Table A6. At Toolik, dominant remains included plants (6 samples), insects (3 samples), fungi (3 samples), and a mix of plants and insects (5 samples). At Atigun, dominant remains included insects (12 samples), plants (2 samples), and a mix of plants and insects (4 samples).

Fungal spores were identified by appearance. Four fungal spore types were readily identifiable to family level, but many of the structures were of unfamiliar morphologies and could not be confidently identified. Two types of segmented teliospores were observed, as well as a smooth ellipsoidal ascomycete spore (perhaps *Geopora*) and two distinctive larger globose spore types that appeared to be Glomalean.

Fungal spores were noted primarily in July and September collections from Toolik. We identified three basidiomycete families by spore morphology, consisting of four distinct types of boletoid spores (*Boletaceae*), three types of cortinarioid spores (*Cortinariaceae*), and one or two types of russuloid spores (*Russulaceae*). Only one ascomycetous spore was observed across all samples. Large intact mycelial wefts were also present in several samples. These clearly indicated fungivory but their identities could not be ascertained.

Although variability in sample dilution precludes statistical analysis of samples, feces collected at the Toolik site generally had more spore content than those from Atigun, and samples collected in July and September generally had more spores than those collected in May. No patterns were apparent between sex or age and spore content.

TABLE 4

Eriophorum radiocarbon in structural and protein component and estimated ages of those components.

Year	Structural $\Delta^{14}\text{C}$ \pm error (‰)	Age (years)	Protein $\Delta^{14}\text{C}$ \pm error (‰)	Age (years)	Atmospheric $\Delta^{14}\text{CO}_2$ (‰)
1986	215.1 \pm 2.3	1–2	235.2 \pm 2.7	3	191.1
1995	110.3 \pm 2.5	0	128.7 \pm 2.3	1	115.5
2000	89.8 \pm 2.2	0	91 \pm 2.3	0	87

TABLE 5

$\delta^{15}\text{N}$, $\delta^{13}\text{C}$, and the \log_e of C/N of arctic ground squirrel feces collected at Atigun and Toolik in 2008 ($n = 31$ and 3, respectively) and 2012 ($n = 51$ and 28, respectively). Samples in 2008 were collected from 24 to 31 August, and samples in 2012 were collected from 9 May to 8 September. The sites were compared using t -tests. Data are means \pm se.

Parameter	Year	Atigun	Toolik	t -test
$\delta^{15}\text{N}$ (‰)	2008	0.27 \pm 0.23	6.39 \pm 0.93	0.017
$\delta^{13}\text{C}$ (‰)	2008	−28.04 \pm 0.13	−28.07 \pm 0.26	0.929
\log_e C/N	2008	2.74 \pm 0.04	2.59 \pm 0.11	0.292
$\delta^{15}\text{N}$ (‰)	2012	0.13 \pm 0.12	2.51 \pm 0.33	<0.001
$\delta^{13}\text{C}$ (‰)	2012	−28.18 \pm 0.08	−27.33 \pm 0.08	<0.001
\log_e C/N	2012	2.72 \pm 0.05	2.81 \pm 0.08	0.351

TABLE 6

Effects of site, year, date of sampling, \log_e of C/N, the interaction of site and date, and the interaction of date and \log_e of C/N on $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of arctic ground squirrel feces collected in 2008 and 2012 ($n = 109$). For $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$, adjusted $r^2 = 0.281$ and 0.700 , respectively, $p < 0.001$ for both. %Var is the % of variance explained by a specific factor. Date unit is 100 days, e.g., July 12 would be 1.94. Stepwise regression tests to minimize AICc are in online Appendix Table A4.

$\delta^{13}\text{C}$				$\delta^{15}\text{N}$			
Term	%Var	Value \pm se (‰)	P	Term	%Var	Value \pm se (‰)	P
Intercept	—	−27.75 \pm 0.06	<0.001	Intercept	—	1.82 \pm 1.54	0.240
Site (Atigun) ^a	81.8	−0.38 \pm 0.06	<0.001	Site (Atigun) ^a	69.6	−1.62 \pm 0.12	<0.001
^b (Date − 1.94) ³	18.2	−1.33 \pm 0.45	0.004	Year (2008) ^c	6.3	0.54 \pm 0.13	<0.001
				Date	6.7	1.51 \pm 0.36	<0.001
				(Date − 1.94) ²	9.9	3.90 \pm 0.77	<0.001
				\log_e C/N	3.9	−1.21 \pm 0.38	0.002
				Site ^a \times Date ^c	3.6	−1.00 \pm 0.33	0.003

^aAtigun would have the value added, Toolik would have the value subtracted.

^bMean of date values was 1.94; the cube of the date was significant for $\delta^{13}\text{C}$ and the square of the date was significant for $\delta^{15}\text{N}$.

^c2008 would have the value added, 2012 would have the value subtracted.

DISCUSSION

Fecal Isotopes and Diet

Stable isotope ratios in feces reflect recent food consumption and accordingly provide information complementary to the long-term and integrated record provided by ratios in hair (Flaherty et al., 2010). Whereas isotopes in feces reflect the overall recent diet, those in hair pri-

marily reflect dietary protein when the hair grew, because metabolic routing means that much of animal protein is derived directly from dietary amino acids (Hobbie, 2017). The few fecal samples from arctic ground squirrels at Toolik in 2008 were higher in $\delta^{15}\text{N}$ than the Atigun samples (Table 5), corroborating our evidence of greater fungal consumption by Toolik animals. The larger data set of 2012 supported this interpretation, because there was a





FIGURE 6. Fecal isotope patterns in 2012 plotted against day of year of collection. Toolik samples are indicated by clear circles and Atigun samples by filled triangles. (A) $\delta^{15}\text{N}$; (B) $\delta^{13}\text{C}$; (C) C/N.

significant interaction of date and site for $\delta^{15}\text{N}$ (Table 6), and late-season samples from Toolik were generally high in $\delta^{15}\text{N}$ (Table 6 and Fig. 7), low in C/N (Table 7), and also had abundant fungal spores, unlike early-season samples (online Appendix Table A6). Fungal sporocarps are low in C/N (Hasselquist and Höglberg, 2014) relative to arctic plants at Toolik.

Insect parts in some of the feces from both locations indicated that we did not have isotopic data on all potential food sources. Insect consumption has yet to be quantified in arctic ground squirrels, but small percentages of insects by volume have been recorded in related species of ground squirrels such as in the golden-mantled ground squirrel (*Callospermophilus lateralis*) in coniferous forests (Tevis, 1953; percentages were 1% in spring and fall, rising to 16% in summer) and the desert-dwelling antelope squirrel (*Ammospermophilus leucurus*) (~10%; Bradley, 1968).

Arthropod consumption was not modeled in MixSIAR because we lacked data on the appropriate isotopic end members for this potential food source. Published work indicated that insects are generally higher in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ than plants (Ikeda et al., 2010; Hyodo, 2015).

However, ongoing work in other biomes indicates that arthropods are intermediate between plants and fungi in isotopic composition (personal communication, R. Stephens), and the dominant dietary component in feces was not a factor in the AICc-selected regression analyses on fecal $\delta^{15}\text{N}$, $\delta^{13}\text{C}$, and C/N (online Appendix Table A7). Because both arthropods and fungi appear higher in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ than plants, including arthropods in mixing models would necessarily reduce the estimated dietary contribution of fungi. The maximum volume of animal matter (5%) recorded in stomachs of




FIGURE 7. The relative effect of date and site on fecal $\delta^{15}\text{N}$ at Atigun and Toolik, illustrating the greater seasonality in the $\delta^{15}\text{N}$ signal at Toolik. The graph was computed by combining the effect on fecal $\delta^{15}\text{N}$ of site, date, site \times date, and date squared as given in the multiple regression analysis in Table 6.

TABLE 7

Effects of site, date of sampling, and the interaction of date and site on C/N of arctic ground squirrel feces collected in 2012. Date unit is 100 days. The adjusted r^2 for C/N is 0.746 ($P < 0.001$, $n = 75$).

Term	%Variance	Parameter \pm se	P
Intercept	—	32.96 \pm 3.02	<0.001
Site (Atigun)	10.8	-1.37 \pm 0.64	<0.001
Date (100 days)	73.4	-8.64 \pm 1.54	<0.001
Date ¹ \times Site ²	15.8	4.00 \pm 1.54	<0.001

¹Date - 1.885 (mean of date values was 1.88).

²Atigun.

arctic ground squirrels (Batzli and Sobaski, 1980) suggested that arthropods are an unlikely dominant protein source. In addition, investigators in the Toolik area have carried out intense live-trapping studies on these sites for multiple years to examine life history constraints and have not documented carnivory. Thus, mammal-derived protein is also unlikely to explain our observations.

Interpreting Stable Isotope Patterns

Several of the independent variables in addition to site correlated significantly with $\delta^{13}\text{C}$. In the regression analysis (Table 2), the calculated 0.35‰ ^{13}C enrichment of adults relative to other animals presumably reflected the reported ^{13}C depletion in milk of nursing mothers relative to the diet caused by the high content of ^{13}C -depleted milk lipids (Miller et al., 2008).

Hair samples from Toolik were sufficiently high in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ to suggest a relatively large contribution of sporocarps to dietary protein, given the high $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ in sporocarps (Table 1) relative to graminoids and dicots (Fig. 3). In contrast, Atigun hair samples appeared to incorporate little fungal protein. These results were supported by the MixSIAR analyses that indicated 30–62% contribution of fungi at Toolik and 0–4% contribution of fungi at Atigun (Table 3).

The half-life of formation of hair carbon in rodents is 40–50 days (Tieszen et al., 1983; Caut et al., 2008), and the carbon of course persists for longer. Arctic ground squirrels molt in the spring and grow underwool in late summer, but it was not possible to determine when hair had molted. Hair sampled in April and May most likely reflected carbon and nitrogen that were assimilated in the prior season from the late summer molt (August–September), with hair sampled from the beginning of July reflecting current year diet from the early molt (May–July). Males molt in late May and June, and females in June and July after they wean their young (Butterworth, 1958), thus some sexual differences in molting time may influence isotopic patterns in hair, such as

the small (0.4‰) depletion in ^{15}N in females relative to males (Table 2).

Data on stomach and cheek pouch contents of arctic ground squirrels near Atqasuk (70°29'N, 157°25'W) in northern Alaska from Batzli and Sobaski (1980) provide detailed direct measurements of arctic ground squirrel diets for July and August to compare against the current estimates. In their study, herbaceous dicots (47%), seeds (18%), monocots (8.5%), and *Equisetum* (8%) were the primary food sources. Fungal consumption was not mentioned, but the animal matter (apparently insects) contribution to stomach contents was 4.9% in May–June, 0.4% in July, and 1.3% in August. In parallel choice experiments of individual plant taxa, highest average consumption of offered foods was for *Equisetum* (75%, one taxon), followed by forbs (62 \pm 22%, 34 species), deciduous shrubs (46 \pm 17%, 8 species), monocots (17 \pm 16%, 10 species), and evergreen shrubs (13 \pm 11%, 5 species) (values \pm sd) (Batzli and Sobaski, 1980). The Atqasuk site was similar in some respects to Atigun, with most burrows located in sandy areas or river bluffs, and the species compositions of forbs and grasses indicate non-acidic tundra at both sites (personal communication, Donald Walker). However, the diet estimates from stomach contents (Atqasuk) and from isotopes in hair (Atigun) are rather different, but that is hardly surprising given probable wide differences among plants in protein content as well as inherent site differences between the northern Alaskan coastal plain (Atqasuk) and the Brooks Range foothills (Atigun).

Interpreting Radiocarbon Results

Toolik hair averaged 27‰ higher in $\Delta^{14}\text{C}$ than Atigun hair, indicating that Toolik squirrels incorporated older carbon than Atigun squirrels. We interpret this difference to reflect the incorporation into food sources of old, soil-derived amino acids that were first synthesized when ^{14}C content of the atmosphere was considerably higher. Because arctic ground squirrels

primarily consume plant matter (Batzli and Sobaski, 1980), we considered the possibility that protein from non-mycorrhizal plants may themselves preserve a ^{14}C signal of soil organic nitrogen. The limited data on *Eriophorum* radiocarbon indicated that protein carbon of some plants can be slightly older than structural carbon, by 0–3 years (Table 4). However, *Eriophorum* is not a significant dietary source for soil-derived organic nitrogen, as these ages are insufficient to account for the ^{14}C enrichments seen in arctic ground squirrels at Toolik. This indicated that Toolik squirrels incorporated some old organic nitrogen into hair, presumably via fungi, and that the low ^{14}C levels in Atigun squirrels reflected little fungal consumption. These results pointed to the probable persistence of organic nitrogen compounds in the soil, their incorporation into ectomycorrhizal fungi (Tibbett et al., 1998; Clemmensen et al., 2008; Hobbie et al., 2013), and the subsequent assimilation of that organic nitrogen by fungivorous mammals.

Implications

The large differences in isotope values between the two arctic ground squirrel locations analyzed in this study indicate that the two populations choose food based on availability. This feeding strategy should allow arctic ground squirrels to adapt to changing conditions despite the fact that forbs, one of their important food sources (Holowaychuk et al., 1966; Batzli and Sobaski, 1980), are expected to decrease across the Arctic with climate change (Chapin et al., 1995). However, the current effects of climate change specifically at Toolik show a warming in the fall but later snowmelt in the spring (Cherry et al., 2014). Expanding ectomycorrhizal shrub habitat in the Arctic (Myers-Smith et al., 2011) and at Toolik may increase arctic ground squirrel fungivory as ectomycorrhizal sporocarps become more common.

Ectomycorrhizal plants at Toolik and Atigun could provide sugars for the ^{13}C -enriched ectomycorrhizal sporocarps that may be consumed by arctic ground squirrels. However, the proportion of ectomycorrhizal plants appeared to differ; heath, shrub, and tussock vegetation types at Toolik had more *Betula* and less *Salix* (Shaver and Chapin, 1991), whereas vegetational quadrats at Atigun indicated between 1% and 20% *Dryas*, 4% and 16% *Salix*, 1% and 9% *Arctostaphylos*, and <1% *Betula* (Shamhart, 2010). The lesser proportion of *Betula* may translate into less sporocarp production at Atigun than at Toolik, although the substrate itself (glacial till at Toolik, dune sand at Atigun) may be the controlling factor. No data are available on the relative sporocarp productivity of *Betula nana* systems versus *Salix* spp.

systems in the Arctic, or on the influence of substrate on sporocarp productivity. However, ectomycorrhizal sporocarp productivity decreases with increasing pH in forest communities (Rao et al., 1997; Höglberg et al., 2007), and soil pH at Atigun should be higher than at Toolik because of the contribution of eroding limestone to Atigun (Walker et al., 1981). The proportion of fungi in diets of arctic ground squirrels probably integrates sporocarp productivity. Protein assimilation from sporocarps may be particularly important because hibernation in arctic ground squirrels requires gluconeogenesis from lean muscle mass, unlike in other squirrel hibernators (Boonstra et al., 2011). Thus, protein demand is likely higher than in other squirrel species. The clear elevation of $\Delta^{14}\text{C}$ signatures of arctic ground squirrels at Toolik relative to Atigun indicated that ectomycorrhizal fungi mobilized and assimilated considerable soil organic nitrogen that could then be transferred to animals as amino acids. Arctic ground squirrels are effectively functioning then as integrators of the organic nitrogen cycled through ectomycorrhizal fungi, and may prove useful in detecting if soil warming in the Arctic is increasing the mobilization of soil-derived nitrogen.

Data Accessibility

Data will be archived at the website of the Arctic Long-Term Ecological Research site at <http://ecosystems.mbl.edu/ARC/datacatalog.html>, and will be searchable by subject (title, abstract, or keywords), owner, spatial criteria, or taxonomic criteria.

The following appendices are available online for this article: Table A1 (isotopic values of plant food items at the Atigun site from samples collected in 2007 and 2008); Table A2 (stepwise multiple regressions of hair $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$); Table A3 (MixSIAR results for dietary estimates of consumption by arctic ground squirrels of dicots, graminoids, and fungi at Toolik and Atigun); Table A4 (stepwise multiple regression of fecal $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$); Table A5 (stepwise multiple regressions of fecal C/N); Table A6 (fungal taxa recorded and other observations in fecal samples from Atigun and Toolik); and Table A7 (fecal $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and C/N for samples that also had information on dietary components from microscopy).

ACKNOWLEDGMENTS

We thank Rebecca Rowe, Ryan Stephens, Andy Maguire, and several anonymous reviewers for comments on previous versions. This study was supported by the NSF LTER network and grants from U.S. National Science Foundation OPP-1108074 and IOS-1147232.

REFERENCES CITED

- Batzli, G. O., and Sobaski, S. T., 1980: Distribution, abundance, and foraging patterns of ground squirrels near Atkasook, Alaska. *Arctic and Alpine Research*, 12(3): 501–510.
- Beavan, N. R., and Sparks, R. J., 1998: Factors influencing ^{14}C ages of the Pacific rat *Rattus exulans*. *Radiocarbon*, 40(2): 601–613.
- Boonstra, R., Krebs, C. J., and Kanter, M., 1990: Arctic ground squirrel predation on collared lemmings. *Canadian Journal of Zoology*, 68(4): 757–760.
- Boonstra, R., Bradley, A. J., and Delehanty, B., 2011: Preparing for hibernation in ground squirrels: adrenal androgen production in summer linked to environmental severity in winter. *Functional Ecology*, 25(6): 1348–1359.
- Bradley, W. G., 1968: Food habits of the antelope ground squirrel in southern Nevada. *Journal of Mammalogy*, 49(1): 14–21.
- Brown, J., and Kreig, R., 1983: *Elliott and Dalton Highways, Fox to Prudhoe Bay, Alaska: Guidebook to Permafrost and Related Features*. Anchorage, Alaska: Alaska Division of Geological and Geophysical Surveys.
- Butterworth, B. B., 1958: Molt patterns in the barrow ground squirrel. *Journal of Mammalogy*, 39(1): 92–97.
- Carleton, S. A., Kelly, L., Anderson-Sprecher, R., and del Rio, C. M., 2008: Should we use one-, or multi-compartment models to describe ^{13}C incorporation into animal tissues? *Rapid Communications in Mass Spectrometry*, 22(19): 3008–3014.
- Castellano, M. A., Trappe, J. M., and Maser, C., 1989: *Key to Spores of the Genera of Hypogeous Fungi of North Temperate Forests with Special Reference to Animal Mycophagy*. Eureka, California: Mad River Press.
- Caut, S., Angulo, E., and Courchamp, F., 2008: Discrimination factors ($\Delta^{15}\text{N}$ and $\Delta^{13}\text{C}$) in an omnivorous consumer: effect of diet isotopic ratio. *Functional Ecology*, 22(2): 255–263.
- Chapin, F. S., Shaver, G. R., Giblin, A. E., Nadelhoffer, K. J., and Laundre, J. A., 1995: Responses of Arctic tundra to experimental and observed changes in climate. *Ecology*, 76(6): 694–711.
- Cherry, J. E., Déry, S. J., Cheng, Y., Stieglitz, M., Jacobs, A. S., and Pang, F., 2014: Climate and hydrometeorology of the Toolik Lake region and the Kuparuk River Basin: past, present, and future. In Hobbie, J. E., and Kling, G. W. (eds.), *Alaska's Changing Arctic: Ecological Consequences for Tundra, Streams, and Lakes*. New York: Oxford University Press, 21–60.
- Clemmensen, K. E., Sorensen, P. L., Michelsen, A., Jonasson, S., and Strom, L., 2008: Site-dependent N uptake from N-form mixtures by arctic plants, soil microbes and ectomycorrhizal fungi. *Oecologia*, 155(4): 771–783.
- Colgan, W., III, Carey, A. B., and Trappe, J. M., 1997: A reliable method of analyzing dietaries of mycophagous small mammals. *Northwestern Naturalist*, 78: 65–69.
- Flaherty, E. A., Ben-David, M., and Smith, W. P., 2010: Diet and food availability: implications for foraging and dispersal of Prince of Wales northern flying squirrels across managed landscapes. *Journal of Mammalogy*, 91(1): 79–91.
- Froehle, A. W., Kellner, C. M., and Schoeninger, M. J., 2010: FOCUS: effect of diet and protein source on carbon stable isotope ratios in collagen: follow up to Warinner and Tuross (2009). *Journal of Archaeological Science*, 37(10): 2662–2670.
- Gannes, L. Z., Obrien, D. M., and del Rio, C. M., 1997: Stable isotopes in animal ecology: assumptions, caveats, and a call for more laboratory experiments. *Ecology*, 78(4): 1271–1276.
- Hamilton, T. D., 1979: Radiocarbon dates and Quaternary stratigraphic sections, Philip Smith Mountains Quadrangle, Alaska. *U.S. Geological Survey Open-file Report*, 79-86643.
- Hasselquist, N. J., and Högborg, P., 2014: Dosage and duration effects of nitrogen additions on ectomycorrhizal sporocarp production and functioning: an example from two N-limited boreal forests. *Ecology and Evolution*, 4(15): 3015–3026.
- Hata, H., Shibata, J., Omori, K., Kohda, M., and Hori, M., 2015: Depth segregation and diet disparity revealed by stable isotope analyses in sympatric herbivorous cichlids in Lake Tanganyika. *Zoological Letters*, 1(1): 1.
- Hobbie, E. A., 2005: Using isotopic tracers to follow carbon and nitrogen cycling of fungi. In Dighton, J., Oudemans, P., and White, J. (eds.), *The Fungal Community: Its Organization and Role in the Ecosystem*. New York: Marcel Dekker, 361–381.
- Hobbie, E. A., 2017: Dietary protein content and tissue type control ^{13}C discrimination in mammals: an analytical approach. *Rapid Communications in Mass Spectrometry*, 31(7): 639–648.
- Hobbie, E., and Moore, J., 2017: Carbon and nitrogen isotopes and concentrations in terrestrial plants from a six-year (2006–2012) fertilization experiment at the Arctic LTER, Toolik Field Station, Alaska. Environmental Data Initiative. <http://dx.doi.org/10.6073/pasta/3ff5ffb62bb6e9c9ee2c523115ba6dbb>.
- Hobbie, S. E., Gough, L., and Shaver, G. R., 2005: Species compositional differences on different-aged glacial landscapes drive contrasting responses of tundra to nutrient addition. *Journal of Ecology*, 93(4): 770–782.
- Hobbie, J. E., Hobbie, E. A., Drossman, H., Conte, M., Weber, J., Shamhart, J., and Weinrobe, M., 2009: Mycorrhizal fungi supply nitrogen to host plants in Arctic tundra and boreal forests: ^{15}N is the key signal. *Canadian Journal of Microbiology*, 55(1): 84–94.
- Hobbie, E. A., Quimette, A. P., Schuur, E. A., Kierstead, D., Trappe, J. M., Bendiksen, K., and Ohenoja, E., 2013: Radiocarbon evidence for the mining of organic nitrogen from soil by mycorrhizal fungi. *Biogeochemistry*, 114(3): 381–389.
- Högborg, M. N., Högborg, P., and Myrold, D. D., 2007: Is microbial community composition in boreal forest soils determined by pH, C-to-N ratio, the trees, or all three? *Oecologia*, 150(4): 590–601.
- Holawaychuk, N., Petro, J., Finney, H., Farnham, R., and Gersper, P., 1966: Soils of Ogotoruk Creek watershed. In Wilimovsky, N. J., and Wolfe, J. N. (eds.), *Environment of the Cape Thompson Region, Alaska*. Oak Ridge, Tennessee: U.S. Atomic Energy Commission, 221–273.
- Hua, Q., Barbetti, M., and Rakowski, A. Z., 2013: Atmospheric radiocarbon for the period 1950–2010. *Radiocarbon*, 55(4): 2059–2072.

- Hubbs, A. H., and Boonstra, R., 1997: Population limitation in Arctic ground squirrels: effects of food and predation. *Journal of Animal Ecology*, 66(4): 527–541.
- Hyodo, F., 2015: Use of stable carbon and nitrogen isotopes in insect trophic ecology. *Entomological Science*, 18(3): 295–312.
- Ikeda, H., Kubota, K., Kagawa, A., and Sota, T., 2010: Diverse diet compositions among harpaline ground beetle species revealed by mixing model analyses of stable isotope ratios. *Ecological Entomology*, 35(3): 307–316.
- Johnson, A. W., and Packer, J. C., 1967: *Distribution, Ecology, and Cytology of the Ogotoruk Creek Flora and the History of Beringia*. Stanford, California: Stanford University Press.
- Karels, T. J., and Boonstra, R., 2000: Concurrent density dependence and independence in populations of arctic ground squirrels. *Nature*, 408(6811): 460–463.
- Kurle, C. M., Koch, P. L., Tershy, B. R., and Croll, D. A., 2014: The effects of sex, tissue type, and dietary components on stable isotope discrimination factors ($\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$) in mammalian omnivores. *Isotopes in Environmental and Health Studies*, 50(3): 307–321.
- Mayer, W. V., 1953: A preliminary study of the Barrow ground squirrel, *Citellus parryi barrowensis*. *Journal of Mammalogy*, 34(3): 334–345.
- McIlwee, A. P., and Johnson, C. N., 1998: The contribution of fungus to the diets of three mycophagous marsupials in *Eucalyptus* forests, revealed by stable isotope analysis. *Functional Ecology*, 12(2): 223–231.
- McKane, R. B., Johnson, L. C., Shaver, G. R., Nadelhoffer, K. J., Rastetter, E. B., Fry, B., Giblin, A. E., Kielland, K., Kwiatkowski, B. L., Laundre, J. A., and Murray, G., 2002: Resource-based niches provide a basis for plant species diversity and dominance in arctic tundra. *Nature*, 415(1): 68–71.
- McLean, I. G., 1985: Seasonal patterns and sexual differences in the feeding ecology of arctic ground squirrels (*Spermophilus parryi plesius*). *Canadian Journal of Zoology*, 63(6): 1298–1301.
- Miller, J. F., Millar, J. S., and Longstaffe, F. J., 2008: Carbon- and nitrogen-isotope tissue-diet discrimination and turnover rates in deer mice, *Peromyscus maniculatus*. *Canadian Journal of Zoology*, 86(7): 685–691.
- Myers-Smith, I. H., Forbes, B. C., Wilmking, M., Hallinger, M., Lantz, T., Blok, D., Tape, K. D., Macias-Fauria, M., Sass-Klaassen, U., and Lévesque, E., 2011: Shrub expansion in tundra ecosystems: dynamics, impacts and research priorities. *Environmental Research Letters*, 6(4): 045509.
- O'Donoghue, M., 1994: Early survival of juvenile snowshoe hares. *Ecology*, 75(6): 1582–1592.
- Pastor, J., Dewey, B., and Christian, D. P., 1996: Carbon and nutrient mineralization and fungal spore composition of fecal pellets from voles in Minnesota. *Ecography*, 19(1): 52–61.
- Rao, C., Sharma, G., and Shukla, A., 1997: Distribution of ectomycorrhizal fungi in pure stands of different age groups of *Pinus kesiya*. *Canadian Journal of Microbiology*, 43(1): 85–91.
- Salvarina, I., Yohannes, E., Siemers, B. M., and Koselj, K., 2013: Advantages of using fecal samples for stable isotope analysis in bats: evidence from a triple isotopic experiment. *Rapid Communications in Mass Spectrometry*, 27(17): 1945–1953.
- Semmens, B. X., Ward, E. J., Moore, J. W., and Darimont, C. T., 2009: Quantifying inter- and intra-population niche variability using hierarchical Bayesian stable isotope mixing models. *PLoS one*, 4(7): e6187.
- Shamhart, J., 2010: *Isotopic Analysis of Arctic Ground Squirrel Tissues and Potential Food Sources*. M.S. thesis, University of New Hampshire, Durham, New Hampshire.
- Shaver, G. R., and Chapin, F. S., 1991: Production: biomass relationships and element cycling in contrasting arctic vegetation types. *Ecological Monographs*, 61(1): 1–31.
- Sheriff, M. J., Kenagy, G. J., Richter, M., Lee, T., Tøien, Ø., Kohl, F., Buck, C. L., and Barnes, B. M., 2011: Phenological variation in annual timing of hibernation and breeding in nearby populations of Arctic ground squirrels. *Proceedings of the Royal Society of London B: Biological Sciences*, 278(1716): 2369–2375.
- Sheriff, M. J., Fridinger, R. W., Tøien, Ø., Barnes, B. M., and Buck, C. L., 2013: Metabolic rate and prehibernation fattening in free-living arctic ground squirrels. *Physiological and Biochemical Zoology*, 86(5): 515–527.
- Soudzilovskaia, N. A., Douma, J. C., Akhmetzhanova, A. A., Bodegom, P. M., Cornwell, W. K., Moens, E. J., Treseder, K. K., Tibbett, M., Wang, Y. P., and Cornelissen, J. H., 2015: Global patterns of plant root colonization intensity by mycorrhizal fungi explained by climate and soil chemistry. *Global Ecology and Biogeography*, 24(3): 371–382.
- Sponheimer, M., Robinson, T., Ayliffe, L., Passey, B., Roeder, B., Shipley, L., Lopez, E., Cerling, T., Dearing, D., and Ehleringer, J., 2003a: An experimental study of carbon-isotope fractionation between diet, hair, and feces of mammalian herbivores. *Canadian Journal of Zoology*, 81(5): 871–876.
- Sponheimer, M., Robinson, T., Ayliffe, L., Roeder, B., Hammer, J., Passey, B., West, A., Cerling, T., Dearing, D., and Ehleringer, J., 2003b: Nitrogen isotopes in mammalian herbivores: hair $\delta^{15}\text{N}$ values from a controlled feeding study. *International Journal of Osteoarchaeology*, 13(1): 80–87.
- Stock, B. C., and Semmens, B. X., 2013: MixSIAR GUI User Manual, Version 1.0, <http://conserver.iugo-cafe.org/user/brice.semmens/MixSIAR>, accessed on 27 January 2017.
- Tevis, L., 1953: Stomach contents of chipmunks and mantled squirrels in northeastern California. *Journal of Mammalogy*, 34(3): 316–324.
- Tibbett, M., Sanders, F. E., Minto, S. J., Dowell, M., and Cairney, J. W. G., 1998: Utilization of organic nitrogen by ectomycorrhizal fungi (*Hebeloma* spp.) of arctic and temperate origin. *Mycological Research*, 102(9): 1525–1532.
- Tieszen, L. L., Boutton, T. W., Tesdahl, K. G., and Slade, N. A., 1983: Fractionation and turnover of stable carbon isotopes in animal tissues—Implications for $\delta^{13}\text{C}$ analysis of diet. *Oecologia*, 57(1): 32–37.
- Walker, D., Short, S., Andrews, J., and Webber, P., 1981: Late Holocene pollen and present-day vegetation, Prudhoe Bay and Atigun River, Alaskan North Slope. *Arctic and Alpine Research*, 13(2): 153–172.
- Walker, D. A., and Maier, H. A., 2008: *Vegetation in the vicinity of the Toolik Field Station, Alaska*. Fairbanks: University of Alaska, Institute of Arctic Biology.

MS submitted 26 September 2016

MS accepted 12 June 2017