

Avoidance or Coexistence? The Spatiotemporal Patterns of Wild Mammals in a Human-dominated Landscape in the Western Himalaya

Authors: Srivastava, Nimisha, Krishnamurthy, Ramesh, and Sathyakumar, Sambandam

Source: Mountain Research and Development, 40(2)

Published By: International Mountain Society

URL: <https://doi.org/10.1659/MRD-JOURNAL-D-19-00046.1>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Avoidance or Coexistence? The Spatiotemporal Patterns of Wild Mammals in a Human-dominated Landscape in the Western Himalaya

Nimisha Srivastava^{1*}, Ramesh Krishnamurthy^{2,3}, and Sambandam Sathyakumar⁴

* Corresponding author: nimisha.mammals@gmail.com

¹ Wildlife Institute of India, Chandrabani, Post Box 18, Dehradun, Uttarakhand 248001, India

² Department of Landscape Level Planning and Management, Wildlife Institute of India, Chandrabani, Post Box 18, Dehradun, Uttarakhand 248001, India

³ Faculty of Forestry, University of British Columbia, Vancouver, BC, V6T 1Z4, Canada

⁴ Department of Endangered Species Management, Wildlife Institute of India, Chandrabani, Post Box 18, Dehradun, Uttarakhand 248001, India

© 2020 Srivastava et al. This open access article is licensed under a Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>). Please credit the authors and the full source.



Human-wildlife interfaces are increasing rapidly due to the disproportionate growth of human and wildlife populations in a spatial context. The Himalayan system, a global biodiversity hotspot, is subject to

landscape modification from various anthropogenic activities. In this study, we offer insights into the human-wildlife interface, reflecting avoidance or coexistence, with implications for local and landscape management strategies. We investigated fine-scale space use and temporal activity patterns of mammalian wildlife in a human-dominated landscape, outside a protected area. The research methods involved robust digital camera trap sampling ($n = 131$) across the target area (116 km^2) with a total human population of 153,585. We developed a new sampling strategy that accounted for spatial heterogeneity in the habitats and variations in mammalian community composition. Our results showed that, in spite of high usage and the presence of humans

across the study area, 16 wild mammal species used the area with varying intensities, exploiting habitat and forage availability. Of the camera traps placed in the study area, 70.23% had overlapping captures for humans on foot and wild mammal species. Generalist species used natural, modified, and altered habitats, while herbivores remained in natural and modified areas. However, some mammals that used modified/changed areas avoided humans by modifying their temporal activity. In the context of management of large landscapes, including areas outside the protected area network, the results of this study highlight the significant plasticity exhibited by wild mammals in negotiating natural and human-modified habitats. This offers an opportunity to develop conservation management strategies focusing on these fine-scale patterns and human actions.

Keywords: agricultural landscapes; animal behavior; Himalaya; human-wildlife interface; landscape management; activity pattern.

Peer-reviewed: December 2019 **Accepted:** May 2020

Introduction

Human population growth and related modification and loss of natural habitat continue to pose a serious threat to biodiversity (Noss et al 1996; Wilcove et al 1998; Western 2001; Tigas et al 2002). Due to alterations in land-use types and growing human populations around protected areas (PAs), these natural areas are becoming increasingly isolated patches (Wittemyer et al 2008). Hence, a landscape-level conservation approach that is ecologically representative and socially inclusive is needed. Many recent studies have focused on the role of agricultural landscapes, also termed anthropogenic landscapes, in the conservation of different taxa and have revealed their importance in conservation (Pimentel et al 1992; Harvey et al 2008). The biodiversity has been attributed to the habitat heterogeneity that such areas provide (Benton et al 2003; Bennett et al 2006; Gardner et al 2009; Anand et al 2010). This led the Convention on Biological Diversity (CBD) to issue Aichi Biodiversity Target 7, which focuses on the sustainable management of

agroecosystems to conserve biodiversity. The CBD also proposed the concept of “Other Effective Area-based Conservation Measures” to achieve Aichi Biodiversity Target 11, which aims to protect 17% of terrestrial area to promote landscape conservation, particularly in biodiversity-rich areas, such as the Himalaya (CBD Plan 2011–2020; CBD 2018).

Studies in anthropogenic landscapes have suggested that relatively more adaptable species can live in a matrix of natural and human-dominated habitats (Macdonald 1979; Fedriani et al 2001; Fuller et al 2010; Rodewald and Gehrt 2014) and show more tolerance of human-mediated disturbance (Doncaster and Macdonald 1991; Tigas et al 2002).

The current trend of conservation discourse highlights the importance of landscape management, and hence the need to assess the importance of areas outside PAs. About 90% of tropical forests lie outside the PA network (WWF 2002) and are threatened by continual pressures of habitat modification and deforestation (DeFries et al 2005). Most

studies conducted outside PAs are from North America, in and around the United Kingdom, and Australia (Tigas et al 2002; Beckmann and Berger 2003; Contesse et al 2004; Ditchkoff et al 2006; Newsome et al 2014); a few are from other regions of the world (Abay et al 2011; Athreya et al 2013). In India, however, little is known about the ecology of mammals found in proximity to human habitations. Most studies have focused on the negative interactions (Treves and Karanth 2003; Bhatia et al 2013; Naha et al 2018, 2019), with very few examples of the habitat-use patterns of wildlife in agricultural landscapes (Kumara et al 2004; Athreya et al 2013; Banerjee et al 2013; Ghoshal et al 2016). Some studies showing the importance of a matrix of forested and unforested areas have been conducted in different parts of India but are mostly restricted to birds (Elsen et al 2017), invertebrates (Dolia et al 2008; Anand et al 2010), and plants (Rawat et al 1999).

The Indian Himalayan region has a network of 138 PAs, covering 10.3% of the geographical region (WII NBWL, 2020), with a total forest cover of 33.7% (FSI 2017). Some of these are managed by local communities and are termed *van panchayat* (Agrawal 1999). Forests interspersed with human habitations are potential wildlife habitats outside PAs (Cuiti et al 2012). As a world biodiversity hotspot, holding 66% of Indian mammalian diversity (Chandra et al 2018), it is important to understand the significance of areas outside the PAs in the Indian Himalayan region.

Animals living in proximity to human-dominated landscapes are known to alter their behavior spatially and/or temporally (Graham et al 2009; Cuiti et al 2012; Valeix et al 2012; Carter et al 2015). In a region such as the Himalaya, we hypothesized that in order to use human-dominated landscape, wild animals would negotiate human presence in space and/or time to avoid direct encounters with humans. We carried out this study to understand the spatiotemporal patterns in wildlife populations using the human-dominated landscape of the Western Himalaya. Following Gardner et al (2009), we defined modified areas in our study as former native vegetation areas that have not been significantly altered to urban areas but to land uses such as agriculture, orchards, and settlements. We designed a sampling protocol to take into account all land-use types and the sampling biases that may generally occur in mountainous landscapes to investigate this idea.

Materials and methods

Study area

We carried out this study in the Mandal valley (area: 116 km²; 30°22'–30°30'N; 79°13'–79°21'E), a mid- to high-elevation (900–3400 meters above sea level [masl]) Himalayan area in the state of Uttarakhand, northern India (Figure 1). The Mandal valley touches the southern boundary of the Kedarnath Wildlife Sanctuary. This region provides access to Hindu shrines and tourist destinations in the region. A road connecting the famous Hindu shrines Kedarnath and Badrinath passes through this area, resulting in high vehicular traffic during May to October. Two Hindu shrines, Tungnath (3500 masl) in the northeast of the study area and Rudranath (3400 masl) in the northwest, attract significant numbers of pilgrims and tourists. Figure 2 shows the main land uses in the study area.

The study area is a densely populated, with a matrix of 11 urban and 35 small and large rural habitations (DCOU 2011), including the district headquarters, Gopeshwar/Chamoli.

This provides a gradient of human disturbance levels based on village/town type and human population density. Figure 3 shows the distribution of humans in the study area: They are not restricted to settlements and agriculture but instead are spread across the study area. The total human population within the intensive study area was 153,585 (DCOU 2011).

Kedarnath Wildlife Sanctuary (975 km²) has a wide diversity of mammalian fauna, most notably, mountain ungulates and large carnivores (Sathyakumar 1994). Local people are mostly dependent on forest resources and agriculture as their main sources of livelihood and include a few pastoral communities. Pressure from lopping and cutting varies in different *van panchayat* forests. However, the majority of the villagers, as observed during the study, value forests and are aware of their importance. They were also positive about conservation of wildlife and would agree to coexist if the harm caused by wild animals was stopped (personal observation, N. Srivastava).

Within the Mandal valley, we selected an intensive study area that ranged from 900 masl to 2500 masl with a diversity of land-use/land-cover types. Based on the objectives of this study, we deployed cameras up to 2500 masl; human settlements are not present above this.

Study design

The study area has high habitat heterogeneity, and, in order to cover the landscape configuration, we plotted grids of 1 km² in the study area. The heterogeneous landscape includes protected forests; *van panchayat* forests, which are dense to open and disturbed, due to lopping and grazing pressure; degraded/abandoned lands that have been taken over by scrub species, such *Lantana camara*, *Rubus* spp, *Berberis* spp, and others; agricultural areas; and human settlements. We placed camera traps with a minimum of 1 to maximum of 3 traps per grid based on habitat heterogeneity in the grid. Within a grid, the area was intensively searched for wild animal signs (droppings/tracks/scent marks) and for resources (crop fields, water). In crop fields, trails leading to forests and/or connecting two villages were chosen to optimize captures. Camera trap locations were selected after 45 days of reconnaissance. Sampling was done once per grid without repetitions, shifting cameras to other grids after each session (mean sampling days = 15) from lower elevations (January 2017 onwards) to higher elevations (April 2017 onwards) to maintain temperature effects. This approach of placing camera traps in human-dominated mountain areas has not been attempted before and, hence, can be considered a model to be tested for its efficacy in such landscapes.

Field methods

In total, 131 camera trap points were sampled with 40 Cuddeback C1 camera traps, covering areas from the most highly populated towns to the smallest villages (within the study area), as well as agricultural fields, protected and unprotected forests and orchards, garbage dump sites, planted forests, and fallow land. Given the mountainous terrain and fine-scale sampling, most locations were not accessible by road; accordingly, the study involved a

FIGURE 1 Map of the Mandal river subbasin (intensive study area). Kedarnath Wildlife Sanctuary (WLS) is situated in the north of the study area.




FIGURE 2 Land-use map of the study area. We used Sentinel 2A data acquired on 28 March 2017, tile no. T44RLU. The accuracy assessment was 75% for supervised and 66.2% for unsupervised classification (kappa statistics can be found in Appendixes S2 and S3, respectively; *Supplemental material*, <https://doi.org/10.1659/MRD-JOURNAL-D-19-00046.1.S1>). Shape files used for agriculture and settlements were manually digitized using Google Earth Pro and were overlaid on the classified shape file.





FIGURE 3 Heat map representing the distribution of humans in the study area based on a kernel density estimate.



substantial amount of walking (~1300 km). After a month of reconnaissance, camera traps were placed strategically and were set to capture photographs at 5 second intervals to maximize detection. At certain points, where the probability of false captures was high, the time interval was set to 10–15 seconds. Camera traps were placed 2–3 m above the ground to avoid vandalism. The cameras were adjusted in a way that covered the area of focus (eg animal trail) and were checked for detection sensitivity to ensure functioning. This was done by putting the camera on test mode and walking on the trail at different heights, as per study mammals heights. We did not use baits for our study. Photos captured after 5 minute intervals were selected for analysis in order to minimize chances of redundancy and reduce the possibility of biased photo capture rates for camera traps with a 10–15 second

interval between captures. We categorized six land-use types: settlements, crop fields, mosaic, fallow lands, dry forest, and moist forest. The criteria set for each habitat type are provided in Appendix S1 (*Supplemental material*, <https://doi.org/10.1659/MRD-JOURNAL-D-19-00046.1.S1>).

Analytical methods

A matrix of camera traps with habitat types was constructed for all species in MS Excel 2016, and mean and standard errors of photo capture rate per 100 trap nights were calculated for each wild species in different habitat types. In order to visualize the distribution of humans in the study area, we interpolated photo capture rates of humans and created a heat map. We used the software ArcMap 10.6.1 to

TABLE 1 List of mammal species photo-captured along with overall photo capture rate and number of photo captures recorded in Mandal river sub-basin (January 2017–May 2017).

Species name	Overall photo capture rate (no./100 trap nights) [Total no. photo captures]
Indian crested porcupine (<i>Hystrix indica</i>)	12.16 [236]
Sambar (<i>Rusa unicolor</i>)	9.17 [178]
Rhesus macaque (<i>Macaca mulatta</i>)	8.66 [168]
Barking deer (<i>Muntiacus muntjak</i>)	8.60 [167]
Wild pig (<i>Sus scrofa</i>)	7.83 [152]
Himalayan langur (<i>Semnopithecus schistaceus</i>)	6.75 [131]
Common leopard (<i>Panthera pardus</i>)	2.37 [48]
Leopard cat (<i>Prionailurus bengalensis</i>)	2.16 [42]
Himalayan palm civet (<i>Paguma larvata</i>)	2.16 [42]
Red fox (<i>Vulpes vulpes</i>)	1.55 [30]
Himalayan grey goral (<i>Naemorhedus goral bedfordi</i>)	1.55 [30]
Jungle cat (<i>Felis chaus</i>)	1.49 [29]
Himalayan serow (<i>Capricornis thar</i>)	1.13 [22]
Yellow-throated marten (<i>Martes flavigula</i>)	1.08 [21]
Golden jackal (<i>Canis aureus</i>)	0.15 [3]
Asiatic black bear (<i>Ursus thibetanus</i>)	0.05 [1]

create kernel density plots, using log-transformed capture rates. We overlaid the shape files of settlements and agriculture fields (digitized manually using Google Earth Pro imagery, where polygons were drawn around agricultural fields and settlements using the add polygon tool) over the map to understand the distribution of humans in the human-modified landscape.

Time stamping in each photo capture was used to estimate temporal activity patterns. For this analysis, time of photo captures was used and divided into two broad habitats, natural habitats and modified habitats, which included fallow land, mosaic habitats, crop fields, and settlements. Most of the modified areas were also close to human settlements, and it was expected that animals would show differences in activity patterns between the areas.

We used R studio, package camtrapR (Niedballa et al 2016), to summarize information on date and time and created a capture matrix for all species. Circular statistics that examined the temporal pattern of animal captures, reflecting time-specific behavioral responses of the species, were calculated and plotted using ORIANA software version 4 (Kovach 2011). For angular data, such as time, we conducted a nonparametric Watson U^2 test (Zar 1999), which was used to compare two sets of data using mean square deviations. We did this to see if our study species had significant difference in temporal activity patterns between natural and human-modified areas. We tested for temporal overlap between wild animals and humans by plotting the time of photo captures using MS Excel. This was done for the overall study area for diurnal and nocturnal species as found in our study. We went on to explore fine-scale temporal

overlap/avoidance in natural, modified, and settlement areas. For temporal analyses, we used the raw data of photo capture (ie no time interval was considered; the 5 minute interval was used for other analyses), as our objective was to look at overlap. We plotted kernel density plots using R 3.4.2 (R Core Team 2014). These plots helped to estimate the coefficient of overlap between the two data sets using the von Mises kernel for circular distributed data (Meredith and Ridout 2014).

Results

Spatial pattern

We recorded 16 species of wild mammals using the study area, despite the high ($n = 3733$) presence of humans across the land-cover/land-use types (Figure 3). The mean number of sampling sessions in the study area was 14.82 ± 0.67 trap nights. The number of photo captures of each species is listed in Table 1. Golden jackal (*Canis aureus*) and Asiatic black bear (*Ursus thibetanus*) had low photo captures ($n = 3$ and $n = 1$, respectively) and hence were excluded from further analyses. We found that 70.23% of the camera traps had overlapping captures for humans on foot and wild species. There were species-specific variations in the capture rates in different habitat types in the study area. The mean values of log-transformed photo captures per 100 trap nights are presented in Table 2. The two primates showed different habitat-use patterns: While rhesus macaques (*Macaca mulatta*) showed high usage in areas near human settlements (0.73 ± 0.88) and in fallow land (0.46 ± 0.93), Himalayan langurs (*Semnopithecus ajax*) were never captured in fallow land and

TABLE 2 Mean capture rates (number of captures/100 trap nights, log transformed) of wild mammals in different habitat types of Mandal river subbasin (December 2016–April 2017).

Species	Habitat type (mean \pm SD)					
	Settlement	Crop field	Mosaic	Fallow land	Dry forest	Moist forest
Himalayan langur	0.26 \pm 0.76	0.20 \pm 0.58	0.23 \pm 0.42	0.00	0.19 \pm 0.47	0.25 \pm 0.55
Rhesus macaque	0.73 \pm 0.88	0.28 \pm 0.60	0.16 \pm 0.40	0.46 \pm 0.93	0.06 \pm 0.28	0.06 \pm 0.26
Golden jackal	0.07 \pm 0.28	0.06 \pm 0.23	0.00	0.00	0.00	0.00
Himalayan masked palm civet	0.07 \pm 0.28	0.29 \pm 0.57	0.17 \pm 0.41	0.08 \pm 0.25	0.02 \pm 0.10	0.15 \pm 0.42
Jungle cat	0.13 \pm 0.37	0.36 \pm 0.69	0.11 \pm 0.32	0.17 \pm 0.35	0.00	0.02 \pm 0.12
Leopard cat	0.14 \pm 0.44	0.13 \pm 0.36	0.06 \pm 0.26	0.30 \pm 0.46	0.16 \pm 0.46	0.20 \pm 0.40
Red fox	0.00	0.15 \pm 0.41	0.11 \pm 0.37	0.00	0.14 \pm 0.33	0.11 \pm 0.37
Yellow-throated marten	0.10 \pm 0.31	0.05 \pm 0.22	0.09 \pm 0.26	0.07 \pm 0.20	0.04 \pm 0.19	0.15 \pm 0.39
Barking deer	0.00	0.33 \pm 0.53	0.57 \pm 0.78	0.36 \pm 0.72	0.23 \pm 0.49	0.52 \pm 0.65
Himalayan grey goral	0.00	0.09 \pm 0.35	0.00	0.00	0.34 \pm 0.47	0.18 \pm 0.41
Sambar	0.00	0.62 \pm 1.04	0.28 \pm 0.62	0.00	0.04 \pm 0.18	0.21 \pm 0.53
Himalayan serow	0.00	0.09 \pm 0.35	0.00	0.00	0.04 \pm 0.21	0.11 \pm 0.36
Wild pig	0.12 \pm 0.35	0.29 \pm 0.71	0.70 \pm 0.73	0.47 \pm 0.50	0.18 \pm 0.50	0.32 \pm 0.54
Indian crested porcupine	0.42 \pm 0.64	0.53 \pm 0.67	0.84 \pm 0.70	0.92 \pm 0.74	0.48 \pm 0.71	0.53 \pm 0.64
Asiatic black bear	0.00	0.00	0.00	0.00	0.05 \pm 0.26	0.00
Common leopard	0.09 \pm 0.27	0.06 \pm 0.23	0.30 \pm 0.48	0.39 \pm 0.63	0.11 \pm 0.38	0.21 \pm 0.47

had high usage of natural habitats, both dry (0.19 ± 0.47) and moist (0.25 ± 0.55). Small- to medium-sized carnivores were captured in almost all habitat types, except for red foxes (*Vulpes vulpes*), which were never captured in areas near human settlements and fallow land. We observed an interesting trend of ungulates avoiding areas near human settlements, except wild pig (0.12 ± 0.35). Goral (*Naemorhedus goral bedfordi*) and serow (*Capricornis thar*) were never captured in other disturbed areas, such as mosaic and fallow land, except for one instance for each, in crop fields. Indian crested porcupine, a rodent, was photo captured in all habitat types with similar frequency and was also the most frequently photo-captured species among all wild mammals ($n = 236$). Leopards used all habitat types, though they were photo-captured relatively more in mosaic (0.30 ± 0.48), fallow land (0.39 ± 0.63), and moist forest (0.21 ± 0.47).

Temporal pattern

Rose diagrams of temporal activity for study species are presented in Figure 4. In cases where the line indicating standard deviation is red, deviation from the expected directionality was wider than expected. Results of the Watson U^2 test showed that species such as Himalayan masked palm civet, red fox, yellow-throated marten, and barking deer showed no significant difference in temporal activity pattern between the two habitats. Other species, such as Himalayan langur (0.74 , $P < 0.001$), rhesus macaque (0.33 , $P < 0.005$), wild pig (0.58 , $P < 0.001$), and Indian crested porcupine (0.38 , $P < 0.005$), showed a significant difference in their temporal activity between the two habitats. While there was no significant difference in usage

between the two habitats by common leopard (*Panthera pardus*), leopard cat (*Prionailurus bengalensis*), and sambar (*Rusa unicorn*), it is interesting to note that these species had an altered peak activity period after midnight in modified areas (Figure 4). Jungle cat, serow, and goral were captured only once in either habitat, and so a difference in temporal pattern could not be determined. However, others that did not show a significant difference in temporal activity between natural and modified areas showed different activity peaks in the two habitat types (Figure 4). The average period of daylight during the study was 7:00 h to 18:00 h.

The temporal-overlap graphs for human and wild species (diurnal and nocturnal) (Figure 5) show that the activity of diurnal species, such as langur, macaque, barking deer, goral, and marten, overlapped with human activity; however, barking deer was also found to be active in nocturnal hours. The human–animal temporal overlap (Table 3) highlights the observation that nocturnal animals such as wild pig (Dhat value = 0.04), sambar (0.04), common leopard (0.04), and jungle cat (0.04) had minimal temporal overlap with humans in settlements, though this was not significantly different from other land-use types. Hence, these animals have adapted to avoid direct encounters with humans, but they still access these areas for resources. Except for langur (Dhat value = 0.44, 0.39 modified and settlement habitats, respectively), macaque (0.57, 0.61), barking deer (0.62 in modified habitat), and yellow-throated marten (0.45 in modified habitat), all other species had a significantly low overlap percentage in modified and/or settlement areas.

FIGURE 4 Temporal rose diagrams of wild animals in natural (f) and modified (nf) areas.




TABLE 3 Temporal overlap of wild mammals with humans in three habitat types of Mandal river subbasin (December 2016–April 2017).

Species	Habitat	Sample size	Dhat value
Himalayan langur	Natural forest	55	0.73
	Modified	29	0.44
	Settlement	47	0.39
Rhesus macaque	Natural forest	55	0.51
	Modified	76	0.57
	Settlement	81	0.61
Barking deer	Natural forest	82	0.42
	Modified	85	0.62
	Settlement	0	NA
Himalayan grey goral	Natural forest	28	0.66
	Modified	2	NA
	Settlement	0	NA
Sambar	Natural forest	29	0.09
	Modified	149	0.04
	Settlement	0	NA
Himalayan serow	Natural forest	20	0.40
	Modified	2	NA
	Settlement	0	NA
Wild pig	Natural forest	42	0.16
	Modified	106	0.12
	Settlement	4	0.02
Common leopard	Natural forest	22	0.25
	Modified	22	0.39
	Settlement	2	0.04
Jungle cat	Natural forest	1	NA
	Modified	24	0.14
	Settlement	4	0.04
Leopard cat	Natural forest	25	0.08
	Modified	10	0.07
	Settlement	07	0.20
Himalayan palm civet	Natural forest	22	0.43
	Modified	18	0.13
	Settlement	2	0.13
Red fox	Natural forest	20	0.11
	Modified	10	0.10
	Settlement	0	NA

TABLE 3 Continued.

Species	Habitat	Sample size	Dhat value
Yellow-throated marten	Natural forest	15	0.63
	Modified	4	0.45
	Settlement	2	0.51
Indian crested porcupine	Natural forest	98	0.05
	Modified	122	0.12
	Settlement	16	0.07


NA, not applicable.

Discussion

Our results show that the majority of species used modified areas, but with some adjustments in usage in space and time. Herbivores such as barking deer, goral, serow, and sambar avoided areas near settlements, as evident from zero to very low captures in such areas. Species such as goral and serow did not use areas that had been modified, such as agricultural lands. This was in accordance with other studies that found these animals were sensitive to human-mediated disturbances, such as trails used frequently by humans (eg tourist trails; Bhattacharya et al 2012), and to agricultural/modified areas and settlements (Paudel and Kindlmann 2012). Species such as barking deer and wild pig have been reported to avoid areas near settlements (Azlan 2006; Paudel and Kindlmann 2012).

Small mammals used almost all habitat types; however, species such as jungle cat, red fox, and civet showed high

FIGURE 5 Temporal patterns of humans and wildlife based on camera trap photo captures in the study area: (A) diurnal species; (B) nocturnal species.



capture rates in crop fields. This could be because of their diet and/or the type of habitat they prefer (Mudappa 2013). For example, rodents are major pests in crop fields (Stenseth et al 2003), and they are prey for many of the small carnivores, which are consequently attracted to these habitats. Indian crested porcupine was the most common wild animal using all the habitats in study area; being nocturnal, it could directly avoid humans. Studies done elsewhere found similar results for small carnivores, which in general are omnivorous (Fedriani et al 2001; Fuller et al 2010; Rodewald and Gehrt 2014). They are also reported to be tolerant of human-mediated disturbance (Doncaster and Macdonald 1991; Tigas et al 2002). Small and large carnivore species such as red fox (Gloor 2002; Contesse et al 2004; Ditchkoff et al 2006; Baker 2007; Díaz-Ruiz et al 2016; Ghoshal et al 2016), coyotes (*Canis latrans*) (McClure et al 1995; Fedriani et al 2001), spotted hyena (*Crocuta crocuta*) (Abay et al 2011), golden jackal (Macdonald 1979; Yom-Tov et al 1995), and Australian dingo (*Canis lupus dingo*) (Newsome et al 2014) have all been found to be dependent on anthropogenic subsidies in semirural to urban landscapes, shifting their activity to become primarily nocturnal in response to humans.

The large carnivore, common leopard, used fallow and mosaic land more than other natural and nonnatural habitats. The use of shrub-dominated areas by leopards was also found by Aggrawal et al (2011) and Bhattacharjee (2006). This could be due to the cover that this vegetation offers, which is particularly important where chances of human encounters are higher, as is the case in the modified areas. Golden jackal, which is fairly common in these areas in the early 1990s (S Sathyakumar, personal observation), is clearly in decline. This was also reported by people in the study area and is evident from the low number of photo captures. This may be due to an increase in the population of dogs in the human-dominated landscapes. The decline in populations of jackal has been observed in other areas and is a concern for this “once common” species (Pillay et al 2011; Srivastava 2019).

With humans present throughout the study area, it is almost impossible for wildlife to avoid humans spatially, and so we looked at how they negotiated humans temporally. Most of the diurnal species did not show much difference in temporal activity pattern, as was expected. According to Gaynor et al (2018), in order to use anthropogenic areas, wild animals shift to a nocturnal activity. In our study, we found that only a few species reported to have diurnal activity pattern in natural areas or PAs, such as wild pig (Azlan 2006; Grassman et al 2006) and common leopard (Ngoprasert et al 2007; Odden et al 2014; Carter et al 2015; Van Cleave et al 2018), were found to be predominantly nocturnal throughout the study area, including inside forests. However, other diurnal species, such as barking deer, were found to be active also in the night hours; this has not been reported in other studies examining human presence in a landscape, such as those by Griffiths and Schaik (1993), Carter et al (2015), and Ota et al (2019).

The activity of wild pig in modified areas was clumped within a short range of time. Similar patterns were mentioned in Prater (1965). A narrow range of temporal activity was also observed in leopard cat, where the peak activity period in modified areas was from 00:00 h to 03:00 h. Yellow-throated martens were active during daytime, while

they followed a more crepuscular pattern in the forest. To the best of our knowledge, this study is the first to describe the time activity for this species. Large carnivores such as tigers (*Panthera tigris*) (Carter et al 2012) and leopards (Ngoprasert et al 2007; Odden et al 2014) have been found to shift their diurnal activity pattern to nocturnal while using human-dominated areas (Azlan and Sharma 2006), as found in our study. Leopards in our study area were predominately nocturnal.

In a previous study, moose (*Alces alces*), a large-bodied ungulate, was found to alter its temporal activity in modified areas, using the area only at night (Bjørneraas et al 2011). A similar large herbivore in our study, the sambar, exhibited nocturnal activity; however, this animal generally follows nocturnal or crepuscular activity in undisturbed areas (Matsubayashi et al 2007). Diurnal species such as barking deer, goral, and marten showed no significant difference in temporal activity pattern between natural and modified areas and in relation to humans. This suggests adaptation to human presence, adopting a strategy of flight if humans are encountered, rather than shifting temporal activity. Sightings of barking deer, followed by goral, were the most common sightings during the present study period. In all these encounters, barking deer ran off into the woods within seconds. This suggests that these animals are habituated to human presence, as also reported by Griffiths and Schaik (1993). We also found that, although a few species had different activity patterns in natural or modified areas, others had similar temporal overlaps with humans in natural and other habitat types, indicating behavioral adaptations allowing animals to survive in human-dominated landscapes. Hence, it seems that adjustment of spatial and temporal patterns of habitat use helps wild animals to utilize modified areas for different purposes. Availability of subsidized food resources (crops for herbivores [Bayani et al 2016; Thinley et al 2017] and rodents/livestock for carnivores [Sekhar 1998; Madhusudan 2003; Rajaratnam et al 2007]), allowing optimal foraging, and cover (use of habitats such as mosaic and secondary scrubland in the study area) seem to govern the use of human-modified area by wild animals. However, long-term studies are required to support our findings and establish the ultimate causes of such patterns of habitat use.

The results of this study highlight that the wild mammals exhibit significant plasticity, allowing them to thrive in human-dominated landscapes. However, co-occurrence of wildlife and humans is often negative (Yom-Tov et al 1995; Baker 2007; Oro et al 2013), involving, for example, crop raiding (Madhusudan 2003; Charoo et al 2011), livestock depredation (Naha et al 2018), attacks on humans (Dhanwatey et al 2013; Naha et al 2019), and retaliatory killings by humans (Mishra et al 2003; Kissui 2008; Pandey 2019). In our study, despite having high overlap with humans, coexistence seemed to be largely positive (personal observation). This is likely to continue as long as the status quo is maintained, but the factors that could cause a tipping point/threshold and the time table over which this might occur can only be understood by long-term observation.

This approach of investigating wild-animal use of human-modified systems outside of PAs has never been carried out in the Indian Himalayan region, making this study important from a conservation point of view. In addition, with the current trajectory of development goals in the Indian Himalaya, such as hydropower plants (Grumbine and Pandit

2013) and urbanization (Pandit et al 2014), this is a critical time to study how mountain ecosystems can be developed sustainably. Our study demonstrates a new design for estimating habitat use by mammals in a heterogeneous area of human-modified landscapes in mountain systems. Since mountain ecosystems are similar all over the world (Smith 2014), with terrace farming and villages interspersed with forests (Heimbuch 2020), our study has wider implications.

Our findings are preliminary and showcase how animals adapt to live in human-dominated areas, but they also show that certain species, especially herbivores, can be negatively affected by modifications of natural habitats. Although we suggest that development activities should be carried out in a sustainable way, further studies investigating the extent to which different species are affected by various land-use types are required to provide better insights.

ACKNOWLEDGMENTS

We thank the Uttarakhand State Forest Department and the divisional forest officer of the Kedarnath Wildlife Division for granting us permission for the study. We are grateful to the village heads and people of Mandal valley. We especially thank our field assistants for their support and cooperation during fieldwork. We thank the Wildlife Institute of India for providing logistic support for the study and the funding provided from the fees paid by my parents (Srivastava).

REFERENCES

- Abay GY, Bauer H, Gebrihiwot K, Deckers J. 2011. Peri-urban spotted hyena (*Crocuta crocuta*) in northern Ethiopia: Diet, economic impact, and abundance. *European Journal Wildlife Research* 57:759–765.
- Aggrawal M, Chauhan DS, Goyal SP, Qureshi Q. 2011. Managing human-leopard conflicts in Pauri Garhwal, Uttarakhand, India, using geographical information system and remote sensing. *International Journal of Scientific & Engineering Research* 2(9):1–7.
- Agarwal R. 1999. Van panchayats in Uttarakhand: A case study. *Economic and Political Weekly* 34(39):2779–2781. <https://www.jstor.org/stable/4408446>; accessed on 23 July 2020.
- Anand MO, Krishnaswamy J, Kumar A, Bali A. 2010. Sustaining biodiversity conservation in human-modified landscapes in the Western Ghats: Remnant forests matter. *Biological Conservation* 143:2363–2374.
- Athreya V, Odden M, Linnell JD, Krishnaswamy J, Karanth U. 2013. Big cats in our backyards: Persistence of large carnivores in a human dominated landscape in India. *PLoS One* 8(3):e57872. <https://doi.org/10.1371/journal.pone.0057872>.
- Azlan JM. 2006. Mammal diversity and conservation in a secondary forest in Peninsular Malaysia. *Biodiversity & Conservation* 15:1013–1025.
- Azlan JM, Sharma DS. 2006. The diversity and activity patterns of wild felids in a secondary forest in peninsular Malaysia. *Oryx* 40(1):36–41.
- Baker RO. 2007. A review of successful urban coyote management programs implemented to prevent or reduce attacks on humans and pets in southern California. In: Nolte DL, Arjo WM, Stalman DH, editors. *Proceedings of the 12th Wildlife Damage Management Conference*. Internet Center for Wildlife Damage Management, abstract 58. https://digitalcommons.unl.edu/icwdm_wdmconfproc/58/; accessed on 23 July 2020.
- Banerjee K, Jhala YV, Chauhan KS, Dave CV. 2013. Living with lions: The economics of coexistence in the Gir forests, India. *PLoS One* 8:e49457.
- Bayani A, Tiwade D, Dongre A, Dongre AP, Phatak R, Watve M. 2016. Assessment of crop damage by protected wild mammalian herbivores on the western boundary of Tadoba-Andhari Tiger Reserve (TATR), central India. *PLoS One* 11(4):e0153854. <https://doi.org/10.1371/journal.pone.0153854>.
- Beckmann JP, Berger J. 2003. Using black bears to test ideal-free distribution models experimentally. *Journal of Mammalogy* 84:594–606.
- Bennett AF, Radford JQ, Harlem A. 2006. Properties of land mosaics: Implications for nature conservation in agricultural environments. *Biological Conservation* 133:250–264.
- Benton TG, Vickery JA, Wilson JD. 2003. Farmland biodiversity: Is habitat heterogeneity the key? *Trends in Ecology and Evolution* 18:182–188.
- Bhatia S, Athreya V, Grenyer R, Macdonald DW. 2013. Understanding the role of representations of human-leopard conflict in Mumbai through media-content analysis. *Conservation Biology* 27:588–594.
- Bhattacharjee S. 2006. *Monitoring Leopard (Panthera pardus) and its Prey in Different Human-Leopard Conflict Zones (High, Medium and Low), Pauri Garhwal, Uttarakhand* [PhD dissertation]. Dehradun, India: Forest Research Institute.
- Bhattacharya T, Bashir T, Poudyal K, Sathyakumar S, Saha GK. 2012. Distribution, occupancy and activity patterns of goral (*Nemorhaedus goral*) and serow (*Capricornis thar*) in Khangchendzonga Biosphere Reserve, Sikkim, India. *Mammal Study* 37:173–181.
- Bjørneraas K, Solberg EJ, Herfindal I, Moorter BV, Rolandsen CM, Tremblay JP, Skarpe C, Sæther BE, Eriksen R, Astrup R. 2011. Moose *Alces alces* habitat use at multiple temporal scales in a human-altered landscape. *Wildlife Biology* 17(1):44–54.
- Carter N, Jasny M, Gurung B, Liu J. 2015. Impacts of people and tigers on leopard spatiotemporal activity patterns in a global biodiversity hotspot. *Global Ecology and Conservation* 3:149–162.
- Carter NH, Shrestha BK, Karki JB, Pradhan NMB, Liu J. 2012. Coexistence between wildlife and humans at fine spatial scales. *Proceedings of the National Academy of Sciences of the United States of America* 109(38):15360–15365.
- CBD [Convention on Biological Diversity]. 2018 *Aichi Biodiversity Targets*. Strategic Plan 2011–2020//Aichi Targets. Montreal, Canada: CBD. <https://www.cbd.int/sp/targets>; accessed on 9 June 2019.
- Chandra K, Gupta D, Gopi KC, Tripathy B, Kumar V. 2018. *Faunal Diversity of Indian Himalaya*. Kolkata, India: Zoological Survey of India.
- Charoo SA, Sharma LK, Sathyakumar S. 2011. Asiatic black bear-human interactions around Dachigam National Park, Kashmir, India. *Ursus* 22(2):106–113.
- Contesse P, Hegglin D, Gloor S, Bontadina F, Deplazes P. 2004. The diet of urban foxes (*Vulpes vulpes*) and the availability of anthropogenic food in the city of Zurich, Switzerland. *Mammalian Biology* 69:81–95.
- Cuiti S, Northrup JM, Muhly TB, Simi S, Musiani M, Pitt JA, Boyce MS. 2012. Effects of humans on behaviour of wildlife exceed those of natural predators in a landscape of fear. *PLoS One* 7:50611.
- DCOU [Directorate of Census Operations, Uttarakhand]. 2011. *District Census Handbook, Chamoli: Village and Town Wise Primary Census Abstract (PCA)*. Dehradun, India: Directorate of Census Operations, Uttarakhand. https://censusindia.gov.in/2011census/dchb/0502_PART_B_DCHB_CHAMOLI.pdf; accessed on 6 October 2020.
- DeFries R, Hansen A, Newton AC, Hansen MC. 2005. Increasing isolation of protected areas in tropical forests over the past twenty years. *Ecological Applications* 15:19–26.
- Dhanwatey HS, Crawford JC, Abade LA, Dhanwatey PH, Nielsen CK, Sillero-Zubiri C. 2013. Large carnivore attacks on humans in central India: A case study from the Tadoba-Andhari Tiger Reserve. *Oryx* 47(2):221–227.
- Díaz-Ruiz F, Caro J, Delibes-Mateos M, Arroyo B, Ferreras P. 2016. Drivers of red fox (*Vulpes vulpes*) daily activity: Prey availability, human disturbance or habitat structure? *Journal of Zoology* 298(2):128–138.
- Ditchkoff SS, Saalfeld ST, Gibson CJ. 2006. Animal behavior in urban ecosystems: Modifications due to human-induced stress. *Urban Ecosystems* 9:5–12.
- Dolia J, Devy MS, Aravind NA, Kumar A. 2008. Adult butterfly communities in coffee plantations around protected area in the Western Ghats, India. *Animal Conservation* 11(1):26–34.
- Doncaster CP, Macdonald DW. 1991. Drifting territoriality in the red fox *Vulpes vulpes*. *The Journal of Animal Ecology* 60(2):423–439. <https://doi.org/10.2307/5288>.
- Elsen PR, Kalyanaraman R, Ramesh K, Wilcove DS. 2017. The importance of agricultural lands for Himalayan birds in winter. *Conservation Biology* 31:416–426.
- Fedriani JM, Fuller TK, Sauvajot RM. 2001. Does availability of anthropogenic food enhance densities of omnivorous mammals? An example with coyotes in southern California. *Ecography* 24:325–331.
- FSI [Forest Survey of India]. 2017. *State of Forest Report*. Dehradun, India: Forest Survey of India, Ministry of Environment, Forests and Climate Change, India. <http://fsi.nic.in/forest-report-2017>; accessed on 12 July 2019.
- Fuller TK, DeStefano S, Warren PS. 2010. Carnivore behavior and ecology, and relationship to urbanization. In: Gehrt SD, Riley SPD, Cypher BL, editors. *Urban Carnivores: Ecology, Conflict, and Conservation*. Baltimore, MD: Johns Hopkins University, pp 13–9.
- Gardner TA, Barlow J, Chazdon R, Ewers RM, Harvey CA, Peres CA, Sodhi N. 2009. Prospects for tropical forest biodiversity in a human-modified world. *Ecology Letters* 12:561–582.
- Gaynor KM, Hohnowski CE, Carter NH, Brashares JS. 2018. The influence of human disturbance on wildlife nocturnality. *Science* 360(6394):1232–1235. <https://doi.org/10.1126/science.aar7121>.
- Ghoshal A, Bhatnagar YV, Mishra C, Suryawanshi K. 2016. Response of the red fox to expansion of human habitation in the Trans-Himalayan mountains. *European Journal of Wildlife Research* 62:131–136.
- Gloor S. 2002. *The Rise of Urban Foxes (Vulpes vulpes) in Switzerland and Ecological and Parasitological Aspects of a Fox Population in the Recently Colonised City of Zurich* [PhD dissertation]. Zurich, Switzerland: University of Zurich.
- Graham MD, Douglas-Hamilton I, Adams WM, Lee PC. 2009. The movement of African elephants in a human-dominated land-use mosaic. *Animal Conservation* 12:445–455.
- Grassman LI, Haines AM, Janečka JE, Tewes ME. 2006. Activity periods of photo-captured mammals in north central Thailand/Périodes d'activité des mammifères photo-capturés en Thaïlande. *Mammalia* 70(3/4):306–309.
- Griffiths M, Schaik CP. 1993. The impact of human traffic on the abundance and activity periods of Sumatran rain forest wildlife. *Conservation Biology* 7(3):623–626.
- Grumbine RE, Pandit MK. 2013. Threats from India's Himalaya dams. *Science* 339(6115):36–37.
- Harvey CA, Komar O, Chazdon R, Ferguson BG, Finegan B, Griffith DM, Martínez-Ramos M, Morales H, Nigh R, Soto-Pinto L, et al. 2008. Integrating agricultural landscapes with biodiversity conservation in the Mesoamerican hotspot. *Conservation Biology* 22:8–15.

- Heimbuch J.** 2020. Examples of terrace farming around the world. *Treehugger*. <https://www.treehugger.com/examples-terrace-farming-around-world-4869737>; accessed on 24 July 2020.
- Kissui BM.** 2008. Livestock predation by lions, leopards, spotted hyenas, and their vulnerability to retaliatory killing in the Maasai steppe, Tanzania. *Animal Conservation* 11(5):422–432.
- Kovach WL.** 2011. *Oriana—Circular Statistics for Windows*, Ver. 4. Pentraeth, United Kingdom: Kovach Computing Services.
- Kumara HN, Kumar M, Sharma AK, Sushma HS, Singh M, Singh M.** 2004. Diversity and management of wild mammals in tea gardens in the rainforest regions of the Western Ghats, India: A case study from a tea estate in the Anaimalai Hills. *Current Science* 87:1282–1287.
- Macdonald DW.** 1979. The flexible social system of the golden jackal, *Canis aureus*. *Behavioral Ecology and Sociobiology* 5:17–38.
- Madhusudan MD.** 2003. Living amidst large wildlife: Livestock and crop depredation by large mammals in the interior villages of Bhadra Tiger Reserve, South India. *Environmental Management* 31(4):466–475.
- Matsubayashi H, Lagan P, Sukor JRA, Kitayama K.** 2007. Seasonal and daily use of natural licks by sambar deer (*Cervus unicolor*) in a Bornean tropical rain forest. *Tropics* 17(1):81–86.
- McClure MF, Smith NS, Shaw WW.** 1995. Diets of coyotes near the boundary of Saguaro National Monument and Tucson, Arizona. *The Southwest Naturalist* 40:101–104.
- Meredith M, Ridout M.** 2014. Overview of the Overlap Package. <https://cran.uib.no/web/packages/overlap/vignettes/overlap.pdf>; accessed on 23 July 2020.
- Mishra C, Allen P, McCarthy TOM, Madhusudan MD, Bayarjargal A, Prins HH.** 2003. The role of incentive programs in conserving the snow leopard. *Conservation Biology* 17(6):1512–1520.
- Mudappa D.** 2013. Herpestids, viverrids and mustelids. In: Johnsingh AJT, Manjrekar N, editors. *Mammals of South Asia*. Vol. 1. Hyderabad, India: University Press, pp 471–498.
- Naha D, Sathyakumar S, Dash S, Chettri A, Rawat GS.** 2019. Assessment and prediction of spatial patterns of human–elephant conflicts in changing land cover scenarios of a human-dominated landscape in North Bengal. *PLoS One* 14(2):e0210580. <https://doi.org/10.1371/journal.pone.0210580>.
- Naha D, Sathyakumar S, Rawat GS.** 2018. Understanding drivers of human–leopard conflicts in the Indian Himalayan region: Spatio-temporal patterns of conflicts and perception of local communities towards conserving large carnivores. *PLoS One* 13(10):e0204528.
- Newsome TM, Ballard GA, Fleming PJ, van de Ven R, Story GL, Dickman DR.** 2014. Human-resource subsidies alter the dietary preferences of a mammalian top predator. *Oecologia* 175:139–150.
- Ngoprasert D, Lynam AJ, Gale GA.** 2007. Human disturbance affects habitat use and behaviour of Asiatic leopard *Panthera pardus* in Kaeng Krachan National Park, Thailand. *Oryx* 41(3):343–351.
- Niedballa J, Sollmann R, Courtiol A, Wilting A.** 2016. camtrapR: An R package for efficient camera trap data management. *Methods in Ecology and Evolution* 7(12):1457–1462.
- Noss RF, Quigley HB, Hornocker MG, Merrill T, Paquet PC.** 1996. Conservation biology and carnivore conservation in the Rocky Mountains. *Conservation Biology* 10:949–963.
- Odden M, Athreya V, Rattan S, Linnell JD.** 2014. Adaptable neighbours: Movement patterns of GPS-collared leopards in human dominated landscapes in India. *PLoS One* 9(11):e112044. <https://doi.org/10.1371/journal.pone.0112044>.
- Oro D, Genovart M, Tavecchia G, Fowler MS, Martínez-Abraín A.** 2013. Ecological and evolutionary implications of food subsidies from humans. *Ecology Letters* 16:1501–1514.
- Ota A, Takagi E, Yasuda M, Hashim M, Hosaka T, Numata S.** 2019. Effects of nonlethal tourist activity on the diel activity patterns of mammals in a national park in peninsular Malaysia. *Global Ecology and Conservation* 20:e00772.
- Pandey A.** Tigris beaten to death in UP, villagers make video with commentary. NDTV [New Delhi], 26 July 2019. <https://www.ndtv.com/india-news/uttar-pradesh-pilibhit-tigris-beaten-to-death-villagers-recorded-video-2075578>; accessed on 5 December 2019.
- Pandit MK, Manish K, Koh LP.** 2014. Dancing on the roof of the world: Ecological transformation of the Himalayan landscape. *BioScience* 64(11):980–992.
- Paudel PK, Kindlmann P.** 2012. Human disturbance is a major determinant of wildlife distribution in Himalayan midhill landscapes of Nepal. *Animal Conservation* 15(3):283–293.
- Pillay R, Johnsingh AJT, Raghunath R, Madhusudan MD.** 2011. Patterns of spatiotemporal change in large mammal distribution and abundance in the southern Western Ghats, India. *Biological Conservation* 144:1567–1576.
- Pimentel D, Stachow U, Takacs DA, Brubaker HW, Dumas AR, Meaney JJ, Onsi DE, Corzilius DB.** 1992. Conserving biological diversity in agricultural/forestry systems. *BioScience* 42:354–362.
- Prater SH.** 1965. Pig. In: Prater, SH, editor. *The Book of Indian Animals*. Vol. 2. Bombay, India: Bombay Natural History Society, pp 298–300.
- R Core Team.** 2014. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. <http://www.R-project.org/>; accessed on 26 September 2019.
- Rajaratnam R, Sunquist M, Rajaratnam L, Ambu L.** 2007. Diet and habitat selection of the leopard cat (*Prionailurus bengalensis borneoensis*) in an agricultural landscape in Sabah, Malaysian Borneo. *Journal of Tropical Ecology* 23(2):209–217.
- Rawat GS, Sathyakumar S, Prasad SN.** 1999. Plant species diversity and community structure in the outer fringes of Kedarnath Wildlife Sanctuary, western Himalaya: Conservation implications. *Indian Forester* 125:873–882.
- Rodewald AD, Gehrt SD.** 2014. Wildlife population dynamics in urban landscapes. In: McCleery RA, Moorman C, Peterson MN, editors. *Urban Wildlife Conservation*. Berlin, Germany: Springer, pp 117–147.
- Sathyakumar S.** 1994. *Habitat Ecology of Major Ungulates in Kedarnath Musk Deer Sanctuary, Western Himalaya* [PhD Dissertation]. Rajkot, India: Saurashtra University.
- Sekhar N.** 1998. Crop and livestock depredation caused by wild animals in protected areas: The case of Sariska Tiger Reserve, Rajasthan, India. *Environment Conservation* 25(2):160–171.
- Smith JMB.** 2014. *Mountain Ecosystems*. Chicago, IL: Encyclopedia Britannica. <https://www.britannica.com/science/mountain-ecosystem>; accessed on 30 April 2020.
- Srivastava N.** 2019. Are the “rice-eating jackals” of Rann of Kachchh gone? *Journal of Bombay Natural History Society* 116:91. <https://doi.org/10.17087/jbnhs/2019/v116/145219>.
- Stenseth NC, Leirs H, Skonhoff A, Davis SA, Pech RP, Andreassen HP, Singleton GR, Lima M, Machang'u RS, Makundi RH, Zhang Z.** 2003. Mice, rats, and people: The bio-economics of agricultural rodent pests. *Frontiers in Ecology and the Environment* 1(7):367–375.
- Thinley P, Lassoie JP, Morreale SJ, Curtis PD, Rajaratnam R, Vernes K, Lek L, Phuntsho S, Dorji T, Dorji P.** 2017. High relative abundance of wild ungulates near agricultural croplands in a livestock-dominated landscape in western Bhutan: Implications for crop damage and protection. *Agriculture, Ecosystems & Environment* 248:88–95.
- Tigas LA, Van Vuren DH, Sauvajot RM.** 2002. Behavioral responses of bobcats and coyotes to habitat fragmentation and corridors in an urban environment. *Biological Conservation* 108:299–306.
- Treves A, Karanth KU.** 2003. Human–carnivore conflict and perspectives on carnivore management worldwide. *Conservation Biology* 17:1491–1499.
- Valeix M, Hemson G, Loveridge AJ, Mills G, Macdonald DW.** 2012. Behavioural adjustments of a large carnivore to access secondary prey in a human-dominated landscape. *Journal of Applied Ecology* 49:73–81.
- Van Cleave EK, Bidner LR, Ford AT, Caillaud D, Wilmers CC, Isbell LA.** 2018. Diel patterns of movement activity and habitat use by leopards (*Panthera pardus pardus*) living in a human-dominated landscape in central Kenya. *Biological Conservation* 226:224–237.
- Western D.** 2001. Human-modified ecosystems and future evolution. *Proceedings of the National Academy of Sciences of the United States of America* 98(10):5458–5465.
- WII NBWL [Wildlife Institute of India, National Board for Wildlife].** 2020. *National Wildlife Database*. Dehradun, India: Wildlife Institute of India. Data available upon request through https://wii.gov.in/national_wildlife_database; accessed on 6 October 2020.
- Wilcove DS, Rothstein D, Dubow J, Phillips A, Losos E.** 1998. Quantifying threats to imperiled species in the United States: Assessing the relative importance of habitat destruction, alien species, pollution, overexploitation, and disease. *BioScience* 48:607–615.
- Wittemyer G, Elsen P, Bean WT, Burton ACO, Brashares JS.** 2008. Accelerated human population growth at protected area edges. *Science* 321(5885):123–126.
- WWF [World Wide Fund for Nature].** 2002. *Forest Management Outside Protected Areas*. Position paper. Gland, Switzerland: WWF International.
- Yom-Tov Y, Ashkenazi S, Viner O.** 1995. Cattle predation by the golden jackal *Canis aureus* in the Golan Heights, Israel. *Biological Conservation* 73:19–22.
- Zar JH.** 1999. *Biostatistical Analysis*. 4th edition (1st edition 1974). Upper Saddle River, NJ: Prentice Hall.

Supplemental material

APPENDIX S1 Criteria used to categorize habitat types described in the study area.

APPENDIX S2 Supervised classification accuracy assessment report.

APPENDIX S3 Unsupervised classification accuracy assessment report.

Found at: <https://doi.org/10.1659/MRD-JOURNAL-D-19-00046.1.S1>.