

## **The presence of the coffee berry borer, *Hypothenemus hampei*, in Puerto Rico: fact or fiction?**

Authors: Vega, Fernando E., Franqui, R. A., and Benavides, Pablo

Source: Journal of Insect Science, 2(13) : 1-3

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.002.1301>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## The presence of the coffee berry borer, *Hypothenemus hampei*, in Puerto Rico: fact or fiction?

Fernando E. Vega<sup>1</sup>, R. A. Franqui<sup>2</sup> and Pablo Benavides<sup>3</sup>

<sup>1</sup>Insect Biocontrol Laboratory, USDA, ARS, Bldg. 011A, Beltsville, MD 20705.

<sup>2</sup>Estación Experimental Agrícola, Universidad de Puerto Rico, P. O. Box 21360, Río Piedras, PR 00928

<sup>3</sup>Department of Entomology, Purdue University, 1158 Smith Hall, West Lafayette, IN 47906  
[vegaf@ba.ars.usda.gov](mailto:vegaf@ba.ars.usda.gov)

Received 28 May 2002, Accepted 18 June 2002, Published 15 July 2002

### Short Communication

The coffee berry borer, *Hypothenemus hampei* Ferrari (Coleoptera: Scolytidae), is widely considered to be the most devastating pest of coffee. Endemic to Central Africa, the coffee berry borer can now be found in most coffee growing regions throughout the world (Le Pelley 1968). Annual losses caused by this insect have been estimated at over \$500 million annually (P. Baker, CABI Bioscience, UK; personal communication). One of the most widely cited references on coffee berry borer presence in different countries is that of Le Pelley (1968), where more than 20 countries, including Puerto Rico, are listed as having the insect.

A survey for the coffee berry borer in Puerto Rico in the major coffee growing areas of Lares, Utuado, and Adjuntas, in the summer of 1998 by F. E. Vega, G. Mercadier and N. Bayron Justiniano, followed by another survey of coffee samples from Utuado, Adjuntas, Lares, Yauco, and Maricao in June of 2002 by F. E. Vega, A. Sidor, and R. Franqui, indicated that the insect was not present on the island. This was interesting because if the insect had indeed been reported in the island, then its eventual eradication would be noteworthy. We therefore decided to investigate the origins of statements indicating the insect had been reported in Puerto Rico. Le Pelley (1968) cited a report by Roque (1946) which reads as follows:

“Status of infestation of coffee by the coffee bean borer *Stephanoderes* sp. - The percentage of infestation of coffee berries at the two farms on the Lares-Adjuntas road where the pest was first noted has not increased since last year. At the Federal Experiment Station in Mayagüez, the infestation has been greatly reduced by a very thorough collection of the berries. Two new infestations have been discovered at the Demonstration Farm of the Agricultural Extension Service, in San Sebastián, and at three farms on the Lares-Utuado road. In

the newly discovered infestations, the percentage of infested berries is very low. All infestations discovered so far have been in *Coffea excelsa*. None in *Coffea arabica*, although under laboratory infestation of *C. arabica* has taken place.”

The accuracy of this report was questioned by Bergamin (1946), due to Roque referring to the insect as *Stephanoderes* sp. without specifying the species (*hampei*) whose damage and importance was quite well known at the time among coffee producing countries. In addition, he found it odd that the reported infestations only occurred in *Coffea dewevrei* Wildem & T. Dur. (= *excelsa*) which is known to be less preferred by the insect than *C. arabica* L. and *C. canephora* Pierre ex Froehner (= *robusta*) (Le Pelley 1968).

We have located an even earlier report (Nolla 1944) which reads as follows:

“The coffee bean beetle. - The coffee bean-beetle has been reported from the Lares region. An infestation of 20 per cent was found this year in berries that had dried on trees in the Excelsa variety. It was found, however that the beetles do not continue to breed in these dried berries. Three months after collection all beetles had died. In Excelsa coffee the crops overlap-berries and various stages of development are found throughout the year. In the Arabian coffee all berries ripen more or less at the same time. To survive unless it has some other host, the insect requires the conditions offered by Excelsa coffee. No other host has yet been found. It may be found advisable to recommend the destruction of all Excelsa trees as a means of preventing the increase and spread of the insect.”

This report is noteworthy because (1) there is no taxonomic

identification of the insect whatsoever; (2) a non-traditional name for the insect is used (the coffee bean beetle); (3) the insect is reportedly found on *Coffea dewevrei* (= *excelsa*) and not on *C. arabica* which is known to be more susceptible to attack and “is used almost exclusively in Puerto Rico” (Vicente Chandler *et al.* 1968; see also Martorell 1976); and (4) the coffee berry borer can survive for more than three, and even up to eight months on dried or overripe berries (Baker 1999). Neither Nolla (1944) nor Roque (1946) specify whether the insect was found feeding on the seed, which is a unique trait of *H. hampei*.

The evidence indicates that the insect reported by Nolla (1944) and Roque (1946) in coffee berries in Puerto Rico could not be *H. hampei*. We found additional evidence after finding a 1948 report by Wolcott in which he recounts that the insects collected from coffee berries in 1941 were identified by M. W. Blackman, a Scolytidae expert with the U. S. Department of Agriculture, as *Stephanoderes* “near, but not *hampei* Ferrari” (Wolcott 1948). The insect reported by Nolla (1944) and Roque (1946) might have been another species of Scolytidae, *H. seriatus* (Eichoff), which is known as the “false coffee berry borer” and can enter the coffee berry, but does not feed on the seed (Fonseca 1938). Unfortunately, we were not able to locate the specimens identified by Blackman in the Coleoptera Collection at the National Collections of Insects and Mites of the National Museum of Natural History, Smithsonian Institution nor at the Museo de Entomología y Biodiversidad Tropical of the Agricultural Experiment Station, University of Puerto Rico.

Our findings indicate that the coffee berry borer has never been present in Puerto Rico and that the citation by Le Pelley is based on a mistaken report. Insect misidentifications create a situation in which scientists make use of a source - in this case Le Pelley’s book - to perpetuate a mistake, even though Bergamin had already questioned the veracity of Roque’s report the same year it was published (1946); Le Pelley’s mistaken attribution was published 20 years later. The consequences of these mistakes could have important implications, such as coffee being quarantined from foreign markets. For example, the Mexican government lists Puerto Rico among those countries from which coffee imports are prohibited (Diario Oficial 1996). Thus, even though the coffee berry borer has been reported in other Caribbean countries, e.g. Jamaica in 1978 (Reid 1983), Cuba in 1994 (Hernández 2002) and the Dominican Republic in 1995 (IICA 1999), so far Puerto Rico has remained free of the insect. This situation might change due to the increased coffee imports from the Dominican Republic and Costa Rica, where the insect is present. In order to prevent the entrance of this devastating insect to Puerto Rico it is imperative that requirements for 10% roasting of imported coffee be strongly enforced in order to kill coffee berry borers that might be present in the coffee seeds, and that coffee be shipped in plastic bags inside cloth bags, as is currently being done. A sampling program for the insect should be implemented in coffee growing regions, especially near roasting factories in Puerto Rico into which imported coffee is being brought. It is also imperative to initiate an extension program aimed at providing training to coffee growers on how to identify and prevent the spread of the insect.

## Acknowledgments

We thank Francisco Infante (El Colegio de la Frontera Sur, Tapachula, Mexico) for comments on an earlier draft of this manuscript and Natalia J. Vandenberg at the National Collections of Insects and Mites of the National Museum of Natural History, Smithsonian Institution, for her help in trying to locate specimens in the Coleoptera collection. Also, Evelio Hernández López and Wigmar González (Agricultural Experiment Station, Adjuntas, Puerto Rico), Miguel Monroig (Coffee Specialist, Agricultural Extension Service, University of Puerto Rico) and Alfonso Pérez and Juan E. Pérez Nieves (Administration for Agricultural Services and Development, Adjuntas, Puerto Rico) for their cooperation and hospitality during our recent visit to Puerto Rico.

## References

- Baker, P. S. 1999. The coffee berry borer in Colombia. Final report of the DFID- Cenicafé - CABI Bioscience IPM for coffee project. Chinchiná (Colombia), DFID - Cenicafé. 154 pp.
- Bergamin, J. 1946. A “broca do café” em Porto Rico? *Boletim da Superintendência dos Serviços do Café* 21:340.
- Diario Oficial. 1996. Norma Oficial Mexicana NOM-019-FITO-1995, por la que se establece la cuarentena exterior para prevenir la introducción de plagas del café. *Diario Oficial, Primera Sección*, Martes 10 de diciembre de 1996. Available at: <http://ns1.oirsa.org.sv/DI07/Di0703/Di070301/Di07030102/NORMA19.htm>
- Fonseca, J. P. 1938. A broca verdadeira e a falsa broca do café. *Revista do Instituto do Café* 13:63-64.
- Hernández, L. R. 2002. El bumerang maldito. *Encuentro en la Red: Diario Independiente de Asuntos Cubanos*, Año III, Edición 278, Lunes 14 de Enero de 2002. Available at: <http://www.cubaencuentro.com/ecologia/2002/01/14.html>.
- IICA. 1999. Instituto Interamericano de Cooperación para la Agricultura - Agencia de Cooperación Técnica en la Republica Dominicana: Resumen Histórico de los Proyectos y Actividades Desarrollados para la Agencia de Cooperación Técnica del IICA en Republica Dominicana, Santo Domingo, Republica Dominicana. Available at: <http://www.iicard.org/espanol/30%20Anos/default.htm>
- Le Pelley, R. H. 1968. *Pests of Coffee*. Longmans, Green and Co., Ltd., London. 590 pp.
- Martorell, L. F. 1976. *Annotated Food Plant Catalog of the Insects of Puerto Rico*. Agricultural Experiment Station, University of Puerto Rico, 303 pp.
- Nolla, J. A. B. 1944. *Annual Report for the Fiscal Year 1942-43*, University of Puerto Rico, Agricultural Experiment Station, Río Piedras, Puerto Rico.
- Reid, J. C. 1983. Distribution of the coffee berry borer (*Hypothenemus hampei*) within Jamaica, following its discovery in 1978. *Tropical Pest Management* 29:224-230.
- Roque, A. 1946. *Annual Report for the Fiscal Year 1943-44*, University of Puerto Rico, Agricultural Experiment Station, Río Piedras, Puerto Rico.

Vega FE, Franqui RA, Benavides P. 2002. The presence of the coffee berry borer, *Hypothenemus hampei*, in Puerto Rico: fact or fiction? 3 pp. *Journal of Insect Science*, 2.13. Available online: [insectscience.org/2.13](https://insectscience.org/2.13)

Vicente Chandler J, F. Abruña, R. Bosque Lugo, and S. Silva. 1968. Intensive coffee culture in Puerto Rico. Agricultural Experiment Station, University of Puerto Rico, Bulletin 211.

Wolcott, G. N. 1948. The Insects of Puerto Rico. The Journal of Agriculture of the University of Puerto Rico Vol. 32, No. 2.