

South American Spider Mites: New Hosts and Localities

Authors: Mendonça, Renata S, Navia, Denise, Diniz, Ivone R, and Flechtmann, Carlos HW

Source: Journal of Insect Science, 11(121) : 1-18

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.011.12101>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



South American spider mites: New hosts and localities

Renata S Mendonça^{1a*}, Denise Navia^{2b}, Ivone R Diniz^{1c}, Carlos HW Flechtman^{3d}

¹ Universidade de Brasília, UNB, Instituto de Ciências Biológicas, Departamento de Zoologia, Campus Universitário Darcy Ribeiro, ICC Sul, Sala AT-159, Asa Norte, 70910-900, Brasília-DF, Brazil; Université Montpellier II, Montpellier, France

² Embrapa Recursos Genéticos e Biotecnologia, Laboratório de Quarentena Vegetal, Parque Estação Biológica, W5 Norte Final, Caixa Postal 02372, 70.770-900 Brasília-DF, Brazil

³ CNPq Researcher, Escola Superior de Agricultura Luiz de Queiroz (ESALQ), Universidade de São Paulo, Cx. Postal 09, 13.418-900, Piracicaba-SP, Brazil

Abstract

In order to contribute to taxonomic information on Tetranychid mites (Acari: Tetranychidae) in South America, surveys were conducted in Brazil (15 States and the Federal District) and Uruguay (one Department); 550 samples of 120 plant species were collected. Tetranychid mite infestations were confirmed in 204 samples, and 22 species belonging to seven genera of the Bryobiinae and Tetranychinae subfamilies were identified on 58 different host plants. Thirty-six new plant hosts were found in Brazil, South America, and worldwide for the following species: *Eutetranychus banksi* (McGregor); *Mononychellus tanajoa* (Bondar); *Oligonychus anonae* Paschoal; *O. mangiferus* (Rahman and Sapro); *Tetranychus bastosi* Tuttle, Baker and Sales; *T. desertorum* Banks, 1900, *T. evansi* Baker and Pritchard; *T. ludeni* Zacher; *T. mexicanus* (McGregor); *T. neocaledonicus* André; and *T. urticae* Koch. Four new localities in Brazil were reported for *Eotetranychus tremae* De Leon; *O. anonae*; *Panonychus ulmi* (Koch); and *T. gloveri* Baker and Pritchard.

Keywords: Eurytetranychini, Hystrichonychini, neotropical region, systematics, taxonomy, Tetranychini

Correspondence: ^a renatasm@cenargen.embrapa.br, ^b navia@cenargen.embrapa.br, ^c irdiniz@unb.br, ^d chwflech@esalq.usp.br. * Corresponding author

Received: 16 September 2010, **Accepted:** 14 November 2010

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 11, Number 121

Cite this paper as:

Mendonça RS, Navia D, Diniz IR, Flechtman CHW. 2011. South American spider mites: New hosts and localities. *Journal of Insect Science* 11:121 available online: insectscience.org/11.121

Introduction

The Tetranychidae Donnadieu family includes a large number of strictly phytophagous mites; a few species are significant agricultural crop pests worldwide (Jeppson et al. 1975; Helle and Sabelis 1985).

The number of described tetranychids remained stable for 75 years and then increased when their economic incidence for agriculture became more significant (Bolland et al. 1998). In 1950, McGregor initially listed 102 species in 15 genera. Five years later, this number increased to 204 species in 18 genera (Pritchard and Baker 1955). In 1998 there were 1189 species in 71 genera registered in the World Catalogue of the Spider Mite Family (Acari: Tetranychidae) (Bolland et al. 1998), and currently there are approximately 1257 species in 76 genera listed on the Spider Mite Web, which is a comprehensive database on the Tetranychidae (Migeon and Dorkeld 2006).

In Brazil, taxonomic studies on the Tetranychidae were initially conducted between 1920 and 1930. Among the early findings was the report of *Tetranychus gloveri* Banks 1900 and the description of *Mononychellus tanajoa* (Bondar 1938) based on specimens collected in Brazil by G Bondar, a Russian researcher living in the state of Bahia (Bondar 1930, 1938). In the 1960s numerous contributions were made by pioneers in agricultural mite studies in Brazil, such as CHW Flechtmann (Flechtmann 1967, 1967a, 1967b, 1972, 1975, 1976; Flechtmann and Baker 1970, 1975) and AD Paschoal (Paschoal 1970, 1970a), followed by RJF Feres in subsequent decades (Feres 1986, 1992; Feres and Flechtmann 1986, 1986a, 1988, 1995, 1995a, 1995b). From that time

forward these and other researchers have conducted major studies on tetranychids in several fields such as taxonomy, ecology, biology, control, and resistance, bringing important advances to knowledge of Brazilian tetranychids. Currently 185 of the 1257 described tetranychid species are reported in South America, of which 104 are found in Brazil (Bolland et al. 1998; Migeon and Dorkeld 2006).

The most intensive search for Tetranychidae mites in Brazil, the largest country in South America, has been centered in the southeast region and is usually associated with economically important crops. Agriculture continues to expand, approaching biological reserves and other regions in the country. It is important to continue tetranychid research in previously unexplored areas in hopes of increasing our understanding of this family. This paper presents new information about the occurrence and distribution of Tetranychidae in five Brazilian regions and in Uruguay, on plants ranging from ornamental, fruit-bearing, vegetable, grass, oilseed, to large crops species. New hosts and localities for tetranychid mites in Brazil, South America, and the world are reported.

Materials and Methods

A legal permit to collect zoological material for scientific purposes is required by the **Brazilian Institute** for the **Environment** and **Renewable** Natural Resources (IBAMA). A permit was requested and permit number 231/2006 CGFAU/IBAMA was issued.

Mite collection

Mites were collected between October 2004 and July 2008. A total of 550 samples were collected from several host plants found in




Figure 1. States and their respective sampling points for collecting tetranychid mites in Brazil between October 2004 and July 2008. ARCGIS 9.0 (ESRI, www.esri.com). Latitude and longitude coordinates for the single sample point from Uruguay were 58.1312 S and 32.6368 W. The sample point was not mapped. High quality figures are available online.

five regions in Brazil that include the states of Acre, Bahia, Ceará, Espírito Santo, Goiás, Minas Gerais, Mato Grosso do Sul, Paraná, Pernambuco, Rio Grande do Norte, Rio Grande do Sul, Roraima, Santa Catarina, São Paulo, Sergipe, and the Federal District (Figure 1; Appendix 1). One additional collection was conducted in Uruguay. Plants were collected when symptoms of tetranychid attacks were observed, according to Moraes and Flechtman (2008). The geographical points of collection were logged into a Garmin 12 GPS using WGS84 (World Geodetic System 84) datum; maps were created using ARCGIS 9.0 (ESRI, www.esri.com).

Inspecting and processing plant samples

Mites were collected from the plants using a washing/sieving extraction method (Miranda et al. 2007). The resulting 70% ethanol solution was inspected under Stemi SV6 (www.zeiss.com) and Olympus SZX 122 (www.olympus.com) stereomicroscopes

connected to a Highlight 3000-1 transilluminator at 50× magnification. Finally, Tetranychidae mites were directly mounted on microscopic slide preparations in Hoyer's medium. From each population, 25 females were mounted in dorso-ventral position and ten isolated males were mounted in lateral position to identify the species by morphological analysis. When only a few males were present in a sample, all of them were slide-mounted.

Identifying mite species

Microscopic slide preparations of specimens were examined under phase contrast (Leitz Wetzlar, www.leica-microsystems.com) and interference microscopes (Nikon Eclipse 80i, www.nikon.com) in 40× and 100× objectives and. Morphological identification was conducted by examining relevant taxonomic characteristics of male and female Tetranychidae systematics. The shape of the aedeagus, or male genitalia, was used to classify the species (Pritchard and Baker 1955, Meyer 1974, 1987, Baker and Tuttle 1994, Ehara, 1999), which were then compared to specimens that had been deposited in the Reference Mite Collection of the Laboratory of Plant Quarantine, Embrapa Genetic Resources and Biotechnology, Brasília, Brazil. Materials from this study were deposited in this collection as voucher specimens.

Results and Discussion

Tetranychid mites were found on 37.1% (n = 204) of the 550 plant material samples that were collected (Appendix 1). Twenty-one species of Tetranychinae and one of Bryobiinae were identified on the 204 samples from different localities and inspected hosts (Table 1).

Table 1. Tetranychid mite species (Tetranychinae) collected in Brazil between October 2004 and July 2008.

Species	Localities	Hosts	Collections
1. <i>Eutetranychus banksi</i> (McGregor, 1914)	10	10	13
2. <i>Eotetranychus tremae</i> De Leon, 1957	1	1	1
3. <i>E. smithi</i> Pritchard & Baker, 1955	1	1	1
4. <i>Monoceronychus ichencensis</i> Mendonça, Navia & Flechtmann, 2010	1	1	1
5. <i>Mononychelhus planki</i> (McGregor, 1950)	18	5	26
6. <i>M. tanajoa</i> (Bondar, 1938)	17	4	23
7. <i>Oligonychus ananae</i> Paschoal, 1970	1	1	1
8. <i>O. ilicis</i> (McGregor, 1917)	1	1	1
9. <i>O. fileno</i> Mendonça, Navia & Flechtmann, 2010	1	1	1
10. <i>O. mangiferus</i> (Rahman & Sapra, 1940)	2	3	3
11. <i>O. aff. mangiferus</i>	1	1	1
12. <i>Panonychus ulmi</i> (Koch, 1836)	1	1	1
13. <i>Tetranychus abacae</i> Baker & Pritchard, 1962	1	1	1
14. <i>T. bastosi</i> Tuttle, Baker & Sales, 1977	4	5	5
15. <i>T. desertorum</i> Banks, 1900	15	11	19
16. <i>T. evansi</i> Baker & Pritchard, 1960	1	2	2
17. <i>T. gloveri</i> Baker & Pritchard, 1962	1	1	1
18. <i>T. ludeni</i> Zacher, 1913	15	10	20
19. <i>T. mexicanus</i> (McGregor, 1950)	14	12	23
20. <i>T. neocaledonicus</i> André, 1933	4	5	5
21. <i>T. ogmophallos</i> Ferreira & Flechtmann, 1997	1	1	1
22. <i>T. urticae</i> Koch, 1836	55	34	101

* The columns do not equal a total of 207 collections due to repeated hosts in some localities.

Several samples contained more than one species of tetranychid on the same leaf. This occurred on bean (*T. urticae* and *Eutetranychus banksi*; *T. urticae* and *M. planki*; *T. urticae*, *T. ludeni*, and *T. desertorum*; *T. desertorum* and *M. planki*; *T. ludeni* and *M. planki*), soybean (*T. urticae* and *M. planki*; *T. urticae* and *T. desertorum*; *T. urticae*, *T. desertorum*, and *M. planki*; *T. ludeni* and *M. planki*; *T. desertorum* and *M. planki*), cotton (*T. urticae* and *M. planki*; *T. ludeni* and *T. mexicanus*), okra (*T. neocaledonicus* and *M. planki*), and eggplant (*T. urticae* and *M. tanajoa*). *T. mexicanus*, *P. ulmi*, *Oligonychus aff. mangiferus*, and *O. fileno* were found on grapes in the municipality of Pirapora. Foott (1962, 1963) reported phytophagous mite species coexisting on a single host. The fact that different species coexist on a single host requires careful sampling and collecting larger numbers of specimens in microscopic preparations. This allows for greater representation of mite fauna and increases the probability of identifying all Tetranychidae species present on a specific host and/or at a specific locality.

Table 2. New hosts and localities for tetranychid mites species collected in Brazil and Uruguay between October 2004 and July 2008 with information on collection sites, host plants, collecting dates, and collectors.

Collection sites				Voucher N°	Host plant			Date	Mite Species	Collectors
Country & State	LONG.	LAT.	Municipality		Common name	Scientific name	Family			
BR BA	-39.1019	-12.6700	Cruz das Almas	139	siratro	<i>Macropodium atropurpureum</i> (L.) #	Fabaceae	09.VI.2006	<i>T. urticae</i> #	A Noronha
BR DF	-47.8864	-15.8063	Brasília	200	red mulberry	<i>Morus rubra</i> L. *	Moraceae	23.VI.2008	<i>T. neocaledonicus</i> *	RS Mendonça
BR DF	-47.9167	-15.7833	Brasília	85	sweet basil	<i>Ocimum basilicum</i> L. *	Lamiaceae	23.VII.2006	<i>T. neocaledonicus</i> *	RS Mendonça
BR DF	-48.1405	-15.9423	Gama	30	butter bean	<i>Phaseolus lunatus</i> L. *	Fabaceae	07.XII.2005	<i>T. urticae</i> +, <i>E. banksi</i> *	RS Mendonça
BR DF	-47.9443	-15.9285	Núcleo Bandeirantes	17	bean	<i>Phaseolus vulgaris</i> L. *	Fabaceae	23.XI.2005	<i>M. tanajoa</i> *	RS Mendonça
BR DF	-47.4888	-15.5224	Planaltina	11	gilo	<i>Solanum gilo</i> Radlk. *	Solanaceae	22.XI.2005	<i>T. desertorum</i> *, <i>E. banksi</i> *	RS Mendonça
BR DF	-47.7222	-15.5299	Planaltina	27	purple granadilla	<i>Passiflora edulis</i> Sims *	Passifloraceae	06.XII.2005	<i>M. tanajoa</i> *	RS Mendonça
BR DF	-48.1180	-15.9087	Recanto das Emas	20	eggplant	<i>Solanum melongena</i> L. *	Solanaceae	28.XII.2005	<i>T. desertorum</i> +, <i>E. banksi</i> *	RS Mendonça
BR DF	-48.1181	-15.9087	Recanto das Emas	21	yellow magnolia	<i>Michelia champaca</i> L. *	Magnoliaceae	28.XII.2005	<i>T. desertorum</i> *	RS Mendonça
BR DF	-48.1182	-15.9087	Recanto das Emas	24	red-ginger	<i>Alpinia purpurata</i> Schum. *	Zingiberaceae	28.XI.2005	<i>T. urticae</i> *	RS Mendonça
BR GO	-47.8367	-16.3443	Luiziana	33	castor oil plant	<i>Ricinus communis</i> L. *	Euphorbiaceae	19.XII.2005	<i>O. mangiferus</i> **	RS Mendonça
BR MG	-43.6744	-15.3383	Juba	84	custard-apple	<i>Annona muricata</i> L. *	Annonaceae	08.IV.2006	<i>O. ananae</i> +, <i>O. ananae</i> *	D Navia LC Miranda RS Mendonça
BR MG	-46.3775	-17.6602	Lagoa Grande	82	purple granadilla	<i>Passiflora edulis</i> Sims *	Passifloraceae	07.IV.2006	<i>E. banksi</i> *	D Navia LC Miranda RS Mendonça
BR MG	-45.0028	-21.2417	Lavras	109	European plum	<i>Prunus domestica</i> L. *	Rosaceae	09.V.2006	<i>T. ludeni</i> +	D Navia LC Miranda RS Mendonça
BR MG	-45.0028	-21.2417	Lavras	110	lemandarin	<i>Citrus limonia</i> L. Osbeck *	Rutaceae	09.V.2006	<i>T. ludeni</i> *	D Navia LC Miranda RS Mendonça
BR MG	-45.0028	-21.2417	Lavras	115	flame-flower	<i>Pyrostegia venusta</i> Miq. *	Bignoniaceae	09.V.2006	<i>T. urticae</i> *	D Navia LC Miranda RS Mendonça
BR MG	-45.0028	-21.2417	Lavras	118	yellow magnolia	<i>Michelia champaca</i> L. *	Magnoliaceae	09.V.2006	<i>T. desertorum</i> +, <i>E. banksi</i> *	D Navia LC Miranda RS Mendonça
BR MG	-43.3006	-15.8025	Nova Porteirinha	98	tree cotton	<i>Gossypium barbadense</i> L. *	Malvaceae	10.IV.2006	<i>T. bastosi</i> *	D Navia LC Miranda RS Mendonça
BR MG	-46.4608	-18.5823	Patos de Minas	40	geranium	<i>Polargonum hortorum</i> Bailey *	Geraniaceae	19.XII.2005	<i>T. ludeni</i> +	RS Mendonça
BR MG	-44.9419	-17.3450	Pirapora	101	grape	<i>Vitis vulpina</i> L. *	Vitaceae	12.IV.2006	<i>T. mexicanus</i> *, <i>T. ulmi</i> *	D Navia LC Miranda RS Mendonça
BR MG	-43.6851	-20.3294	Recanto Novo	135	jessamine	<i>Cestrum intermedium</i> Sandw. *	Solanaceae	13.V.2006	<i>T. neocaledonicus</i> *	D Navia LC Miranda RS Mendonça
BR MG	-43.6851	-20.3294	Recanto Novo	136	pomegranate	<i>Punica granatum</i> L. *	Punicaceae	13.V.2006	<i>Oligonychus mangiferus</i> **	D Navia LC Miranda RS Mendonça
BR MG	-43.6851	-20.3294	Recanto Novo	137	maissenda	<i>Mussaenda sp.</i> *	Rubiaceae	13.V.2006	<i>O. mangiferus</i> *	D Navia LC Miranda RS Mendonça
BR MG	-45.2071	-18.2400	Três Marias	121	coral-tree	<i>Erythrina ilicifolia</i> Lam. *	Fabaceae	09.IV.2006	<i>T. mexicanus</i> +	D Navia LC Miranda RS Mendonça
BR MG	-42.8819	-20.7539	Vigosa	119	fire dragon plant	<i>Aclypha wilkesiana</i> Mull. Arg. *	Euphorbiaceae	13.V.2006	<i>E. tremula</i> *	D Navia LC Miranda RS Mendonça
BR PE	-34.9117	-7.9458	Recife	92	papaya	<i>Carica papaya</i> L. *	Caricaceae	11.VIII.2006	<i>T. evansi</i> *	D Navia LC Miranda RS Mendonça
BR PE	-34.9165	-7.8239	Recife	95	wattle	<i>Acacia sp.</i> #	Fabaceae	11.VIII.2006	<i>E. banksi</i> #	D Navia LC Miranda RS Mendonça
BR RN	-37.3442	-5.1875	Mossoró	148	papaya	<i>Carica papaya</i> L. *	Caricaceae	05.VII.2006	<i>T. gloveri</i> *	Ehom
BR RR	-40.9833	1.9333	Boa Vista	177	purple granadilla	<i>Passiflora edulis</i> Sims *	Passifloraceae	17.V.2007	<i>T. urticae</i> +	D Navia
BR RR	-40.6624	1.6454	Boa Vista	154	watermelon	<i>Citrullus lanatus</i> (Thunb.) *	Cucurbitaceae	29.III.2005	<i>T. ludeni</i> +	D Navia
BR RR	-40.6733	1.9333	Boa Vista	156	watermelon	<i>Citrullus lanatus</i> (Thunb.) *	Cucurbitaceae	12.III.2005	<i>T. ludeni</i> +	D Navia
BR RR	-40.6878	1.7335	Boa Vista	162	watermelon	<i>Citrullus lanatus</i> (Thunb.) *	Cucurbitaceae	30.III.2005	<i>T. ludeni</i> +	D Navia
BR RS	-56.5531	-29.1253	Itaqui	187	perennial ragweed	<i>Ambrosia polysyncha</i> DC. #	Asteraceae	29.IX.2007	<i>T. urticae</i> +	D Navia
BR RS	-52.4000	-28.2667	Passo Fundo	161	wheat	<i>Triticum vulgare</i> L. *	Poaceae	27.IX.2006	<i>T. urticae</i> **	E Shrica
BR RS	-53.8333	-29.6833	Santa Maria	185	wild quinine	<i>Parthenium sp.</i> *	Asteraceae	26.IX.2007	<i>T. urticae</i> **	D Navia
BR SP	-47.0760	-22.8665	Campinas	48	velvet bean	<i>Mucuna uteraria</i> Holland *	Fabaceae	10.I.2006	<i>T. desertorum</i> +, <i>E. banksi</i> *	RS Mendonça
BR SP	-47.3659	-24.2649	Miracatu	56	gilo	<i>Solanum gilo</i> Radlk. *	Solanaceae	16.I.2006	<i>T. desertorum</i> +, <i>E. banksi</i> *	RS Mendonça
UY RN	-58.1312	-32.6368	Rio Negro, San Javier	195	lantana	<i>Lantana camara</i> L. *	Verbenaceae	06.VI.2007	<i>T. urticae</i> -	D Navia E Castiglione

New reports: * = New host for species in the world; ** = First report as host in Neotropical region; + = First reported as host in South America; # First reported as host in Brazil. Species names marked in red indicate new localities in Brazil.

New hosts for Tetranychidae mites in Brazil, South America, and the world

New hosts were found for 11 tetranychid mite species: *E. banksi*, *M. tanajoa*, *O. ananae*, *O.*

mangiferus, *T. bastosi*, *T. desertorum*, *T. evansi*, *T. ludeni*, *T. mexicanus*, *T. neocaledonicus*, and *T. urticae* (Table 2). The new hosts are listed below according to mite species and host plant family.

Tetranychinae Berlese, Eurytetranychini Reck

Eutetranychus banksi

(McGregor 1914)

Tetranychus banksi McGregor, 1914. Type-host: *Ricinus communis* L. Type-locality: Orlando, Florida, United States.

Fabaceae

Acacia sp., (acacia), Universidade Federal Rural de Pernambuco, UFRPE, Recife, Pernambuco, 11.VIII.2006.

Mucuna aterrima Holland (black mucuna), Instituto Agrônômico de Campinas, ICA, Campinas, São Paulo, 10.I.2006.

Phaseolus lunatus L. (lima beans), Embrapa Hortaliças, Gama, Distrito Federal, 07.XII.2005.

The presence of *E. banksi* on *Acacia* sp. was previously reported in Colombia (Urueta 1975). The occurrence on this host from Pernambuco is the first in Brazil.

McGregor (1914) described *E. banksi* on *M. pruriens* in the United States and Livshits and Salinas-Croche (1968) located it on *Mucuna* sp. in Cuba. Garret et al. (1967) reported *E. banksi* infestations on *P. lunatus* in Hawaii. This is the first report of *E. banksi* on *M. aterrima* and *P. lunatus* in South America.

Magnoliaceae

Michelia champaca L., (yellow magnolia), Faculdade da Terra, Recanto das Emas, Federal District, 28.XI.2005.

Passifloraceae

Passiflora edulis Sims (sour passion fruit), Lagoa Grande, Minas Gerais, 07.IV.2006.

Solanaceae

Solanum gilo Raddi (scarlet eggplant), Pípiripau, Planaltina, Distrito Federal, 22.XI.2005 and Caturra farm, Vale do Ribeira, Miracatu, São Paulo, 16.I.2006. Caturra farm, Miracatu, Vale do Ribeira, São Paulo, 16.I.2006.

Solanum melongena L. (eggplant), Faculdade da Terra, Recanto das Emas, Federal District, 28.XI.2005.

Michelia champaca, *P. edulis*, *S. gilo* and *S. melongena* are new hosts for *E. banksi* in the world.

Tetranychinae Berlese, Tetranychini Reck

Mononychellus tanajoa

(Bondar 1938)

Tetranychus tanajoa Bondar, 1938. Type-host: *Manihot utilissima* and *M. aipim*. Type-locality: Bahia, Brazil.

Fabaceae

Phaseolus vulgaris L. (beans), Vargem Bonita, Núcleo Bandeirantes, Distrito Federal, 23.XI.2005.

This is the first report of *M. tanajoa* infesting beans (*P. vulgaris*) under field conditions. This mite mainly infests species from the *Manihot* genus (Euphorbiaceae) and is also reported on plants from the Asteraceae,

Caesalpinaceae, Curcubitaceae, Passifloraceae, Malvaceae, Rubiaceae, and Solanaceae families in the northeastern region of Brazil (Tuttle et al. 1977; Moraes et al. 1995). Plants of the Fabaceae family were cited as hosts for the *M. tanajoa* in Brazil (*Canavalia Braziliensis* Mart. ex Benth., *Macroptilium martii* Benth) (Moraes et al. 1995), in Central America (*Erythrina* sp., *Gliricidia maculata* Kunth) (Andrews and Poe 1980; Gutierrez 1986; Bolland et al. 1998, Migeon and Dorkeld 2006), and Mexico ([*Gliricidia sepium* (Jacq.) Kunth)]) (Tuttle et al. 1976).

Passifloraceae

Passiflora edulis Sims (yellow passion fruit), Faculdades Integradas da União Pioneira de Integração Social, UPIS, Planaltina, Distrito Federal, 06.XII.2005.

Passiflora edulis is a new host for *M. tanajoa*. This mite was previously reported on *P. cincinnata* Mart. by Moraes et al. (1995) in the northeast region of Brazil (Moraes et al. 1995). The passion fruit plants (*P. edulis*) and the bean plants (*P. vulgaris*) were near a cassava plantation (*Manihot esculenta* Crantz) that was highly infested with *M. tanajoa*. According to Moraes et al. (1995), high levels of *M. tanajoa* in cassava plantations can cause to the species to be dispersed by the wind to nearby plants and temporarily infest alternative hosts.

Oligonychus anonae

Paschoal 1970

Oligonychus anonae Paschoal, 1970. Type-host: *Annona muricata* L. Type-locality: Jaboticabal, Brazil.

Annonaceae

Annona squamosa L. (sweetsop), in the Jaíba, Mucambinho Project, Minas Gerais, 08.IV.2006.

This species was reported in São Paulo on soursop (*A. muricata*) (Annonaceae) (Paschoal 1970) and was collected from Lauraceae plants (*Persea americana* Mill.) and Vitaceae (*Vitis vinifera* L.) in Brazil (Paschoal 1970a). *Annona squamosa* (sweetsop) is a new host for the *O. anonae*.

Oligonychus mangiferus

(Rahman and Sapro 1940)

Paratetranychus mangiferus Rahman and Sapro, 1940. Type-host: *Mangifera indica* L. Type-locality: Pakistan.

Euphorbiaceae

Ricinus communis L., (castor oil plant), BR 040, km 40, Luziânia, Goiás, 19.XII.2005.

Punicaceae

Punica granatum L. (pomegranate), Recanto Novo, Minas Gerais, 13.V.2006.

Rubiaceae

Mussaenda sp. (pink mussaenda), Recanto Novo, Minas Gerais 13.V.2006.

This is the first report of *O. mangiferus* on castor oil plants and pomegranates in the Neotropics. This mite was previously found on these host plants in India (Gupta 1976, Gupta and Gupta 1994). *Mussaenda* L. is reported for the first time as a host for *O. mangiferus*.

Tetranychus bastosi

Tuttle, Baker, and Sales 1977

Tetranychus (Tetranychus) bastosi Tuttle, Baker, and Sales 1977. Type-host: *Morus rubra* L. Type-locality: Crato, Brazil.

Malvaceae

Gossypium barbadense L. (wild cotton), Empresa de Pesquisa Agropecuária de Minas Gerais – EPAMIG, Nova Porteirinha, Minas Gerais, 10.IV.2006.

This is first report of *T. bastosi* on cotton (*G. barbadense*). This species was observed previously on *Malva rotundifolia* L. (Tuttle et al. 1977).

Tetranychus desertorum

Banks 1900

Tetranychus desertorum Banks, 1900. Type-host: *Larrea tridentata* J. M. Coult, *Phacelia crenulata* Torr. ex S. Watson. Type-locality: Mesilla, United States.

Fabaceae

Mucuna aterrima Holland (black mucuna), Instituto Agrônomo de Campinas, ICA, Campinas, São Paulo, 10.I.2006.

Baker and Pritchard (1962) reported *T. desertorum* on *M. pruriens* in Central America. Therefore, *M. aterrima* is a new registered host in the Fabaceae family for the *T. desertorum* in South America.

Magnoliaceae

Michelia champaca L. (yellow magnolia), Faculdade da Terra de Brasília, Recanto das Emas, Distrito Federal, 28.XI.2005 and the Universidade Federal de Lavras, Lavras, Minas Gerais, 06.V.2006.

Michelia champaca is a new host for *T. desertorum*.

Solanaceae

Solanum melongena L. (eggplant), Faculdade da Terra, Recanto das Emas, Distrito Federal, 28.XI.2005.

Solanum gilo Raddi (scarlet eggplant), Pípiripau, Planaltina, Distrito Federal, 22.XI.2005 and Caturra farm, Vale do Ribeira, Miracatu, São Paulo, 16.I.2006. Caturra farm, Miracatu, Vale do Ribeira, São Paulo, 16.I.2006.

This is the first report of *T. desertorum* on eggplant (*S. Melongena*) in South America previously registered in Japan by Ehara (1956). Infestations of *T. desertorum* in Brazil were registered on *Acnistus cauliflorus* (Flechtmann 1967),; *Brugmansia suaveolens* (Furtado et al. 2006),; *Brunfelsia* sp, (Flechtmann 2004),; *Lycopersicon esculentum* (Flechtmann 1967), and *Solanum tuberosum* (Paschoal 1970a).

Solanum gilo is reported as a new host for *T. desertorum*.

Tetranychus evansi

Baker and Pritchard 1960

Tetranychus evansi Baker and Pritchard,, 1960. Type-host: *Lycopersicon esculentum* L. Type-locality: Mauritius. (Indian Ocean).

Caricaceae

Carica papaya L. (papaya), Mumbecas farm, Recife, Pernambuco, 11.VIII.2006.

Tetranychus evansi was originally described using samples collected from tomato plants (Baker and Pritchard 1960). However, this mite had already been discovered in Brazil by Silva (1954) who described it as *T. marianae* McGregor. Since then it has been reported in the United States (Bolland et al. 1998) and

more recently in Africa, Argentina, China, Spain, France, Greece, Israel, Italy, Puerto Rico, Portugal, and Taiwan (Migeon and Dorkeld 2006). *Tetranychus evansi* is currently reported on 93 host species (Migeon and Dorkeld 2006), most of which belong to the Solanaceae family (Bolland et al. 1998; Moraes et al. 1987). Infestations were reported in Europe of this mite across several plant families (Ferragut and Escudero 1999). However, the only reports on the biology of *T. evansi* are for those found on tomato plants. *Carica papaya* is a new host for the *T. evansi*.

Tetranychus ludeni

Zacher 1913

Tetranychus ludeni Zacher, 1913 (Zacher 1913 *apud* Pritchard and Baker 1955). Type-host: *Cucurbita* sp., *Salvia splendens* Ker Gawl. Type-locality: St. Cloud (near Paris), , France.

Cucurbitaceae

Citrullus lanatus (Thunb.) Matsum and Nakai (watermelon), Embrapa Roraima, Boa Vista, Roraima, 29.III.2005, 12.III.2005 e 30.III.2005.

Geraniaceae

Pelargonium hortorum L. H. Bailey (geranium), Patos de Minas, Minas Gerais, 19.XII.2005.

Rosaceae

Prunus domestica L. (plumb), Universidade Federal de Lavras, Lavras, Minas Gerais, 09.V.2006.

Tetranychus ludeni was previously found on *C. lanatus*, *P. hortorum* and *P. domestica* in Africa, Central America, (El Salvador), Australia, and India (Meyer and Ryke 1959; Davis 1968; Andrews and Poe 1980;

Gutierrez and Schicha 1983; Gupta and Gupta 1994). This is the first report of these hosts in South America.

Rutaceae

Citrus limonia L. (mandarin-lime), Universidade Federal de Lavras, Lavras, Minas Gerais, 09.V.2006.

This is the first report of *Citrus limonia* as a host for *T. ludeni*.

Tetranychus mexicanus

(McGregor 1950)

Tetranychus mexicanus (McGregor 1950). Type-host: *Citrus sinensis* L. Type-locality: Mexico.

Fabaceae

Erythrina indica Lam. (coral tree), Três Marias, Minas Gerais, 09.IV.2006.

This is the first report of *T. mexicanus* infesting the *E. indica* ornamental plant in South America. According to Migeon and Dorkeld (2006), *T. mexicanus* infests these hosts of the Fabaceae family: *Arachis hypogaea* L., *Bauhinia* sp., *Centrosema pubescens* Benth, *Crotalaria retusa* L., *Erythrina poeppigiana* O. F. Cook, and *Phaseolus vulgaris* L.

Vitaceae

Vitis vinifera L. (grapes.), Pirapora, Minas Gerais, 12.IV.2006.

This is the first report of *T. mexicanus* on plants of the Vitaceae family.

Tetranychus neocaledonicus

André 1933

Eotetranychus neocaledonicus André,, 1933.
Type-host: *Gossypium* sp. Type-locality: New Caledonia (Southwest Pacific Ocean).

Lamiaceae

Ocimum basilicum L. (basil), Embrapa Recursos Genéticos e Biotecnologia - Cenargen, Brasília, Distrito Federal, 23.VII.2006.

Moraceae

Morus rubra L. (mulberry), SQS 202 Sul, Brasília, Distrito Federal, 23.VI.2008

Solanaceae

Cestrum intermedium Sendtn. (night blooming jasmine), Recanto Novo, Minas Gerais, 13.V.2006

Cestrum intermedium, *O. basilicum*, and *M. rubra* are new hosts for *T. neocaledonicus* in the world. However, the presence of *T. neocaledonicus* on *Cestrum* sp. has been registered in the state of Rio de Janeiro (Flechtman 1981). Other plant species belonging to the *Ocimum* and *Morus* genera (*O. sanctum* L., *M. alba* L., *M. australis* Poir, *M. indica* L., *M. nigra* L., *M. sp.*) were registered as hosts for *T. neocaledonicus* in Cuba, India, Japan, and Thailand (Livshits and Salinas-Croche 1968; Baker 1975; Gupta 1992; Gupta and Gupta 1994; Ehara and Yamaguchi 2001).

Tetranychus urticae

Koch 1836

Tetranychus urticae Koch, 1836. (Koch 1836 *apud* Pritchard and Baker 1955). Type-host: *Urtica* sp. Type-locality: Regensburg, Germany.

Asteraceae

Ambrosia polystachya DC. (ragweed), Itaquí, Rio Grande do Sul, 29.IX.2007.

Parthenium sp. (parthenium), Universidade Federal de Santa Maria, Rio Grande do Sul, 26.IX.2007.

Paschoal (1970b) first confirmed the presence of *T. urticae* on *A. trifida* L. in the state of São Paulo. *Ambrosia polystachya* is a new host for *T. urticae* in Brazil and in the world.

Reports of *Tetranychus urticae* on *Parthenium* sp. were made in India (Gupta and Gupta, 1994). In the Americas, the first report of its presence on *Parthenium* sp. occurred in Rio Grande do Sul.

Bignoniaceae

Pyrostegia venusta Miers (flame vine), Universidade Federal de Lavras, Lavras, Minas Gerais, 09.V.2006.

Vargas et al. (1996) described *T. urticae* infestations on *P. ignea* C. Presl. in Costa Rica. This was the first report of *T. urticae* infesting *P. venusta* in the world.

Fabaceae

Phaseolus lunatus L. (lima beans), Embrapa Hortaliças, Gama, Distrito Federal, 07.XII.2005.

Macroptilium atropurpureum (L.) (siratro), Caliman farm, Cruz das Almas, Bahia.

Infestations of *T. urticae* on *P. lunatus* were reported in the United States, India, and Thailand (Reeves 1963; Gupta and Gupta 1994; Ho et al. 1997) and on *M. atropurpureum* in Greece (Hatzinikolis 1969). *Phaseolus lunatus* and *M. atropurpureum* are new hosts for *T. urticae* in South America.

Passifloraceae

Passiflora edulis Sims (yellow passion fruit), Embrapa Roraima, Boa Vista, Roraima, 17.V.2007.

Tetranychus urticae on passion fruit (*P. caerulea* L. and *P. edulis*) was reported in Venezuela (Doreste 1968) and Australia (*Passiflora* sp.) (Gutierrez and Schicha 1983). This is the first report of *T. urticae* on passion fruit (*P. edulis*) in Brazil.

Poaceae

Triticum aestivum L. (wheat), Embrapa Trigo, Passo Fundo, Rio Grande do Sul, 27.IX.2006.

The first *T. urticae* infestation of wheat crops (*Triticum* sp.) was registered in Greece (Hatzinikolis 1969). This wheat infestation in Brazil is the first occurrence in the Americas.

Zingiberaceae

Alpinia purpurata K. Schum (alpinia), Faculdade da Terra de Brasília, Recanto das Emas, Distrito Federal, 28.XI.2005.

Cases of *Tetranychus urticae* on Zingiberaceae (*Curcuma longa* L.) were reported in India (Gupta and Gupta 1994). This is the first report of *T. urticae* on alpinia (*A. purpurata*) in the world.

The single sample collected in Uruguay revealed a new host for the two-spotted spider mite (*T. urticae*).

Verbenaceae

Lantana camara L. (wild sage), Rio Negro, San Javier, Uruguay, 06.VI.2007. (-58.1312 S; -32.6368 W – collected by D. Navia and E. Castiglione).

Tetranychus urticae infesting *L. camara* was first observed in the United States and in India

(Thewke and Enns 1969; Gupta and Gupta 1994); this is the first record in South America.

New localities for Tetranychidae mites in Brazil

New localities have been registered for *Eotetranychus tremae*, *O. anonae*, *T. gloveri*, *P. ulmi*, and *E. smithi* (Table 2).

Tetranychinae Berlese, Tetranychini Reck

Eotetranychus tremae

De Leon 1957

Eotetranychus tremae, De Leon 1957. Type-host: *Trema floridanum*. Type-locality: Coral Gables, United States.

Reports of *Eotetranychus tremae* infesting *Acalypha* sp. ornamental plant have been registered in Viçosa, Minas Gerais. Previous reports in Brazil were registered in the states of Rio de Janeiro (Flechtmann 1981) and São Paulo (Flechtmann 2004; Daud and Feres 2005; Feres et. al. 2005).

Oligonychus anonae

Paschoal 1970

Oligonychus anonae Paschoal, 1970. Type-host: *Annona muricata* L., *Persea americana* Mill., *Vitis vinifera* L. Type-locality: Brazil.

Paschoal (1970a, 1970b) described this species from samples collected from soursop (*Annona. muricata*) in the state of São Paulo. Its presence on *A. squamosa* in the municipality of Mocimbinho, in northern Minas Gerais, indicates a new locality for this species in Brazil.

Tetranychus gloveri

Banks 1900

Tetranychus gloveri Banks, 1900. Type-host: *Gossypium hirsutum* L. Type-locality: Baton Rouge, United States

Tetranychus gloveri was reported on beans in Bahia (Bondar 1930) and on papaya at an (unspecified location) (Migeon and Dorkeld 2006). Reports of its presence on papaya (*Carica papaya* L.) in the state of Rio Grande do Norte indicate a new locality for this species.

Panonychus ulmi
(Koch 1836)

Tetranychus ulmi Koch, 1836. Koch (1836). Type-host: *Ulmus* sp. Type-locality: Regensburg, Germany.

Reports of the European red mite (*P. ulmi*) on grape vines (*Vitis vinifera* L.) were registered for the first time in Brazil in 2008 when damages caused by these mites were first noticed on crops in Rio Grande do Sul (Ferla and Botton 2008). In this work, a second occurrence of *P. ulmi* infesting grape vines was reported in Pirapora, state of Minas Gerais.

Panonychus ulmi was first reported in Brazil by Flechtmann (1967a), who discovered them on apples imported from Argentina. Soon after, Bleicher (1974) reported their presence on apple trees (*Malus domestica* L.) growing in orchards in the south of the country. However, European countries such as France, Austria, Portugal, Italy, and Greece consider the *P. ulmi* a grape vine pest (Rambier 1958; Artofer 1976; Carmona and Dias 1980; Girolami and Mozzi 1983; Papaioannou-Souliotis et al. 1994). Similar reports were

also registered in Morocco and the United States (Reeves 1963; Tixier et al. 2003).

Despite reports of *P. ulmi* on apple trees as early as 1967 (Flechtmann, 1976a), grape vines in Brazil were not affected, in contrast to the damages the *P. ulmi* caused to vineyards in Europe. Navia et al. (1998) hypothesized that the mites that attacked grape vines in Europe had different biotypes than those that infested apple trees. Moraes and Flechtmann (2008) emphasized the importance of avoiding introduction of European mites into Brazil, stating the reason this species was not present on grape vines in Brazil was unknown. They suggested the possibility of biological differences between Brazilian and European mites.

The spread of *P. ulmi* infestation on grape vines from Rio Grande do Sul (2005/2006) to Minas Gerais (2006) state could be the result of the two states trading plant propagation material. The hypothesis that European mites were introduced into Brazil cannot be discarded considering that *P. ulmi* were discovered approximately 40 years ago and have only recently been causing damage to grape vines. Ferla and Botton (2008) state that the probable causes for *P. ulmi* dispersion in Brazil are: the proximity of apple tree plantations to vineyards in the south of Brazil; the trading of infested plant material within Brazil and abroad; and unbalanced apple orchards due to excessive application of chemical products, especially non-selective fungicides.

New occurrences of phytophagous mites in South America were recently reported in the literature. For instance, the citrus Hindu mite, *Schizotetranychus hindustanicus* (Hirst) was collected from citrus in the municipality of Boa Vista, State of Roraima (Navia and

Marsaro Jr. 2010), and *Eotetranychus smithi* Pritchard & Baker 1955 from roses in Rio Branco, Acre (Mendonça et al. 2010); both states from the Northern region of the country. *Tetranychus roseus* Gutierrez 1969 was first observed in São Paulo in 2007 (Matioli et al. 2008) and its dispersal in the state was reported by Matioli et al (2010).

These new occurrences for mites in Brazil together with the results obtained in this study indicate the importance of intensified surveys on different host plants in the country. Doing this will broaden the understanding of the Tetranychidae family and give light to management and control of these mite species in agricultural systems.

Conclusions

Thirty-five new hosts were registered for 11 tetranychid species already known in Brazil: *E. banksi* (6 hosts), *M. tanajoa* (2), *O. anonae* (1), *O. mangiferus* (3), *T. bastosi* (1), *T. desertorum* (4), *T. evansi* (1), *T. ludeni* (4), *T. mexicanus* (2), *T. neocaledonicus* (3), and *T. urticae* (8), as well as one in Uruguay for *T. urticae*. New localities were registered in Brazil for *E. tremae*, *O. anonae*, *T. gloveri* and *P. ulmi*.

Acknowledgements

The authors would like to thank Fábio Aquino Albuquerque, Embrapa - Centro Nacional de Pesquisa de Algodão, Campina Grande, PA; Adalton Raga and Mario E. Sato, Instituto Biológico de São Paulo, Campinas, SP; Marineide Rosa Vieira, Faculdade de Ciências Agrônomicas, UNESP, Botucatu, SP; Lino Bittencourt Monteiro, Universidade Federal do Paraná, Paraná, RS; Manoel G. C. Gondin, Universidade Federal Rural de Pernambuco, Recife, PE; José Maria Milanez, Embrapa

Trigo, Passo Fundo, RS; Empresa Caliman, Espírito Santo; Aloyséia Cristina Noronha, Embrapa Embrapa Amazônia Oriental, Belém, PA; Marcílio Thomazini, Embrapa Acre, Rio Branco, Acre; Marcione C. M. Chagas, Empresa de Pesquisa Agropecuária do Rio Grande do Norte, Natal, RN; Elton Lúcio de Araújo, Universidade Federal Rural do Semi-Árido, Mossoró, RN; Ricardo Adaime da Silva, Embrapa Agroflorestral, Macapá, AP; Paulo E. Degrande, Universidade Federal da Grande Dourados, Dourados, MS; Marcelo Poletti, PROMIP Ltda, Piracicaba, SP, and students Renato Tratch and Samuel Rogger for assistance in the collection of mite samples. Research for this paper was conducted in part to fulfill requirements for the PhD degree of the senior author in Animal Biology, University of Brasília, and Ecology, Biodiversity and Evolution, University of Montpellier II, France. This work was partially supported by CAPES, Fundação Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Brazil. The authors are also grateful to CNPq, Conselho Nacional de Desenvolvimento Científico e Tecnológico, Brazil, for providing the fellowship.

Editor's note: Paper copies of this article will be deposited in the following libraries. Universitätsbibliothek Johann Christian Senckenberg, Frankfurt Germany; National Museum of Natural History, Paris, France; Field Museum of Natural History, Chicago, Illinois USA; University of Wisconsin, Madison, USA; University of Arizona, Tucson, Arizona USA, Smithsonian Institution Libraries, Washington D.C. USA; The Linnean Society, London, England. The date of publication is given in 'About the Journal' on the JIS website.

References

- Andre M. 1933. Note sur un tétranyque nuisible au cotonnier en Nouvelle-Calédonie. *Bulletin de la Muséum National d'Histoire Naturelle* (2) 5(4): 302-308.
- Andrews KL, Poe SL. 1980. Spider mites of El Salvador, Central America (Acari: Tetranychidae). *Florida Entomologist* 63(4): 502-505.
- Artofer R. 1976. Spider-mites in vine cultivation. *Pflanzenarzt* 29(7): 73.
- Baker EW, Pritchard AE. 1960. The tetranychoid mites of Africa. *Hilgardia* 29(11): 455-574.
- Baker EW, Pritchard AE. 1962. Arañas rojas de América Central (Acarina:Tetranychidae). *Revista de la Sociedad Mexicana de Historia Natural* 23: 309-340.
- Baker EW, Tuttle DM. 1994. *A guide to the spider mites (Tetranychidae) of the United States*. Indira Publishing House.
- Baker EW. 1975. Plant-feeding mites of Thailand (Tetranychidae, Tenuipalpidae, and Tuckerellidae). *Plant Protection Service Technical Bulletin* 35: 43.
- Banks N. 1900. The red spiders of the United States (Tetranychus and Stigmaeus). United States Department of Agriculture, Division of Entomology. *Technical Series Bulletin* 8: 65-77.
- Bleicher E. 1974. Ocorrência do acaro *Panonychus ulmi* (Koch, 1836) Tuttle & Baker (1966) no Estado de Santa Catarina. *Revista o Solo* 66(1): 64-64.
- Bolland HR, Guterrez J, Flechtmann CHW. 1998. *World catalogue of the spider mite family (Acari: Tetranychidae), with references to taxonomy, synonymy, host plants and distribution*. Brill Academic Publishers.
- Bondar G. 1930. Insetos daninhos e moléstias dos feijões na Bahia. *Boletim do Laboratorio de Pathologia Vegetal* 9: 1-83.
- Bondar G. 1938. Notas entomológicas de Bahia III. *Revista de Entomologia, Brasil* 9(3-4): 441-445.
- Carmona MM, Dias JCS. 1980. O complexo Acarina nas culturas portuguesas. I. *Congresso Português de Fitiatria e Fitofarmacologia*. pp. 97-115.
- Daud RD, Feres RJF. 2005. Diversidade e flutuação populacional de ácaros (Acari) em *Mabea fistulifera* Mart. (Euphorbiaceae) de dois fragmentos de mata estacional semidecídua em São José do Rio Preto, SP. *Neotropical Entomolog* 34(2): 191-201.
- Davis JJ. 1968. Studies of Queensland Tetranychidae. 3. Records of the genus *Tetranychus*. *Queensland Journal of Agricultural and Animal Sciences* 25: 27-67.
- De Leon D. 1957. Two new *Eotetranychus* and a new *Oligonychus* from Southern Florida (Acarina: Tetranychidae). *Florida Entomologist* 40(3): 111-113.
- Doreste E. 1968. Primera lista de acaros de importancia agricola en Venezuela. *Agronomia Tropical* 18(4): 449-460.
- Edge VE, James DG. 1986. Organo-tin resistance in two-spotted spider mite, *Tetranychus urticae* Koch (Acarina:

Tetranychidae) in Australia. *Journal of Economic Entomology* 79(6): 1477-1483.

Ehara S, Yamaguchi K. 2001. Discovery of *Tetranychus neocaledonicus* André (Acari: Tetranychidae) from Amami-Oshima Island, Japan. *Plant Protection* 55: 268-272.

Ehara S. 1956. Notes on some tetranychid mites of Japan. *Japanese Journal of Applied Zoology* 21(4): 139-147.

Ehara S. 1999. Revision of the Spider Mite Family Tetranychidae of Japan (Acari, Prostigmata). *Species Divers* 4: 63-141.

Feres RJF, Flechtmann CHW. 1986. Nova espécie de *Oligonychus* Berlese (Acari, Tetranychidae) da região de Botucatu, SP. *Revista Brasileira de Entomologia* 30(1): 163-166.

Feres RJF, Flechtmann CHW. 1988. *Aponychus chiavegato* n.sp. collected From *Citrus* sp. (Rutaceae) in São Paulo, Brazil (Acari, Tetranychidae, Eurytetranychini). *Revista Brasileira de Biologia* 48(4): 959-964.

Feres RJF, Flechtmann CHW. 1995. Mites (Acari) associated with bamboo (*Bambusa* sp., *Poaceae*) in a woody area from Northwestern São Paulo State, Brazil. *Revista Brasileira de Zoologia* 12(3): 533-546.

Feres RJF, Flechtmann CHW. 1995a. A new *Oligonychus* and description of the female allotype of *O. psidium* Estebanes & Baker (Acari: Tetranychidae) from *Qualea grandiflora* Mart. (Vochysiaceae) in northwestern São Paulo, Brazil. *Revista Brasileira de Zoologia* 12(3): 529-532.

Feres RJF, Flechtmann CHW. 1995b. *Sonotetranychus angiopenis* n.sp. (Acari:

Tetranychidae) from *Bauhinia* sp. (Fabaceae) in northwestern São Paulo State, Brazil. *International Journal of Acarology* 21(2): 89-91.

Feres RJF, Flechtmann CHW. 1986a. Duas novas espécies de *Tetranychus* Dufour (Acari, Tetranychidae) dos Estados de São Paulo e Rio de Janeiro. *Revista Brasileira de Biologia* 46(2): 371-376.

Feres RJF. 1986. Dois Novos *Eotetranychus* (Acari, Tetranychidae) do Estado de São Paulo. *Revista Brasileira de Biologia* 46(2): 365-369.

Feres RJF. 1992. *Allonychus brevipenis*, a new species from *Triplaris surinamensis* Cham. in Brazil with a new host records for *A. brasiliensis* (McGregor) and *A. reisi* Paschoal (Acari: Tetranychidae). *International Journal of Acarology* 18(4): 299-301.

Feres RJF, Lofego AC, Oliveira AR. 2005. Ácaros plantícolas (Acari) da "Estação Ecológica do Noroeste Paulista", Estado de São Paulo, Brasil. *Biota Neotropica* 5(1): 1-14.

Ferla NJ, Botton M. 2008. Ocorrência do ácaro vermelho europeu associado à cultura da videira no Rio Grande do Sul, Brasil. *Ciência Rural* 38: 25-35.

Ferragut F, Escudero LA. 1999. *Tetranychus evansi* Baker & Pritchard (Acari, Tetranychidae), una nueva araña roja en los cultivos hortícolas españoles. *Boletín de Sanidad Vegetal, Plagas* 25: 157-164.

Flechtmann CHW. 1967. *Contribuição para o conhecimento dos ácaros de plantas de algumas regiões do estado de São Paulo*. Tese

(Doutorado). Escola Superior de Agricultura Luiz de Queiroz, Universidade de São Paulo.

Flechtmann CHW. 1967a. Ácaros encontrados sobre maçãs de procedência Argentina. *Anais da Escola Superior de Agricultura Luiz de Queiroz* 24: 83-85.

Flechtmann CHW. 1967b. New Tetranychidae (Acarina) from Brazil. *Anais Escola Superior de Agricultura Luiz de Queiroz* 24: 323-331.

Flechtmann CHW. 1972. *Tetranychus (T.) urticae* Koch. 1836 (Acari) praga severa de orquídeas (*Catasetum* sp.) em ripados. *Revista de Agricultura* 47(2): 70.

Flechtmann CHW. 1976. A report on the Tetranychidae (Acari) of Brazil: an emmendation. *Revista Brasileira de Entomologia* 20(2): 115-116.

Flechtmann CHW. 1981. A small collection of spider mites from Jaguanum Island, Rio de Janeiro, Brazil, with description of *Aponychus insularis* (Prostigmata: Tetranychidae). *International Journal of Acarology* 7: 87-90.

Flechtmann CHW. 2004. Two new plant feeding mites from *Brachiaria ruziziensis* in citrus groves in São Paulo, Brazil and new distribution records of other plant mites in Brazil. *Zootaxa* 708: 1-11.

Flechtmann CHW, Baker EW. 1970. A Preliminary report on the Tetranychidae (Acarina) of Brazil. *Annals of the Entomological Society of America* 63(1): 156-163.

Flechtmann CHW, Baker EW. 1975. A report on the Tetranychidae (Acari) of Brazil. *Revista Brasileira de Entomologia* 19(3): 111-122.

Flechtmann CHW, Bastos JAM. 1972. Ácaros Tetranychidae do Estado do Ceará, Brasil. *Ciência Agronômica* 2(2): 83-90.

Foott WH. 1962. Competition between two species of mites. I. Experimental results. *Canadian Entomologist* 94: 365-375.

Foott WH. 1963. Competition between two species of mites. II. Factors influencing intensity. *Canadian Entomologist* 95: 45-57.

Furtado IP, Moraes de GJ, Kreiter S, Knapp M. 2006. Search for effective natural enemies of *Tetranychus evansi* in south and southeast Brazil. *Experimental and Applied Acarology* 40(3-4): 157-174.

Garrett LE, Haramoto FH. 1967. A catalog of Hawaiian Acarina. *Proceedings of the Hawaiian Entomological Society* 19(3): 381-414.

Girolami V, Mozzi A. 1983. Distribution, economic threshold and sampling methods of *Panonychus ulmi* (Koch). In: Cavalloro R, Editor. *Statistical and mathematical methods in population dynamics and pest control*. Proceedings of EC Experts' Group. pp. 90-101. AA Balkema

Gupta SK, Gupta YN. 1994. A taxonomic review of Indian Tetranychidae (Acari: Prostigmata) with description of new species, redescription of known species and keys to genera and species. *Memoirs of the Zoological Survey of India* 18(1): 1-196.

Gupta SK. 1976. Contribution to our knowledge of tetranychid mites (Acarina) with descriptions of three new species from India. *Oriental Insects* 10(3): 327-351.

- Gupta SK. 1992. Arachnida: plant mites (Acari). In: Ghosh AK, Editor. *Fauna of West Bengal* 3: 61-211. Zoological Survey of India.
- Gutierrez J, Schicha E. 1983. The spider mite family Tetranychidae (Acari) in New South Wales. *International Journal of Acarology* 9(3): 99-116.
- Gutierrez J. 1987. The cassava green mite in Africa: one or two species? (Acari: Tetranychidae). *Experimental and Applied Acarology* 3: 163-168.
- Hatzinikolis EN. 1969. Preliminary notes on Tetranychoid and Eriophyid mites infesting cultivated plants in Greece. *Proceedings of the 2nd International Congress of Acarology* 161-167.
- Helle W, Pijnacker LP. 1985. Spider mites: their biology, natural enemies and control. In: Helle W, Sabelis MW, Editors. *World Crop Pests* 1A, 1B. Elsevier Science Publishers.
- Ho CC, Lo KC, Chen WH. 1997. Spider mite (Acari: Tetranychidae) on various crops in Taiwan. *Journal of Agricultural Research of China* 46(4): 333-346.
- Jeppson LR, Keifer HH, Baker EW. 1975. *Mites injurious to economic plants*. University of California Press.
- Koch KL. 1836. Deutschlands Crustaceen Myriapoden und Arachniden. 1835-1844. Pustet, Regensburg.
- Livshits IZ, Salinas-Croche A. 1968. *Preliminares acerca de los acaros "Tetranychos" de Cuba*. Centro Nacional Fitosanid.
- Matioli AL, Gomes JAV, Raga A, Sato ME. 2008. First record of *Tetranychus roseus* Gutierrez (1969) (Acari: Tetranychidae) in Brazil. In: Barros-Battest DM, Fadini MAM, Perez CA, Onofrio VC, Famadas KM, Pallini A, Venzon M, Moraes de GJ, Editors. *Anais do II Simpósio Brasileiro de Acarologia*. pp. 172.
- Matioli AL, Sato ME, Mineiro JLC. 2010. New occurrence of *Tetranychus roseus* Gutierrez, 1969 (Acari: Tetranychidae) in Brazil. In: Moraes de GJ, Castilho RC, Flechtmann CHW, Editors. 13th International Congress of Acarology Abstract Book. pp. 148.
- McGregor EA. 1914. Four new tetranychids. *Annals of the Entomological Society of America* 7: 354-364.
- McGregor EA. 1950. Mites of the family Tetranychidae. *American Midland Naturalist* 44(2): 257-420.
- Mendonça RS, Navia D, Flechtmann CHW. 2010. Two new spider mites (Acari: Tetranychidae) from Brazil: a *Monoceronychus* McGregor (Bryobiinae) from fingergrass and an *Oligonychus* Berlese (Tetranychinae) from grape. *International Journal of Acarology* 36(6): 487-502.
- Meyer MKPS, Ryke PAJ. 1959. A revision of the spider mites (Acarina: Tetranychidae) of South Africa, with descriptions of a new genus and new species. *Journal of the Entomological Society of Southern Africa* 22(2): 330-366.
- Meyer MKPS. 1974. A revision of the Tetranychidae of Africa (Acari) with a key to the genera of the world. *Entomology Memoir* 36: 1-291.

Meyer MKPS. 1987. African Tetranychidae (Acari: Prostigmata) - with reference to the world genera. *Entomology Memoir* 69: 1-175.

Migeon A, Dorkeld, F. 2006. Spider Mites Web: a comprehensive database for the Tetranychidae. Available online, <http://montpellier.inra.fr/CBGP/spmweb>

Miranda, LM, Navia D, Rodrigues JCV. 2007. *Brevipalpus* mites Donnadieu (Prostigmata: Tenuipalpidae) associated with ornamental plants in Distrito Federal, Brazil. *Neotropical Entomology* 36(4): 587-592.

Moraes de GJ, Flechtmann CHW. 2008. *Manual de acarologia – Acarologia básica e ácaros de plantas cultivadas no Brasil*. Holos Editora.

Moraes de GJ, Moreira NA, Delalibera Jr I. 1995. Growth of the mite *Mononychellus tanajoa* (Acari: Tetranychidae) on alternative plant hosts in northeastern Brazil. *Florida Entomologist* 78(2): 350-354.

Navia D, Flechtmann CHW, Moraes de GJ. 1998. Avaliação do risco de introdução de ácaros fitófagos associados à cultura da uva no Brasil. *Embrapa Recursos Genéticos e Biotecnologia Documentos* 32: 1-51.

Navia D, Moraes de GJ, Flechtmann CHW. 2007. Phytophagous mites as invasive alien species: quarantine procedures. In: Morales-Malacara JB, Behan-Pelletier V, Ueckermann E, Pérez TM, Estrada E, Gispert C, Badii M, Editors. *Proceedings of the XI International Congress of Acarology*. pp. 307-316.

Navia D, Marsaro Jr AL. 2010. First report of the citrus Hindu mite, *Schizotetranychus hindustanicus* (Hirst) (Prostigmata:

Tetranychidae), in Brazil. *Neotropical Entomology* 39(1): 140-143.

Papaionnaou-Souliotis P, Ragusa Di Chiara S, Tsolakis H. 1994. Phytophagous mites and their predators on cultivated plants in Greece during 1975-1990. *Annals of the Benaki Phytopathological Institute* 17: 35-87.

Paschoal AD. 1970a. New Brazilian spider mites (Acarina: Tetranychidae). *Anais da Escola Superior de Agricultura Luiz de Queiroz* 27: 139-455.

Paschoal AD. 1970b. *Contribuição ao conhecimento da família Tetranychidae no Brasil (Arachnida: Acarina)*. Tese (Doutorado). *Escola Superior de Agricultura Luiz de Queiroz*.

Pritchard AE, Baker EW. 1955. A revision of the spider mite family Tetranychidae. *Memoirs Series, Pacific Coast Entomological Society* 2: 472.

Rahman KA, Sapra AN. 1940. Mites of the family Tetranychidae from Lyallpur with descriptions of four new species. *Proceedings of the Indian Academy of Science, Ser. B* 11: 17-196.

Rambie A. 1958. Les tétranyques nuisibles à la vigne en France continentale. *Revue de Zoologie Agricole et Appliquée* 1-3: 1-21.

Reeves RM. 1963. Tetranychidae infesting woody plants in New York State, and a life history study of the elm mite *Eotetranychus matthyssei* n. sp. *Cornell University Agricultural Experimental Station Memoir* 380: 99.

Zacher F. 1933. Übersicht der deutschen Spinnmilben. *Mitteilungen aus dem Zoologischen Museum in Berlin* 19: 584-589.

Larger appendix available online.