

Abundance of Apple Maggot, *Rhagoletis pomonella*, Across Different Areas in Central Washington, with Special Reference to Black-Fruited Hawthorns

Authors: Yee, Wee L., Klaus, Michael W., Cha, Dong H., Linn, Charles E., Goughnour, Robert B., et al.

Source: Journal of Insect Science, 12(124) : 1-14

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.012.12401>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Abundance of apple maggot, *Rhagoletis pomonella*, across different areas in central Washington, with special reference to black-fruited hawthorns

Wee L. Yee^{1a*}, Michael W. Klaus^{2b}, Dong H. Cha^{1,6c}, Charles E. Linn, Jr.^{3d}, Robert B. Goughnour^{4e} and Jeffrey L. Feder⁵

¹USDA-ARS, Yakima Agricultural Research Laboratory, 5230 Konnowac Pass Road, Wapato, WA 98951

²Washington State Department of Agriculture, 21 North 1st Avenue, Suite 103, Yakima, WA 98902

³Department of Entomology, Cornell University, 630 W North Street, Geneva, NY 14456

⁴Washington State University, Vancouver, Research and Extension Unit, 1919 NE 78th Street, Vancouver, WA 98665

⁵Department of Biological Sciences, 100 Galvin Life Sciences Center, University of Notre Dame, Notre Dame, IN 46556

⁶Current address: USDA-ARS, Yakima Agricultural Research Laboratory, 5230 Konnowac Pass Road, Wapato, WA 98951

Abstract

The apple maggot fly, *Rhagoletis pomonella* (Walsh) (Diptera: Tephritidae), infests non-commercial apple (*Malus domestica* (Borkh.) Borkh.) and native black-fruited hawthorns (mostly *Crataegus douglasii* Lindl.) in central Washington, but little has been published on the abundance of the fly in this region. In this paper, the abundance of *R. pomonella* across different sites near apple-growing areas in central Washington is documented in order to assess the threat of the fly to commercial apple orchards. The fly was first detected on traps in Klickitat, Yakima, and Kittitas Counties in 1981, 1995, and 1997, respectively. From 1981–2010 in Kittitas and Yakima Counties, only 0 to 4.7% of traps on apple, crabapple, and hawthorn trees were positive for flies, whereas in Klickitat County, located farther from commercial apple orchards, 0 to 41.9% of traps were positive. In 2008, in Yakima County and Goldendale in Klickitat County, 7.8% of black-fruited hawthorn trees were infested, with 0 to 0.00054 larvae per fruit. In 2010, in Kittitas and Yakima Counties and Goldendale in Klickitat County, 25.0% of *C. douglasii* trees were infested, with 0.00042 to 0.00248 larvae per fruit. In 2010, in a remote forested area of Klickitat County far from commercial apple orchards, 94.7% of *C. douglasii* trees were infested, with 0.20813 larvae per fruit. Overall results suggest *R. pomonella* is unlikely to develop high populations rapidly near major commercial apple-growing areas in central Washington, including in black-fruited hawthorns, increasing chances it can be kept out of commercial orchards.

Keywords: *Crataegus douglasii*, Diptera, larval infestation levels, Tephritidae, trapping

Correspondence: ^a wee.yee@ars.usda.gov, ^b mklauus@agr.wa.gov, ^c dong.cha@ars.usda.gov, ^d cell1@cornell.edu, ^e grblakee@hotmail.com, ^f jeffrey.l.feder.2@nd.edu, *Corresponding author

Editor: Henry Hagedorn was editor of this paper.

Received: 3 June 2011, **Accepted:** 11 February 2012

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 12, Number 124

Cite this paper as:

Yee WL, Klaus MW, Cha DH, Linn Jr CE, Goughnour RB, Feder JL. 2012. Abundance of apple maggot, *Rhagoletis pomonella*, across different areas in central Washington, with special reference to black-fruited hawthorns. *Journal of Insect Science* 12:124. Available online: <http://www.insectscience.org/12.124>

Introduction

The apple maggot fly, *Rhagoletis pomonella* (Walsh) (Diptera: Tephritidae), is a quarantine pest of apples (*Malus domestica* (Borkh.) Borkh.) in the Pacific Northwest of the U.S. The species evolved on hawthorns (*Crataegus* spp.) in eastern North America (Bush 1966; Bush and Smith 1998) and Mexico (Rull et al. 2006) and moved from hawthorn onto apple about 150 years ago in the eastern U.S.A. (Walsh 1867), differentiating over time into hawthorn and apple host races (Feder et al. 1988; Bush and Smith 1998). The fly apparently was introduced from eastern North America into the West Coast of the U.S.A., perhaps as early as 1951 into Oregon (AliNiazee and Wescott 1986). The first definitive detection of the fly in the West Coast was of larvae infesting backyard apples in 1979 in Portland, Oregon (AliNiazee and Penrose 1981). In 1980, the fly was detected in nearby Vancouver, Washington, and in 1982, in Spokane, Washington; in 1983–1984 it was detected in California and in Utah (Brunner 1987). In western Washington, the fly infests apples and native and introduced hawthorns (Tracewski et al. 1987). Two native black-fruited hawthorns, black hawthorn, *C. douglasii* var. *douglasii* Lindl., and Suksdorf's hawthorn, *C. suksdorfii* (Sarg.) Kruschke, are infested by *R. pomonella* in Washington (Yee and Goughnour 2008).

Central Washington is the center of the largest apple industry in the U.S.A., producing most of the apples in the state of Washington, which were worth \$1.4 billion in 2009 (USDA 2010). There is zero tolerance for *R. pomonella*-infested apples (WSDA 2001), making the mere presence of the fly near an orchard a potential threat. However, little work has been published on the abundance of *R. pomonella* in central Washington. As a result, there is a lack of information about the general status of the fly as a threat to commercial apples in the region, information that could be of great value for management efforts in Washington and for allaying concerns of regulatory officials in apple export markets. In central Washington, apples appear less utilized than black-fruited hawthorns (Yee 2008), raising the possibility that hawthorns rather than feral apples in some sites could harbor high fly populations that can threaten apple orchards, but few data are available on fly abundance in these hawthorns.

In this paper, the abundance of *R. pomonella* across different sites near apple-growing areas in central Washington is documented in order to assess the threat of the fly to commercial apple orchards. Fly abundance was determined by fly catches during a long-term detection survey in apples, crabapples, and hawthorns, and by larval infestation levels in *C. douglasii* and *C. suksdorfii*. The

implications of the findings for pest management of flies in central Washington are discussed.

Materials and Methods

Adult trapping survey in central Washington (1981–2010)

Washington State Department of Agriculture's (WSDA) annual statewide apple maggot detection survey began in 1981, shortly after *R. pomonella* was detected in Portland (Brunner 1987). The goal of the survey since its inception was to delimit the spread of the fly and keep it out of commercial apple orchards. The survey work reported here was conducted in Kittitas, Yakima, and Klickitat Counties (Figure 1) from 1981–2010, essentially spanning the period since *R. pomonella* was first detected in apple in the Portland area until the time of this paper. Most areas in Kittitas, Yakima, and Klickitat Counties (Figure 1) are relatively dry ponderosa pine or bunchgrass and sagebrush habitats (Lyons and Merilees 1995). The ponderosa pine habitats in Kittitas and Klickitat Counties are wetter than the bunchgrass and sagebrush habitat in Yakima County. Major commercial apple orchards are found throughout the Yakima Valley as well as the Naches Valley (gray areas in Figure 1). The survey emphasized trapping in these areas.

Flies were captured over the course of the 30-year monitoring period using yellow sticky traps (Pherocon® AM (apple maggot) traps (Zoecon Co., <http://zoecon.com/>) or equivalent) baited with one ammonium carbonate or bicarbonate lure containing 10 g of the chemical attractant. Traps were hung in non-commercial apple trees, crabapple, and ornamental hawthorns from mid-June to late September in backyards, roadsides, and rural

settings, including wilderness areas. Native hawthorns were trapped less frequently due to a scarcity of these plants in trapped areas. Each tree had one trap that was replaced every 2–4 weeks, as the trap became covered with debris. The survey traps were categorized as three types: (1) general survey traps were placed on host trees at a density of roughly one per 2.6 km² (1 mile²) in places where no flies had previously been detected; (2) high-density traps were placed in all accessible host trees 0.8 to 1.6 km (½ to 1 mile) around positive fly trees, and were used to delimit the scope of newly detected fly infestations; (3) certification traps were placed in the 1.9 km (1.2 mile) buffer zone around apple orchards, and were deployed in Kittitas and Yakima Counties (many large commercial orchards present) but not in Klickitat County (few commercial orchards present). Certification trapping is required for the exportation of apples from quarantined areas and may be required for exporting apples from fly-free zones. Traps were deployed in pest-free and quarantine areas. Pest-free areas are those where the larval stage of *R. pomonella* have not been found; quarantine areas are those where adults and larvae have been found. Figure 2 shows the typical distribution of the three trap types deployed in Kittitas and Yakima Counties during the trapping survey (in 2009 in this case). The distribution approximately mirrored that of host-trees present near apple-growing areas in the Yakima Valley; areas immediately outside this distribution were generally non-irrigated sagebrush habitat with few or no trees. Klickitat County had not been trapped in some years. Otherwise, numbers of traps varied from 6 to 4,482 per county over the 30 years of the survey. Many of the same trees were trapped every year of the study. Flies were removed from traps every 1–3 weeks each season. As part of a management program,

trees that trapped positive for flies were generally treated weekly with applications of the insecticide Imidan (Gowan Co., <http://www.gowanco.com/>) by county pest control boards. *R. pomonella* were differentiated from *Rhagoletis zephyria* Snow, the snowberry maggot, based on Wescott (1982). Host-tree species were not always separated in the detection survey, and so trap records from different host trees were combined.

Larval abundance survey in hawthorns in central Washington (2008 and 2010)

In addition to the trapping studies of adult abundance, larval surveys were conducted at 10 field sites in 2008 and 2010 to confirm *R. pomonella* infestation in hawthorn fruit. Although unlikely, it is possible that flies captured on yellow traps could conceivably have originated not from hawthorn but elsewhere (e.g., from local apple sources), or that fly larvae feeding in hawthorns were not *R. pomonella* but *R. zephyria*. The larval abundance surveys allowed for these alternative scenarios to be rejected and for estimates of local fly densities to be made. The host-plant identified for *R. zephyria* in Washington is snowberry, *Symphoricarpos albus* (L.) Blake (Bush 1966; Yee 2008). Table 1 and Figure 1 show the 10 hawthorn fruit collection sites surveyed for larval abundance. These sites were located within the WSDA adult trapping survey areas, but no records were found showing native hawthorn trees used for collections had been trapped by WSDA, as emphasis had been placed on monitoring feral apples, crabapples, and ornamental hawthorns (Table 1, site trapping history in footnotes). *Crataegus douglasii* and/or *Crataegus suksdorfii* fruit were collected at the 10 sites in 2008 and 2010. Hawthorn identifications were based on Brunsfeld and Johnson (1990). In Ronald,

Roslyn, and Cle Elum, and in Wenas, Nile, Tampico, and Toppenish, only *C. douglasii* fruit were collected, although *C. suksdorfii* may have been present but not seen. In Klickitat sites, a survey of local hawthorn trees indicated that 86% (n = 29 total) were *C. douglasii*, with the remainder *C. suksdorfii*; in Goldendale, 75% were *C. douglasii* (n = 96 total). No native red-fruited hawthorns (Phipps 1998) were found at the 10 collection sites.

In 2008, randomly selected black-fruited hawthorn fruit from trees at Nile (19 trees), Wenas (40 trees), Tampico (21 trees), Toppenish (4 trees), and Goldendale (34 trees) sites were sampled from mid-August to late September. The number of trees sampled at the sites reflected a combination of how many trees were growing in the area that could be accessed and available labor for picking fruit. Fruit on trees among and within sites ripened asynchronously, but all fruit was collected when it was ripe and black. *Crataegus douglasii* and *C. suksdorfii* at the Goldendale site in 2008 were not distinguished and, thus, pooled results are presented for this site in 2008. However, all other samples were distinguished by host-tree species. Collected fruit were placed in 3.8-L zip lock plastic bags, transported back to the laboratory, and held in an ~25° C constant temperature room with a 16:8 L:D photoperiod. Each fruit sample was weighed when fresh. From one sample, 40 randomly selected fruits were weighed in order to estimate the number of fruit collected in the sample. Fruit were kept in bags for two to four weeks and then, if necessary, placed in tubs at ~20% relative humidity to prevent or slow rotting and molding. Bags and tubs were checked for puparia (used synonymously with larvae here) two or three times a week for six weeks. Puparia were held at 20–22° C for around

three weeks, stored in moist soil, chilled at 3–4° C for six months, and then held at 26–27° C and 16:8 L:D for adult emergence. Adult flies that emerged following laboratory rearing were identified (Wescott 1982). Voucher specimens were kept at the insect collection at the USDA-ARS Yakima Agricultural Research Laboratory in Wapato, WA.

In 2010, fruit from randomly selected *C. douglasii* trees at Ronald, Roslyn, and Cle Elum (19 trees from the 3 sites combined), Wenas (18 trees), Middle Klickitat and Lower Klickitat (19 trees), and Goldendale sites (15 trees) (Figure 1, Table 1) were sampled from mid-August to early October. In addition, *C. suksdorfii* trees at Middle and Lower Klickitat (5 trees) and Goldendale sites (9 trees) (Figure 1, Table 1) were sampled from August to September. Nile, Tampico, and Toppenish sites were dropped in 2010 due to labor limitations, as labor was diverted to Roslyn and Klickitat sites in 2010 because those trees had not been previously sampled, and we wanted to document a wider range of sites. Some trees sampled at Wenas and Goldendale had also been sampled in 2008. Fruit were maintained and puparia were handled as described above for the 2008 collections, except that puparia were held at 20–21° C for 10 weeks under an 8–12 hour day length for five days per week, and two days in darkness. These conditions resulted in the emergence of “non-diapausing” adults over the 10-week rearing period, allowing for rapid species confirmations of flies from samples.

Statistics

Data from the 1981–2010 adult trapping survey and from the host rearing studies were primarily descriptive and were not subjected to statistical analyses. For the fly detection survey, traps were deployed for two to four

weeks, but the exact numbers of days were not always recorded, preventing calculations of numbers of flies captured per trap per day. Instead, the measure used to standardize fly abundance across years was percentage of traps (one trap per tree) positive for at least one fly.

Results

Adult trapping survey in central Washington (1981–2010)

Among the three counties surveyed, *R. pomonella* was first caught in 1981 in Klickitat County on non-commercial apple, in 1995 in Yakima County at Union Gap (47.00, -120.55), and in 1997 in Kittitas County at Ellensburg (46.55, -120.48) (host affiliations for the latter two records are unclear) (Table 2). After these first recorded catches, one fly was caught in 1982 in the town of Klickitat (number of traps uncertain), about 3 km south of the Lower Klickitat site (Figure 1). In 1984, one fly was caught on 70 traps at Goldendale. In the same year, 12 flies were caught on 58 traps in the town of Klickitat. Over the 30 years in Kittitas and Yakima Counties, only 0–4.7% of the traps on apple, crabapple, and hawthorn trees were positive for flies, whereas in Klickitat County, south of Kittitas and Yakima Counties (Figure 1), 0–42.9% of traps were positive. In general, more flies were detected in Klickitat than in Kittitas County, and more in Kittitas than Yakima County over the 30 years when > 30 traps were deployed. The mean percentages of traps capturing flies (averaged from columns in Table 2) in Kittitas, Yakima, and Klickitat Counties were low: 1.5, 0.6, and 6.5%, respectively, starting when flies were first detected at these sites. In contrast, trap catches at western Washington sites on apple trees were high: 100% of yellow panels captured at least one fly during the same time period as traps in central

Washington were deployed in 2003–2005 (Yee et al. 2005). In Klickitat County from 2001–2006 (except 2005), traps were only placed in a few trees in the dry, extreme eastern portion of the county to verify the absence of flies there. As a result, fly catches during this period reflected location differences and trapping intensity rather than reductions in fly abundance in Klickitat County. Fluctuations in fly catches in Kittitas and Yakima Counties among years could be due to emphasis being placed in some years on trapping trees that were positive for flies in previous years (e.g., around Ellensburg in Kittitas County beginning in 2003 and Nile in Yakima County in 2004 and 2005; Table 2). Thus, in some years, overall fly abundance was probably lower than suggested by the survey.

Larval abundance survey in hawthorn in central Washington in 2008

In 2008, in Yakima County and Goldendale in Klickitat County, a total of 7.8% of black-fruited hawthorn trees were infested, with a range of infestation of 0–0.00054 larvae per fruit among sites (Table 3). Regardless of when and where hawthorns were sampled in central Washington in 2008, population densities were low. As a basis for comparison, average larval infestation levels of *C. douglasii* at one site in Skamania County in western Washington in 1982, 1983, and 1984 were 0.027, 0.060, and 0.015 per fruit, respectively (Tracewski et al. 1987), or 28–111 times greater than the highest infestation rate in 2008 in central Washington.

Larval abundance survey in hawthorns in central Washington in 2010

In 2010 in Kittitas and Yakima Counties and Goldendale in Klickitat County, a total of 25.0% of *C. douglasii* trees were infested, with a range of infestation of 0.00042–

0.00248 larvae per fruit among sites (Table 4). In 2010, in a remote forested area of Klickitat County (Middle and Lower Klickitat sites combined) relatively far from commercial apple orchards (Figure 1), 94.7% of *C. douglasii* trees were infested, with 0.20813 larvae per fruit (Table 4). This represents the one exception to the general observation that infestation levels in black-fruited hawthorns in central Washington are low.

Discussion

Over a period of 30 years, only 0–4.7% of traps on apple, crabapple, and hawthorn trees in Kittitas and Yakima Counties were positive for *R. pomonella*, implying that human enacted management measures and/or natural factors have prevented fly populations from reaching levels seen in western Washington (Tracewski et al. 1987; Yee et al. 2005; Yee and Goughnour 2008). The *R. pomonella* monitoring and targeted insecticide control program implemented by the WSDA and county pest control boards during this period could have been partly responsible for slowing the fly's spread. However, natural factors probably have played a larger role, because insecticides for fly control were and are not routinely used in the Roslyn area, Wenas, or Nile (sprayed only once in fall 2004), but these sites still have low abundance of *R. pomonella*. Some aspects of the physical environment are likely keeping these populations in check. One possibility is the climate, which is drier and hotter during the summer and fall than the climate west of the Cascade Mountains in Washington. *R. pomonella* also occurs in higher abundance in hawthorns in western New York (Reissig and Smith 1978), where it is relatively wet, than in central Washington. The low fly populations near or in apple-growing regions in central Washington is probably one reason why no

exported commercial apples from this region have been found to be infested with *R. pomonella* larvae (WSDA 2010), even though feral apples in central Washington are occasionally infested (Yee 2008).

Data from the 30-year detection survey suggest that *R. pomonella* was slow to spread and establish near or within the apple-growing regions in Kittitas and Yakima Counties. The trapping survey first detected the fly in Klickitat County 13 and 15 years before it was found in Kittitas and Yakima Counties, respectively. These data suggest that the fly may have spread relatively rapidly to Klickitat County following its introduction to the Pacific Northwest, but has been slower in expanding its range eastward from Klickitat, perhaps due in part to the difficulty of dispersal from forested to more barren sagebrush habitats with isolated trees and/or reduced opportunity for human mediated transport of infested fruit, likely via apple. The fly could have expanded its range faster than the data indicate if the fewer traps deployed in Kittitas and Yakima Counties in earlier than later years of the survey (Table 2) were insufficient to detect the low populations. However, this still would not contradict the idea of slow fly dispersal and establishment in Yakima County, because traps that numbered in the hundreds in the earlier years in that county consistently failed to catch a fly. The likelihood of flies being present but not detected would have been greater in Kittitas County from 1988–1996, when 39–123 traps were deployed, than in Yakima County. Data from the 2008 and 2010 larval abundance surveys concur with the trapping survey in indicating much higher abundances of flies infesting *C. douglasii* in Klickitat than in Kittitas and Yakima Counties, thus suggesting slow establishment.

It is not clear why black-fruited hawthorns growing in the remote, forested Klickitat sites seem to be better hosts than ones growing in sites in Kittitas and Yakima Counties. The higher larval infestation levels in *C. douglasii* in Klickitat than in other sites could be due to regional variations in hawthorn fruit quality, resulting in differential larval survival. They could also be due to habitat effects, or, as mentioned before, in regards to general fly populations, an establishment time effect influenced by human transport.

Despite the generally low abundance of *R. pomonella* in black-fruited hawthorns, it is still prudent to diligently monitor hawthorns that are near orchards, because of the zero tolerance for larval infestations and the fact that the hawthorn race of the fly can colonize apples (at least in the eastern U.S.A.) (Reissig and Smith 1978). *R. pomonella* has been reported to travel 666 meters within two weeks (Bourne et al. 1934), so perhaps hawthorns > 666 meters from orchards, especially if separated from them by barren landscape, need only be monitored periodically to ensure populations do not dramatically increase. Flies at the Klickitat sites probably pose little threat to major commercial apple orchards, due to the distance and the geographic barriers (Simcoe Mountains and Satus Pass) between the sites and orchards.

In summary, overall results suggest *R. pomonella* is unlikely to develop high populations rapidly near major commercial apple-growing areas in central Washington, including in black-fruited hawthorns, increasing chances it can be kept out of commercial orchards. Whether the detection survey and insecticide spray program described here have contributed to low fly populations is unclear. Habitat, climate, and

genetic factors could contribute to low overall fly populations, a hypothesis that may be experimentally tested.

Acknowledgements

We thank the managers of the Wenas Wildlife Refuge for use of the study site, Pete Chapman and Janine Jewett for field and laboratory assistance, Clark County Heritage Farm and Blair Wolfley and Doug Stienbarger of Washington State University Extension for providing laboratory space, the many employees who assisted in the WSDA apple maggot trapping surveys from 1981–2010, and Grant McQuate (USDA-ARS, Hilo, HI), Dave Horton (USDA-ARS, Wapato, WA), Ed Lisowski (Washington State Department of Agriculture, Yakima, WA), and Michael Bush (Washington State University, Yakima, WA) for helpful comments on the manuscript.

References

- AliNiazee MT, Penrose RL. 1981. Apple maggot in Oregon: a possible new threat to the Northwest apple industry. *Bulletin of the Entomological Society of America* 27: 245–246.
- AliNiazee MT, Wescott RL. 1986. Distribution of the apple maggot, *Rhagoletis pomonella* (Diptera: Tephritidae) in Oregon. *Journal of the Entomological Society of British Columbia* 83: 54–56.
- Bourne AI, Thies WH, Shaw FR. 1934. Some observations on long distance dispersal of apple maggot flies. *Journal of Economic Entomology* 27: 352–355.
- Brunner JF. 1987. Apple maggot in Washington state: a review with special reference to its status in other western states. *Melandria* 45: 33–51.
- Brunsfeld SJ, Johnson FD. 1990. Cytological, morphological, ecological and phenological support for specific status of *Crataegus suksdorfii* (Sarg.) Kruschke. *Madroño* 37: 274–282.
- Bush GL. 1966. The taxonomy, cytology, and evolution of the genus *Rhagoletis* in North America (Diptera, Tephritidae). *Bulletin of the Museum of Comparative Zoology* 134: 431–562.
- Bush GL, Smith JJ. 1998. The genetics and ecology of sympatric speciation: a case study. *Researches on Population Ecology* 40: 175–187.
- Feder JL, Chilcote CA, Bush GL. 1988. Genetic differentiation between sympatric host races of *Rhagoletis pomonella*. *Nature* 336: 61–64.
- Lyons CP, Merilees B. 1995. *Trees, shrubs and flowers to know in Washington and British Columbia*. Lone Pine Publishing.
- Phipps JB. 1998. Introduction to the red-fruited hawthorns (*Crataegus*, Rosaceae) of western North America. *Canadian Journal of Botany* 76: 1863–1899.
- Reissig WH, Smith DC. 1978. Bionomics of *Rhagoletis* in *Crataegus*. *Annals of the Entomological Society of America* 2: 155–159.
- Rull J, Aluja M, Feder J, Berlocher S. 2006. Distribution and host range of hawthorn-infesting *Rhagoletis* (Diptera: Tephritidae) in Mexico. *Annals of the Entomological Society of America* 99: 662–672.

Tracewski KT, Brunner JF, Hoyt SC, Dewey SR. 1987. Occurrence of *Rhagoletis pomonella* (Walsh) in hawthorns, *Crataegus*, of the Pacific Northwest. *Melandieria* 45: 19–25.

USDA. 2010. *National agricultural statistics service*. United States Department of Agriculture. Available online: www.nass.usda.gov

Walsh BD. 1867. The apple-worm and the apple maggot. *American Journal of Horticulture* 2: 338–343.

Wescott RL. 1982. Differentiating adults of apple maggot, *Rhagoletis pomonella* (Walsh) from snowberry maggot, *R. zephyria* Snow (Diptera: Tephritidae) in Oregon. *The Pan-Pacific Entomologist* 58: 25–30.

WSDA. 2001. Washington Administrative Code 16-470-108. *Distribution of infested or damaged fruit is prohibited*. Washington State Department of Agriculture. Available online: <http://agr.wa.gov/>

WSDA. 2010. *Apple maggot – unwanted in Washington*. Washington State Department of Agriculture. Available online: <http://agr.wa.gov/PlantsInsects/InsectPests/AppleMaggot/>

Yee WL. 2008. Host plant use by apple maggot, western cherry fruit fly, and other *Rhagoletis* species (Diptera: Tephritidae) in central Washington state. *The Pan-Pacific Entomologist* 84: 163–178.

Yee WL, Landolt PJ, Darnell TJ. 2005. Attraction of apple maggot (Diptera: Tephritidae) and nontarget flies to traps baited with ammonium carbonate and fruit volatile

lures in Washington and Oregon. *Journal of Agricultural and Urban Entomology* 22: 133–149.

Yee WL, Goughnour RB. 2008. Host plant use by and new host records of apple maggot, western cherry fruit fly, and other *Rhagoletis* species (Diptera: Tephritidae) in western Washington state. *The Pan-Pacific Entomologist*. 84: 179–193.

Table 1. Kittitas, Yakima, and Klickitat County sites sampled for black-fruited hawthorns in 2008 and 2010.

Site	County	Main Location	Approximate Lat., Long.
Ronald	Kittitas	Near Pacific Ave	47.24, -121.03
Roslyn	Kittitas	Washington Ave	47.22, -121.00
Cle Elum	Kittitas	N of Columbia Ave	47.20, -120.92
Wenas	Yakima	Wenas Wildlife Refuge	46.84, -120.72
Nile	Yakima	Near Nile Road	46.84, -120.95
Tampico	Yakima	Ahtanum Road	46.56, -120.91
Toppenish	Yakima	Off Highway 82	46.41, -120.31
Goldendale	Klickitat	Along Hill Road	45.84, -120.91
Middle Klickitat	Klickitat	Liedl Park, Stinson Flats	45.94, -121.12
Lower Klickitat	Klickitat	Ice House Access	45.81, -121.12

Washington State Department of Agriculture’s detection survey trapped mostly feral apples, crabapples, and ornamental hawthorns at these sites (or nearby areas). The Ronald, Roslyn, and Cle Elum sites were trapped sporadically at low levels by the Washington State Department of Agriculture from the 1990s to early 2000s; the Wenas site had never been trapped; the Nile site was not trapped until 2004; areas near Tampico and Toppenish sites had been trapped since the early 1980s; the Goldendale site was trapped from the early to late 1980s; the two Klickitat sites were trapped heavily until the early 1990s.

Table 2. Surveys of *Rhagoletis pomonella* catches on sticky yellow rectangles baited with ammonium carbonate or bicarbonate in native hawthorns, ornamental hawthorn, crabapple, and apple trees in Kittitas, Yakima, and Klickitat Counties in central Washington from 1981–2010. Each tree had one trap at a time over the season, but the trap was replaced when needed. Some trees had multiple fly catches.

Year	Kittitas County			Yakima County			Klickitat County ^a		
	No. Trees	No. Flies	% Traps Positive For Flies	No. Trees	No. Flies	% Traps Positive For Flies	No. Trees	No. Flies	% Traps Positive For Flies
1981	—	—	—	—	—	—	110 ^a	24	10.000 ^a
1982	57	0	0	217	0	0	1,202 ^a	185 ^b	4.243 ^a
1983	161	0	0	369	0	0	2,048 ^a	792	8.545 ^a
1984	229	0	0	250	0	0	1,274	148	6.122
1985	313	0	0	203	0	0	4,365	266	3.872
1986	248	0	0	735	0	0	1,877	36	1.705
1987	214	0	0	802	0	0	1,031	> 9	0.873
1988	39	0	0	641	0	0	355	> 9	2.535
1989	35	0	0	606	0	0	383	> 15	3.916
1990	34	0	0	361	0	0	251 ^a	> 23	9.163 ^a
1991	37	0	0	369	0	0	99	> 3	3.03
1992	26	0	0	427	0	0	248	0	0
1993	37	0	0	643	0	0	9	0	0
1994	82	0	0	1,027	0	0	—	—	—
1995	123	0	0	1,164	1 ^c	0.086	12	—	—
1996	69	0	0	1,617	5	0.309	23	1	4.348
1997	139	1 ^d	0.719	3,850	1	0.026	40	> 8	20
1998	339	2	0.59	3,564	2	0.056	40	0	0
1999	437	1	0.229	3,634	0	0	30	8	3.333
2000	937	12	1.174	1,707	3	0.117	30	48	26.67
2001	1,835	36	1.308	1,869	0	0	—	—	—
2002	2,048	21	0.977	1,483	1	0.067	—	—	—
2003	2,297	26 ^e	1.132	1,402	8	0.571	—	—	—
2004	2,893	37	1.002	2,001	39 ^f	1.399	—	—	—
2005	3,496	145	3.289	2,510	215 ^f	4.741	43	108	41.86
2006	3,481	68	1.695	4,260	18	0.305	6 ^g	0	0
2007	3,650	87	1.918	4,482	13	0.29	8 ^g	0	0
2008	2,801	91	2.785	3,584	8	0.223	7 ^g	0	0
2009	3,044	85	2.497	3,853	19	0.493	6 ^g	0	0
2010	679	7	1.031	4,112	44	0.754	6 ^g	0	0

^aIn 1981, 1982, 1983, and 1990, catches from eastern Skamania County were included in Klickitat County.

^bIncludes first catch (1 fly) in the town of Klickitat.

^cUnion Gap.

^dEllensburg.

^eMostly around Ellensburg beginning in 2003.

^fMostly around Nile in 2004 and 2005; not trapped in subsequent years.

^gLimited trapping in extreme eastern portion of county. — = not applicable, not collected, or records not found.

Table 3. Larval abundance of *Rhagoletis pomonella* in native black-fruited hawthorns in central Washington in 2008.

Site	Collection Date	Species	No. Trees Sampled	Fruit Sample Weight (g)	Approx. No. Fruit ^a	% Trees Positive	No. Larvae	No. Larvae Per g of Fruit	No. Larvae Per Fruit ^a
Nile	12-Aug	<i>C. douglasii</i>	19	7,892	13,153	10.5	2	0.00025	0.00015
Wenas	14-Aug	<i>C. douglasii</i>	8	2,835	4,725	0	0	0	0
	21-Aug	<i>C. douglasii</i>	2	773	1,288	0	0	0	0
	28-Aug	<i>C. douglasii</i>	9	7,842	13,070	11.1	1	0.00013	0.00008
	3-Sep	<i>C. douglasii</i>	13	9,410	15,683	0	0	0	0
	11-Sep	<i>C. douglasii</i>	6	4,471	7,452	0	0	0	0
	23-Sep	<i>C. douglasii</i>	2	1,228	2,047	0	0	0	0
Wenas Totals			40	34,451	57,418	2.5	1	0.00003	0.00002
Tampico	19-Aug	<i>C. douglasii</i>	18	12,202	20,337	22.2	11	0.0009	0.00054
	5-Sep	<i>C. douglasii</i>	3	1,117	1,862	0	0	0	0
Tampico Totals			21	13,319	22,198	19	11	0.00083	0.0005
Toppenish	19-Aug	<i>C. douglasii</i>	4	1,102	1,837	0	0	0	0
Goldendale	26-Aug	<i>C. douglasii</i> , <i>C. suksdorfii</i>	34	12,881	21,468	5.9	2	0.00016	0.00009
Overall Totals			116 ^b	61,753	102,922	7.8	16	0.00026	0.00016

^aApproximate, based on 0.6 g per fruit.
^bTrees sampled multiple times counted once.

Table 4. Larval abundance of *Rhagoletis pomonella* in native black-fruited hawthorns in central Washington in 2010.

Species	Site	Collection Date	No. Trees Sampled	Fruit Sample Weight (g)	Mean g Per Fruit ± SE	No. Fruit	% Trees Positive	No. Larvae	No. Larvae Per g Fruit	No. Larvae Per Fruit
Black Hawthorn, <i>Crataegus douglasii</i>	Ros, Ron, Cle ^a	17-Aug	6	573.739	0.569 ± 0.045	1,014	0	0	0	0
		24-Aug	6	1,739.53	0.569 ± 0.037	2,856	0	0	0	0
		2-Sep	6	2,035.44	0.561 ± 0.084	3,380	0	0	0	0
		8-Sep	19	6,856.09	0.651 ± 0.026	10,105	10.5	6	0.00088	0.00059
		15-Sep	4	2,560.55	0.566 ± 0.095	3,893	50	3	0.00117	0.00077
	Ros, Ron, Cle Totals		19 ^b	13,765.34	—	21,248	21.1	9	0.00065	0.00042
	Wenas	10-Sep	14	4,015.75	0.721 ± 0.023	5,531	14.3	2	0.0005	0.00036
		17-Sep	4	1,908.16	0.756 ± 0.053	2,542	75	18	0.00943	0.00708
	Wenas Totals		18	5,923.91	—	8,073	27.8	20	0.00338	0.00248
	Goldendale	22-Sep	15	2,330.11	0.567 ± 0.036	3,991	26.7	8	0.00343	0.002
	Klickitat ^c	15-Aug	1	—	—	2,109	—	164	—	0.07776
		16-Sep	12	2,546.44	0.550 ± 0.042	4,538	91.7	1,063	0.41745	0.23424
		29-Sep	2	2,980.00	0.617 ± 0.087	4,637	100	1,245	0.41779	0.26849
		6-Oct	4	1,067.39	0.511 ± 0.013	2,044	100	302	0.28293	0.14775
	Klickitat Totals		19	—	—	13,328	94.7	2,774	—	0.20813
	<i>C. douglasii</i> Overall Totals		71 ^b	—	—	46,640	43.7	2,788	—	0.05978
Suksdorf's Hawthorn, <i>Crataegus suksdorfii</i>	Goldendale	22-Sep	9	675.123	0.329 ± 0.018	2,030	11.1	1	0.00148	0.00049
	Klickitat ^c	15-Aug	1	—	—	382	—	1	—	0.00262
		16-Sep	3	27.714	0.233 ± 0.044	124	0	0	0	0
		6-Oct	1	113.789	0.452	252	0	0	0	0
	Klickitat Totals		5	—	—	758	20	1	—	0.00132
	<i>C. suksdorfii</i> Overall Totals		14	—	—	2,788	15.4	3	—	0.00108

^aRoslyn, Ronald, and Cle Elum.
^bTrees sampled multiple times counted once. — = not recorded or incomplete data.
^cMiddle and Lower Klickitat sites combined.



