

## **Microbiology of the Gut of the Kola Nut Weevil, *Balanogastriis kolae***

Authors: Femi-Ola, T. O., and Babalola , A. G.

Source: Journal of Insect Science, 12(84) : 1-6

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.012.8401>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## Microbiology of the gut of the kola nut weevil, *Balanogastris kolae*

T. O. Femi-Ola<sup>a\*</sup>, and A. G. Babalola<sup>b</sup>

Department of Microbiology, Ekiti State University, P.M.B. 5363, Ado-Ekiti, Nigeria

### Abstract

Reports have shown that many insects have microbes in their gut system. Gut microbes are very important for insect vitality and much of their nutrition is derived from products of microbial metabolism. The habitat of *Balanogastris kolae* (Desbrocher des Loges) (Coleoptera: Curculionidae) suggests that they possess the ability to digest varieties of sugars particularly starch and protein materials present in the kola nut, *Cola nitida* Schott & Endlicher (Malvales: Malvaceae). The aim of this study was to characterize the gut bacterial communities of the kola weevil, *B. kolae*. To ascertain this, the gut bacterial community of a kola nut-feeding weevil, *B. kolae* was characterized using culture-dependent methods. The bacterial counts in the foregut, midgut and hindgut were  $7.14 \pm 0.11 \times 10^6$  cfu ml<sup>-1</sup>,  $2.68 \pm 0.13 \times 10^7$  cfu ml<sup>-1</sup> and  $1.43 \pm 0.20 \times 10^6$  cfu ml<sup>-1</sup> respectively. There were no significant differences in the total bacterial count of the foregut, midgut and hindgut. The bacterial species were identified to be *Fusobacterium nucleatum*, *Staphylococcus aureus*, *Bacillus subtilis*, *Corynebacterium fascians*, *Arthrobacter globiformis*, *Serratia marcescens*, *Bacillus brevis*, *Vibrio haemolyticus* and *Flavobacterium breve*. The majority of these isolates were demonstrated to have both proteolytic and amylolytic activities.

**Keywords:** bacteria, *Cola nitida*

**Correspondence:** <sup>a</sup> [titifemi2006@yahoo.com](mailto:titifemi2006@yahoo.com), \*Corresponding author

**Editor:** Allen Cohen was editor of this paper.

**Received:** 21 September 2009, **Accepted:** 15 October 2010

**Copyright :** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 12, Number 84

#### Cite this paper as:

Femi-Ola TO, Babalola AG. 2012. Microbiology of the gut of the kola nut weevil, *Balanogastris kolae*. *Journal of Insect Science* 12:84 available online: [insectscience.org/12.84](http://insectscience.org/12.84)

## Introduction

The weevils and other plantation pests are associated with a rich and complex community of microorganism in their guts and other body regions. This microbiota participates in many kinds of interactions ranging from pathogenesis to obligate mutualism (Dillon and Dillon, 2004). There has been a renewed interest in the understanding of insect gut, because they are potential source of enzymes (Zhang and Brune, 2004) and metabolites (Wilkinson, 2001). It has been reported that the manipulation of the microbial symbionts may be an effective strategy for controlling spread of pathogens that use pests as hosts (Dillon et al., 2005).

Weevils are the major pest associated with the spoilage of nuts and other crop plants. Among the weevil pests, the species found in Nigeria associated with kola nut are *Balanogastrius kolae*, *Sophrorhinus insperatus*, *Sophrorhinus duveroyi* and *Sophrorhinus gbajensis*. The digestion of the kola nut by the kola-feeding weevil *Sophrorhinus insperatus* has been linked to the activities of enzymes present in the gut of the weevil. The dominant enzymes are the carbohydrases and proteases (Adedire, 1994). The enzymatic activities of the weevil have been reported to vary with the different segments of the gut; the foregut, midgut and hindgut. Enzymatic activities have been reported to be more prominent in the midgut (Adedire, 1994). However, there is a dearth of information on the microbial populations in the gut of kola nut feeding weevil and their possible role in the digestion of kola nut in the gut of the weevil. The knowledge of bacterial species composition will facilitate the studies of the functions of the gut microbiota.

The aim of this study was to characterize the gut bacterial communities of the kola weevil, *B. kolae* (Desbrocher des Loges) (Coleoptera: Curculionidae), using classical microbiological methods. This study also evaluates the production of some of the enzymes required for the digestion of kola nut in the gut of the kola feeding weevil.

## Materials and Methods

### Collection of materials

Kola nut weevils were collected from field-infested kola nuts, *Cola nitida* Schott and Endlicher (Malvales: Malvaceae) and were kept until emergence in a woven basket lined with banana leaves at insectary temperature of  $28 \pm 2^\circ\text{C}$  and  $80 \pm 5\%$  RH. The kola weevil samples were identified as *B. kolae*. There was no discrimination with respect to the sex of the insect used.

### Assessment of microflora in the gut of *B. kolae*

Approximately twenty weevil samples were selected for the isolation of gut bacteria. The weevils were anaesthetized in a refrigerator for 15 minutes and then surface sterilized by rinsing them in 10% v/v sodium hypochlorite and distilled water to kill external bacteria. Thereafter they were rinsed in two changes of sterile distilled water. They were aseptically dissected to remove the intestine. Each of the intestines was divided into three regions namely foregut, midgut and hindgut. The respective gut region was homogenized in 10 ml sterile Ringer's solution containing the following ( $\text{g L}^{-1}$ ): NaCl, 10.0; KCl, 0.42;  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , 0.48 and  $\text{Na}_2\text{HCO}_3$ , 0.2 (Femi-Ola and Aderibigbe, 2009). The gut sample was then crushed with a sterile glass rod to release the intestinal contents. Serial dilution of gut sample was carried out in sterile

dilution blanks of distilled water and plated on nutrient agar (NA) plates in replicates and incubated at 37° C for 48h. Colonies on plates were counted to determine the total colony forming units of bacterial (cfu) counts. Pure cultures of the isolates were obtained by sub culturing serially onto sterile nutrient agar plates.

### Characterization and isolation of bacterial isolates

Pure cultures of the isolates were identified on the basis of their cultural, morphological and biochemical characteristics in accordance with the taxonomic scheme of Holt *et al.* (1994) and Barrow and Feltham (1993). The test performed include Gram-stain, spore-stain, motility test, Voges-Proskauer, oxidative/fermentative tests, gelatin hydrolysis, oxygen requirement, growth at different salt concentration and utilization of carbon source.

### Physiological studies of microorganisms in the gut of *B. kolae*

Bacterial isolates were screened for their amylase and protease activities.

Alpha amylase production was tested by inoculating the bacterial culture on starch agar (nutrient agar plus 1% soluble starch). Plates were incubated at 35° C for 3 days, after which they were flooded with iodine solution [0.3% I<sub>2</sub> (w/v) in 3% KI (w/v)]. Amylase was indicated by a clear/brown zone surrounding the colony. Amylase activity was determined quantitatively by the dinitrosalicylic acid (DNSA) method of Bernfield (1955), which measures the increase in the reducing power of the digests in the reaction between starch and the enzyme. One unit of alpha amylase activity was defined as the amount of enzyme required to produce one micromole of maltose from starch under the assay condition.

To determine of protease activity, bacterial isolates were inoculated onto casein agar plates and were incubated at 35° C for 24h. Protease production was indicated by a clear zone of casein hydrolysis. Protease activity was measured quantitatively by the method of Kunitz (1974). One unit of protease activity was defined as the amount of enzyme required to produce one micromole of tyrosine from casein under the assay condition.

### Statistical analysis

Analysis of variance of data obtained was carried out using the statistical package for Social Sciences (SPSS version 11.0).

### Results

The bacterial count in the different region of *B. kolae* is shown in Table 1. The total bacterial count in the foregut, midgut and hindgut were  $7.14 \pm 0.11 \times 10^6$ ,  $2.68 \pm 0.13 \times 10^7$  and  $1.43 \pm 0.20 \times 10^6$  cfu ml<sup>-1</sup> respectively. There was no significant difference in the total bacterial count of the foregut, midgut and hindgut ( $p \geq 0.05$ ). The characterization of the bacterial isolates revealed that nine different species of bacteria were isolated from the gut of *B. kolae*. These bacteria were identified as *Staphylococcus aureus*, *Bacillus subtilis*, *Corynebacterium fascians*, *Fusobacterium nucleatum*, *Arthrobacter globiformis*, *Serratia marcescens*, *Bacillus brevis*, *Vibrio haemolyticus*, *Flavobacterium breves* and *Zymomonas mobilis* (Table 1). The rate of occurrence of the bacterial isolates is shown in Table 2. *B. subtilis*, *C. fascians* and *A. globiformis* had the highest occurrence rate of 66.6%.

Most of the bacterial isolates in this study were both proteolytic and amylolytic (Table

**Table 1.** Total load (cfu/ml) of bacteria in the gut of *Balanogastis kolae*.

Bacterial isolates	Counts (cfu/mL)		
	Foregut	Midgut	Hindgut
<i>Staphylococcus aureus</i>	5.4±0.05 ×10 <sup>6</sup>	-	-
<i>Bacillus subtilis</i>	1.98 ±0.03×10 <sup>5</sup>	4.08 ±0.02×10 <sup>6</sup>	-
<i>Corynebacterium fascians</i>	7.8 ±0.02×10 <sup>5</sup>	4.90± 0.05×10 <sup>5</sup>	-
<i>Fusobacterium nucleatum</i>	7.6 ±0.01×10 <sup>5</sup>	-	-
<i>Arthrobacter globiformis</i>	-	2.10±0.02×10 <sup>7</sup>	9.21±0.03×10 <sup>5</sup>
<i>Serratia marcescens</i>	-	1.25±0.04×10 <sup>6</sup>	-
<i>Bacillus brevis</i>	-	-	3.4±0.06×10 <sup>4</sup>
<i>Vibrio haemolyticus</i>	-	-	1.78±0.03×10 <sup>5</sup>
<i>Flavobacterium breves</i>	-	-	8.0±0.05×10 <sup>4</sup>
<i>Zymomonas mobilis</i>	-	-	2.20±0.03×10 <sup>5</sup>
<b>Total</b>	<b>7.14±0.11×10<sup>6</sup></b>	<b>2.68±0.13×10<sup>7</sup></b>	<b>1.43±0.20×10<sup>6</sup></b>

Cfu = Colony forming unit; - = Absent

**Table 2.** Rate of occurrence of bacterial isolates in the gut of *Balanogastis kolae*.

Bacterial isolates	Incidence (%)
<i>Staphylococcus aureus</i>	33.3
<i>Bacillus subtilis</i>	66.7
<i>Corynebacterium fascians</i>	66.7
<i>Fusobacterium nucleatum</i>	33.3
<i>Arthrobacter globiformis</i>	66.7
<i>Serratia marcescens</i>	33.3
<i>Bacillus brevis</i>	33.3
<i>Vibrio haemolyticus</i>	33.3
<i>Flavobacterium breves</i>	33.3
<i>Zymomonas mobilis</i>	33.3

3), and *B. subtilis* showed the highest amylolytic and proteolytic activities. *S. aureus* and *F. breves* were only proteolytic and *F. nucleatum* did not hydrolyze either starch or casein.

## Discussion

This study has shown that the kola weevil gut contained diverse bacterial species. *B. subtilis*, *A. globiformis* and *C. fascians* were isolated from two different regions of the gut, while the other bacterial species were isolated from a single region. This observation corroborates the report of Dillon and Dillon (2004), that bacteria are associated with a number of different insect species across all major orders of the insects. Eka (1971) had reported that the bulk of carbohydrate digestion takes place in the midgut of kola weevil. Carbohydrases

**Table 3.** Enzymatic (amylase and protease) activities of bacteria isolated from the gut of *Balanogastis kolae* feeding weevil.

Bacteria	Enzyme activity (U/mg protein)	
	Amylase	Protease
<i>Bacillus subtilis</i>	0.66	10.66
<i>Corynebacterium fascians</i>	0.61	5.21
<i>Vibrio haemolyticus</i>	0.46	4.12
<i>Bacillus brevis</i>	0.64	3.62
<i>Flavobacterium breve</i>	0	6.24
<i>Arthrobacter globiformis</i>	0.57	9.38
<i>Zymomonas mobilis</i>	0.43	13.72
<i>Serratia marcescens</i>	0.47	8.32
<i>Staphylococcus aureus</i>	0	6.62
<i>Fusobacterium nucleatum</i>	0	0.08

and proteolytic activities have been reported by Adedire (1994) in the gut of the kola weevil, *Sophrohinus insperatus*, with the bulk of enzymatic activities in the midgut of the insect. In our study, most of the bacterial isolates were recovered from the midgut that is the site where most digestive activities (carbohydrate and protein utilization) were taking place compared to other sites of the gut. The high density of bacterial species observed in this region probably indicate the involvement of these organisms in the digestion of carbohydrate in the kola nut that forms the major diet of the weevil *B. kolae*.

Microorganisms in the gut of various insects have been reported to be involved in the degradative activities in the gut (Todaka et al., 2007; Bera-Maillet et al., 2000). In this study, most of the bacterial isolates were found to be both proteolytic and amylolytic. Some of these isolates may be relevant in relation to

the diet of the insect as they produce enzymes that help to break down carbohydrate, protein and fats constituents of the weevil's diets. The organisms isolated are believed to be living in symbiotic relationship with the kola nut weevil that provides the food substrate for the micro flora in the different regions of the weevil gut. However, further research is needed to establish the roles played by the micro flora in the gut. Molecular studies of the gut will reveal the presence or absence of some non-culturable microorganisms that may be playing major roles in the digestion of kola in the gut of the kola nut feeding weevil.

## References

- Adedire CO. 1994. Distribution of carbohydrases and proteases in the intestine of Kola nut weevil, *Sophrorhinus insperatus* FAUST (Coleoptera: Curculionidae) and response of proteases to inhibitors from Kola nuts. *Applied Entomology and Zoology* 29 (3): 331-338.
- Barrow GI, Feltham RKA. 1993. *Cowan and Steel's Manual for the identification of Medical Bacteria*. (3<sup>rd</sup> Edn) Cambridge University Press.
- Bera-Maillet CL, Arthaud P, Rosso MN. 2000. Biochemical characterization of MI-ENGI a family 5 endoglucanase secreted by the root-knot nematode *Meladogyne incognita*. *European Journal of Biochemistry* 267: 3255-3263.
- Bernfield P. 1955. Amylases  $\alpha$  and  $\beta$ . In: *Methods in Enzymology*. Vol.1. Colowick SP, Kaplan NO, Editors. pp. 149-158. New York Press.
- Daramola AM. 1973. The bionomics of the Kola weevils, *Sophrorhinus* spp FAUST (Coleoptera: Curculionidae). Ph.D thesis University of Ibadan, Nigeria.
- Dillon RJ, Dillon VM. 2004. The gut bacteria of insects: non pathogenic interactions. *Annual Review of Entomology* 49: 71-92
- Dillon RJ, Yennard CT, Buckling A, Charnley AK. 2005. Diversity of locust gut bacteria protects against pathogens invasion. *Ecology Letters* 8: 1291-1298.
- Eka OU. 1971. Chemical composition and uses of Kola nuts. *Journal of West African Science* 16: 167-169.
- Femi-Ola TO, Aderibigbe EY. 2009. Effects of seasonal changes on the microflora in the hindgut of wood-eating termites *Amitermes evuncifer* Silvestri. *Journal of Entomology* 6 (1): 67-71.
- Holt JG, Kreig NR, Sneath PHA, Staley JT, Williams ST. 1994. *Bergey's Manual of Determinative Bacteriology* 9th Edn. Williams and Wilkins, Baltimore.
- Kunitz M. 1974. Crystalline soybean trypsin inhibitor. II. General properties. *Journal of General Physiology* 30: 291-310.
- Todaka N, Moriya S, Saita K, Hondo T, Kiuchi I, Takasu H, Ohkuma M, Piero C, Hayashizaki Y, Kudo T. 2007. Analysis of the genes involved in lignocellulose digestion in the symbiotic protist community of *Reticulitermes insperatus* *FEMS Microbio. Ecol.* 59(3): 592-599.
- Wilkinson T. 2001. Disloyalty and treachery in bug-swapping shocker. *Trends in Ecology and Evolution* 16: 659-661.

Zhang H. Brune A. 2004. Characterization and partial purification of proteinases from highly alkaline midgut of the humivorous larvae of *Pachnoda ephippiata* (Coleoptera: Scarabaeidae) *Soil Biology and Biochemistry* 36: 435-442.