

Genetic diversity and differentiation among populations of the Indian eri silkworm, *Samia cynthia ricini*, revealed by ISSR markers

Authors: Vijayan, K., Anuradha, H. J., Nair, C. V., Pradeep, A. R., Awasthi, A. K., et al.

Source: Journal of Insect Science, 6(30) : 1-11

Published By: Entomological Society of America

URL: https://doi.org/10.1673/2006_6_30.1

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Genetic diversity and differentiation among populations of the Indian eri silkworm, *Samia cynthia ricini*, revealed by ISSR markers

K Vijayan^{1,\$}, H. J. Anuradha¹, C. V. Nair¹, A. R. Pradeep^{1,\$*}, A. K. Awasthi¹, B. Saratchandra^{1,#}, S. A. S. Rahman², K. C. Singh², R. Chakraborti² and S. Raje Urs¹

¹ Seribiotech Research Laboratory, Central Silk Board, CSB Campus, Kodathi, Carmelram P.O., Bangalore 560 035, Karnataka, India

² Central Muga Eri Research and Training Institute, Jorhat, Assam, India

\$ contributed equally

Present address: Central Silk Board, B.T.M. Lay Out, Bangalore – 560035, Karnataka, India

Abstract

Samia cynthia ricini (Lepidoptera: Saturniidae), the Indian eri silkworm, contributes significantly to the production of commercial silk and is widely distributed in the Brahmaputra river valley in North-Eastern India. Due to over exploitation coupled with rapid deforestation, most of the natural populations of *S. cynthia ricini* are dwindling rapidly and its preservation has become an important goal. Assessment of the genetic structure of each population is a prerequisite for a sustainable conservation program. DNA fingerprinting to detect genetic variation has been used in different insect species not only between populations, but also between individuals within a population. Since, information on the genetic basis of phenotypic variability and genetic diversity within the *S. cynthia ricini* populations is scanty, inter simple sequence repeat (ISSR) system was used to assess genetic diversity and differentiation among six commercially exploited *S. cynthia ricini* populations. Twenty ISSR primers produced 87% of inter population variability among the six populations. Genetic distance was lowest between the populations Khanapara (E5) and Mendipathar (E6) (0.0654) and highest between Dhanubhanga (E4) and Titabar (E3) (0.3811). Within population, heterozygosity was higher in Borduar (E2) (0.1093) and lowest in Titabar (E3) (0.0510). Highest gene flow (0.9035) was between E5 and E6 and the lowest (0.2172) was between E3 and E5. Regression analysis showed positive correlation between genetic distance and geographic distance among the populations. The high G_{ST} value (0.657) among the populations combined with low gene flow contributes significantly to the genetic differentiation among the *S. cynthia ricini* populations. Based on genetic diversity, these populations can be considered as different ecotypes and *in situ* conservation of them is recommended.

Keywords: Eri phenotype, geographic isolation, gene flow, heterozygosity

Abbreviations: *Samia cynthia ricini* populations: Nongpoh (E1), Borduar (E2), Titabar (E3), Dhanubhanga (E4), Khanapara (E5), Mendipathar (E6)

Correspondence: *arpradeep23@rediffmail.com, kvijayan01@yahoo.com, anujnh@yahoo.co.in, cvgn_2000@yahoo.com, arvind.awasthi@rediffmail.com, sarat@silkboard.org, sasrahman@hotmail.com kchobasingh@yahoo.co.in, rajenchakravorty13@yahoo.com, srajeurs@yahoo.com

Received: 19.9.2005 | Accepted: 2.4.2006 | Published: 20.10.2006

Copyright: Creative Commons Attribution 2.5

ISSN: 1536-2442 | Volume 6, Number 30

Cite this paper as:

Vijayan K, Anuradha HJ, Nair CV, Pradeep AR, Awasthi AK, Saratchandra B, Rahman SAS, Singh KC, Chakraborti R, Urs SR. 2006. Genetic diversity and differentiation among populations of the Indian eri silkworm, *Samia cynthia ricini*, revealed by ISSR markers. 11pp. *Journal of Insect Science* 6:30, available online: insectscience.org/6.30

Introduction

Understanding and preserving biodiversity is one of the most important global challenges that biologists are facing. Widespread realization that biodiversity is strongly modified by changes in global environment has generated strategies to conserve and protect biodiversity in many parts of the world. Assessment of the genetic diversity present within a species is a prerequisite for developing a sustainable conservation program. North-eastern India belongs to the trans-Himalayan mountainous zone and is characterized by a tropical humid climate with temperature varying from 10 to 24° C, a range of relative humidity from 75–98% and rainfall of 1300–1700 mm per year. 65% of the area is mountainous with evergreen forests and it includes the Brahmaputra river valley. The forested areas are the abode of many valuable flora and fauna including about ten species of wild silk moths (Chowdhury 1983; Thangavelu 1991). The Indian eri silkworm, *Samia cynthia ricini* (Lepidoptera: Saturniidae), a commercial silk producing insect, is believed to have originated in the Brahmaputra valley (Jolly et al. 1979) and has had restricted distribution in India, China and Japan for two centuries (Peigler 1993; Singh and Benchamin 2002). The primary food plant of this polyphagous insect is castor (*Ricinus communis* L.), but it also feeds on a wide range of food plants such as *Heteropanax fragrans* Seem, *Manihot utilissima* Phol, *Evodia flaxinifolia* Hook, *Ailanthus gradulosa* Roxb etc (Suryanarayana et al. 2002). The wild *S. cynthia ricini* silkworm completes one to three generations per year depending on geographical position and climatic conditions of the region, however, up to six generations occur in the domesticated cultures (Neupane et al. 1990). Populations of *S. cynthia ricini*, that have been commercially exploited and are present in different regions of north-east India showed wide variations in morphological and quantitative characters such as absolute silk content, larval weight, cocoon weight, cocoon shell weight and silk ratio (Siddiqui et al. 2000). These populations were named as Nongpoh, Kokrajhar Red, Borduar, Titabar, Sille, Dhanubhanga, Mendipathar and Khanapara, after their place of collection from where they were originally reported and were abundantly present. Due to over exploitation of the silkworms for commercial uses

coupled with deforestation, most of these natural populations are dwindling rapidly. In order to preserve the natural biodiversity present among these populations, attempts are being made to understand the genetic structure of each population. Preliminary studies based on some quantitative traits such as cocoon weight, shell weight, larval weight etc. were made to understand the genetic basis of this phenotypic variability (Siddiqui et al. 2000). However, no systematic studies were made to generate substantial information on the genetic diversity of these populations so as to develop appropriate strategy for its conservation at the natural habitat. The present study was focused on the genetic diversity of six commercial populations of *S. cynthia ricini* present in north-eastern India.

The advent of molecular biological techniques clearly showed the advantages of molecular markers over morphobiochemical markers to analyze population diversity. As the molecular markers are stable and environmentally independent, they are increasingly being preferred to phenotypic traits to detect genetic variation not only among populations but also between individuals within a population. A number of DNA marker systems such as simple sequence repeats (SSR; Kimpton et al. 1993; Estoup et al. 1993; Reddy et al. 1999a; Prasad et al. 2005), random amplified polymorphic DNA (RAPD; Williams et al. 1990; Nagaraja and Nagaraju 1995; Chatterjee and Pradeep 2003), inter-simple sequence repeats (ISSR; Zietkiewicz et al. 1994; Ehtesham et al. 1995; Reddy et al. 1999b; Chatterjee et al. 2004; Kar et al. 2005; Pradeep et al. 2005), expressed sequence tag (EST; Vlachou et al. 1997; Ciolfi et al. 2005) and amplified fragment length polymorphism (AFLP; Vos et al. 1995; Reineke et al. 1998; Katiyar et al. 2000) have been used to study the population genetics of different organisms including insects. Genetic diversity and differentiation among different populations of the wild silkworm *Antheraea mylitta* was examined using ISSR markers (Chatterjee et al. 2004; Kar et al. 2005). Use of SSR and EST primers requires prior knowledge of the genome of the organism whereas development of ISSR primers does not require prior information of the genome. The information available on the genome of *S. cynthia ricini* is

inadequate to develop SSR and EST primers for assessing genetic diversity and genetic differentiation among its populations. As short repeats, such as dinucleotides, are ubiquitously distributed in eukaryotic genomes (Ichimura and Mita 1992; Goldstein and Schlötterer 1999; Prasad et al. 2005; Pradeep *et al.* unpublished), primers representing these repeats (eg. ISSR primers; Zietkiewicz et al. 1994) produce complex fingerprints from distantly related organisms. ISSR markers are reliable, reproducible, polymorphic and have been used to estimate genetic diversity among closely related populations also (Vogel and Scolnik 1997; Abbot 2001; Deshpande et al. 2001). Considering these advantages of ISSR primers, this marker system was used to analyze the genetic variability among *S. cynthia ricini* silkworm populations.

Materials and Methods

Genetic material and DNA extraction

Six morphologically distinct populations of *S. cynthia ricini*, collected from different regions of North-eastern India (Table 1, Figure 1), were used for the study: Nongpoh (E1), Borduar (E2), Titabar (E3), Dhanubhanga (E4), Khanapara (E5) and Mendipathar (E6). From each population, 10 to 15 cocoons were collected and kept until emergence of the adult moth. Genomic DNA from 10 individual moths of each population was extracted separately following the phenol: chloroform extraction method (Suzuki et al. 1972). After RNase incubation, DNA was re-extracted and the purified DNA was diluted in TE (Tris-EDTA; pH 8.0) buffer to obtain the concentration of DNA at 10ng/μl.

Table 1. Key to the phenotypic characters of six different populations of the Indian eri silkworm, *S.cynthia ricini*

Characters	Nongpoh (E1)	Borduar (E2)	Titabar (E3)	Dhanubhanga (E4)	Khanapara (E5)	Mendipathar (E6)
Distribution	Nongpoh, Meghalaya	Kamrup, Assam	Jorhat, Assam	Goalpara, Assam	Kamrup, Assam	East Garo hills, Meghalaya
Voltinism	multivoltine	multivoltine	multivoltine	multivoltine	multivoltine	multivoltine
Larval colour	Yellow, Greenish blue	Yellow, Greenish blue	Yellow, Greenish blue	Yellow, Greenish blue	Greenish blue	Greenish blue
Larval marking	Plain	Zebra	Plain	Zebra	Plain	Plain
Total larval duration	15–18 days	18–20 days	15–20 days	15–17 days	15–18 days	20–30 days
Larval weight (gm).	9.58±1.70	6.87±1.01	5.56±0.68	10.73±1.22	9.61±1.73	9.11±0.81
Cocoon colour	Pale white	Creamy white	White	White	Creamy white	Creamy white
Cocoon shape	Oval	Oval	Oval	Oval	Oval	Oval
Cocoon wt.(gm)	3.19±0.38	3.09±0.24	3.12±0.34	3.45±0.27	3.06±0.39	3.72±0.32
Cocoon shell wt.(gm)	0.45±0.03	0.47±0.017	0.44±0.01	0.50±0.01	0.48±0.02	0.56±0.10
Moth colour	both sexes blackish gray	both sexes blackish gray	both sexes blackish gray	both sexes blackish gray	both sexes blackish gray	males blackish gray females light blackish gray




Figure 1. Map of North-Eastern India showing the places of collection of the eri silkworm, *S.c. ricini* populations. E1–E6 was the six populations as given in Table 1 (map not to scale).

PCR amplification of the genomic DNA with ISSR primers

ISSR primers (Table 2), procured from Dr. John B. Hobbs, Nucleic Acid & Protein Service, Biotechnology Laboratory, University of British Columbia, Vancouver, Canada, (<http://naps.biotech.ubc.ca>) (set # 9), were tested for their efficacy in amplification of DNA. Twenty primers that produced reproducible robust bands, which appeared consistently and distinctly across three different amplifications, were selected for the study. PCR amplification of the DNA was carried out as described earlier (Chatterjee et al 2004) on an MJ Research Thermal-Cycler, PTC 200 (MJ Research Inc. www.mjr.com/), using 20 µl reaction mixture containing 10x PCR buffer, 2 mM dNTPs, 2.5 mM MgCl₂, 0.10 µl *Taq* DNA polymerase (recombinant) (5 U/µl) (all the chemicals from MBI Fermentas; Fermentas Inc, www.fermentas.com/), 2.0 µl of 1.5 µM ISSR primer and 40 ng DNA.

The PCR products were resolved on 1.5% agarose gel, in Tris - Boric Acid–EDTA buffer, stained with ethidium bromide (0.5 µg/ml) and photographed under Bioimaging system (Syngene, www.syngene.com). Binary scoring of the profiles was done on the basis of presence or absence of a band at a particular locus. The amplification of the DNA with each primer was repeated thrice and only the robust bands were scored for the analysis.

Statistical analyses

From the binary data, genetic dissimilarity coefficients among the individuals were estimated using Dice’s coefficient ($D = 1 - [(2N_{ab}) / (2N_{ab} + N_a$

+ $N_b)]$, where N_{ab} is the number of bands that are shared by the genotypes ‘a’ and ‘b’; N_a is the number of bands present in ‘a’ and N_b is the number of bands present in ‘b’ (Sneath and Sokal 1973). A phylogeny tree was generated from the above matrix using the unweighted pair group method with arithmetical averages (UPGMA) (Sneath and Sokal 1973). Bootstrap analysis was conducted using 10000 replicates as implemented in PAUP 4.0 software (Swofford 1999). Nei’s (1972) genetic distance among the phenotypes was calculated using POPGENE Version 1.3 (Yeh 1998). A dendrogram for the six populations was also created using UPGMA analysis (Sneath and Sokal 1973). The PhiPT value for genetic variability was calculated using GenAlEx v 5.1. The percentage of polymorphism, heterozygosity (h), number of alleles (N_a), number of effective alleles (N_e) and Shannon’s information index (I) were calculated for each population. Genetic variability in the populations was also calculated using Nei’s (1973) coefficient of gene differentiation (G_{ST}) in POPGENE version 1.3 (Yeh 1998). In POPGENE the genetic divergence among different populations was calculated using a multiallelic analogue of F_{ST} among a finite number of populations, which is otherwise, called the coefficient of gene differentiation (Nei 1973). This is stated as $G_{ST} = D_{ST} / H_t = (H_t - H_s) / H_t$ where D_{ST} is the average gene diversity between subpopulations, including the comparisons of sub populations with themselves. The $D_{ST} = (H_t - H_s)$. G_{ST} is an extension of Nei’s (1972) genetic distance between a pair of populations to the case of hierarchical structure of populations (Nei 1973). $H_t = (1 - \sum p_i^2)$, where p_i is the frequency of i^{th} allele at a locus in a population

Table 2. Polymorphism detected by 20 ISSR primers employed in the genetic diversity studies on six populations of eri silkworm *Samia cynthia ricini*

Primer code (UBC)	Sequence (5'-3')	Total bands	Polymorphic bands
807	AGAGAGAGAGAGAGT	5	4
808	AGAGAGAGAGAGAGC	3	0
809	AGAGAGAGAGAGAGG	5	2
810	GAGAGAGAGAGAGAT	5	4
811	GAGAGAGAGAGAGAC	4	0
812	GAGAGAGAGAGAGAA	4	3
825	ACACACACACACACT	4	2
826	ACACACACACACACC	6	5
834	AGAGAGAGAGAGAGYT	3	2
835	AGAGAGAGAGAGAGYC	4	0
840	GAGAGAGAGAGAGAYT	4	3
842	GAGAGAGAGAGAGAYG	4	1
855	ACACACACACACACYT	6	2
864	ATGATGATGATGATG	6	2
873	GACAGACAGACAGACA	2	0
880	GGAGAGGAGAGGAGA	6	6
881	GGGTGGGGTGGGGTG	5	4
884	HBHAGAGAGAGAGAG	4	2
885	BHBGAGAGAGAGAGAGA	5	5
886	VDVCTCTCTCTCTCT	7	6
TOTAL		92	53

*Y=(C,T); H=(ACT); B=(C,G,T); V=(A,C,G); D=(A,G,T) any one of the base in the place of the respective letter.

and Σ is the summation of all alleles. Hence, the H_s in the equation were defined in terms of gene diversities. However, for random mating subpopulations, gene diversities can be defined as expected heterozygosities under Hardy-Weinberg equilibrium averaged among sub populations (H_s) and of the total population (H_t). The estimate of gene flow from G_{ST} was calculated as $(Nm) = 0.5 (1 - G_{ST})/G_{ST}$. To analyze the relation between genetic distance and geographic distance between the populations, regression analysis was done using inter- population genetic distance against the geographic distance between them in kilometers. A scatter diagram was prepared and the best-fit linear regression line was applied.

Results

Six populations of *S. cyathia ricini* showed variation in morphological and quantitative characters (Table 1). The color of the mature larvae was greenish blue in E5 and E6, and a mixture of yellowish and greenish blue in E1, E2, E3 and E4. Typical zebra markings were present on larvae of E2 and E4 whereas E1, E3, E5 and E6 were plain

white without any markings. The weight of mature larvae recorded was 9.58 ± 1.70 g in E1, 6.87 ± 1.01 g in E2, 5.56 ± 0.68 g in E3, 10.73 ± 1.22 g in E4, 9.61 ± 1.73 g in E5 and 9.11 ± 0.81 g in E6. Cocoon weight varied from 3.06 ± 0.39 g in E5 to 3.72 ± 0.32 g in E6.

Inter population genetic diversity

Twenty ISSR primers generated a total of 92 bands, of which 53 were polymorphic (Table 2; Figure 2). AMOVA revealed 87% of inter-population genetic variability and the Φ_{IPT} value was 0.872, which was significant at a probability of 0.010. Analysis for pair-wise genetic distance revealed that the genetic distance was minimum between E5 and E6 populations (0.0654) and maximum between E4 and E3 (0.3811) populations (Table 3). The intra-population variability in terms of DNA polymorphism was highest in E2 (31.52%) and least in E5 and E6 (14.13%) (Table 4). The dendrogram shown in Figure 3 that resulted from the genetic dissimilarity matrix using UPGMA (Sneath and Sokal 1973) revealed that E5 and E6 were genetically closer than the other populations, and E3 formed an isolate.




Figure 2. An example of PCR amplification profile generated from genomic DNA of *S.c. ricini* populations with UBC886, resolved on 1.5% agarose gel. M- marker.

Table 3. Genetic distance estimated among the six populations of eri silkworm, *Samia cynthia ricini*

Population	Nongpoh (E1)	Borduar (E2)	Titabar (E3)	Dhanubhanga (E4)	Khanapara (E5)
Borduar	0.222				
Titabar	0.222	0.2688			
Dhanubhanga	0.1729	0.2229	0.3811		
Khanapara	0.1098	0.2636	0.3156	0.139	
Mendipathar	0.1262	0.2798	0.2441	0.1854	0.0654

Maximum and minimum genetic distances are shown in bold face

Table 4. Gene diversity in the six populations of the eri silkworm *Samia cynthia ricini*

Accessions	Number of observed alleles (Na)	Number of effective alleles (Ne)	Heterozygosity (h)	Shannon's information index (I)	Polymorphism (%)
Nongpoh (E1)	1.2609	1.1928	0.108	0.1568	26.09
Borduar (E2)	1.3152	1.1961	0.1093	0.1621	31.52
Titabar (E3)	1.1522	1.0886	0.051	0.0762	15.22
Dhanubhanga (E4)	1.2391	1.177	0.0982	0.1421	23.91
Khanapara (E5)	1.1413	1.1134	0.0609	0.0871	14.13
Mendipathar (E6)	1.1413	1.0914	0.0521	0.0769	14.13
Total	1.6413	1.4003	0.2334	0.3475	64.13




Figure 3. Dendrogram showing grouping of the six populations of *S.c. ricini* using UPGMA analysis based on Dice dissimilarity matrix.

Intra population genetic diversity

When all 60 individuals were taken for analysis without assigning any population status, the individuals grouped according to their population affinity (Figure 4). None of the individuals changed its population cluster. The bootstrap values for each population were highly significant and varied from

65 to 100. The grouping of individuals within each population showed variation. Though the individuals of different populations grouped into sub groups, in most cases, the bootstrap values were insignificant. But significantly high bootstrap values (71–99%) were observed in the subgrouping of the E2 population.




Figure 4. Dendrogram showing grouping of 60 individuals from six populations of *S.c. ricini* based on the genetic distance derived from ISSR markers. Bootstrap values were given at the fork of each group. E1–E6 was the six populations as given in Table 1. E1.1 – One individual from the population E1.

Population genetic parameters

The population diversity analysis using POPGENE revealed that the average number of observed alleles varied from 1.143 in E5 and E6 to 1.315 in E2. The heterozygosity present within a population was higher in E2 (0.109) and was lowest in E3 (0.051) (Table 4). When all populations were considered, the total genetic diversity (H_t) was 0.233. The total genetic differentiation coefficient (G_{ST}) among the populations was 0.657 (Table 5). The pair-wise comparison of the populations

showed that the genetic differentiation varied from 0.356 between E5 and E6 to 0.697 between E3 and E5. Gene flow among the populations was 0.261. Pair-wise analysis of the populations for their genetic differentiation and gene flow showed that the highest gene flow (0.904) was between E5 and E6 and lowest (0.217) was between E3 and E5 (Table 5). E3 showed an average gene flow estimate of 0.257 with all other populations.

Table 5. Pair-wise estimation of genetic differentiation and gene flow among six populations of *Samia cynthia ricini*

Ecoraces	Ht	Hs	Gst	Nm
E1x E2	0.1997	0.1087	0.4558	0.5969
E1x E3	0.2122	0.0795	0.6252	0.2998
E1x E4	0.1766	0.1031	0.4162	0.7013
E1x E5	0.1342	0.0845	0.3706	0.8493
E1x E6	0.1367	0.0801	0.4142	0.7072
E2x E3	0.1904	0.0802	0.5789	0.3637
E2x E4	0.1955	0.1037	0.4693	0.5655
E2x E5	0.193	0.0851	0.5589	0.3947
E2x E6	0.1947	0.0807	0.5854	0.3541
E3x E4	0.2227	0.0746	0.665	0.2519
E3 x E5	0.1848	0.056	0.6971	0.2172
E3x E6	0.1554	0.0516	0.668	0.2484
E4 x E5	0.1412	0.0796	0.4366	0.6453
E4x E6	0.1552	0.0752	0.5158	0.4694
E5x E6	0.0878	0.0565	0.3563	0.9035
Total	0.2334	0.0799	0.6575	0.2605

Genetic diversity and geographical distribution

In north-eastern India, *S. cynthia ricini* was distributed in the Brahmaputra river valley of Assam and in the hilly area. Titabar, where the population E3 was collected from, is located in the Upper Assam area and is approximately 421 kilo meters away from all other places of collection. The distance between the other populations was comparatively smaller. Regression of the genetic distance between the populations against geographic distance between them showed a positive correlation ($R^2 = 0.502$) (Figure 5).

Discussion

Recent advances in silkworm genome analysis showed the presence of large numbers of repeats in

the domesticated mulberry silkworm *Bombyx mori* (Mita et al. 2004; Biology Analysis group et al. 2004; Prasad et al. 2005). *S. cynthia ricini* seemed to share about 13% of its microsatellites with *B. mori* (Prasad et al. 2005). The presence of 250 mariner-like elements was reported in *Philosamia cynthia ricini* and were found to be similar to the *MosI* element from *Drosophila mauritiana* (Prasad et al. 2002). Genetic divergence among closely related populations of *B. mori* and in *A. mylitta* was successfully deciphered using ISSR profiling (Chatterjee et al. 2004; Pradeep et al. 2005; Kar et al. 2005). These studies explicitly report the importance of repeats and transposable elements in the molecular phylogenetic analysis of domesticated and wild silkworms. The higher amount of genetic divergence realized in this study




Figure 5. Regression of genetic distance between the populations of eri silkworm, *S.c. ricini* against geographical distance between their places of collection.

from closely related populations of *S. cynthia ricini*, further demonstrates the efficacy of the ISSR marker system. The ISSR markers revealed different levels of genetic differentiation among the populations of *S. cynthia ricini*. However, when a dominant marker system such as ISSR is used and the PCR products are resolved on agarose gels, a homozygote for a null allele at a particular locus will not produce a band, but both a heterozygote and a homozygote at that locus will produce a band. As a result, the fundamental data available from a population at any given locus are the number of individuals with a band corresponding to that locus and the number of individuals lacking that band. Nevertheless, ISSR markers showed the potential not only to reveal variations among geographically separated populations, but also among individuals within populations.

The genetic diversity revealed in this study is of much use in formulating strategies to conserve the genetic diversity present in this unique silkworm species. The low genetic variation revealed by AMOVA (ca. 13%) within the individuals of a population points to the fact that the populations are homogenous in nature, whereas the higher genetic variability (ca. 87%) among the populations indicated that the populations have already differentiated into separate genetic pools. Hence, these different gene pools should be conserved separately and maintained without any inter-mixing. Comparatively higher polymorphism (31.52%), heterozygosity (0.1093) and high boots trap values (98%) at the subgroup level observed in the E2 population points to significantly higher genetic differentiation at Borduar. Since moths of *S. cynthia ricini* can fly only a maximum of two kilometers, the chances of natural genetic mixing between the geographically separated populations can be ruled out. Transportation of populations for multiplication and unscientific breeding between *S. cynthia ricini* populations leads to genetic mixing of the populations. Such human interventions should be avoided at the core germplasm level to maintain purity of the germplasm.

The high gene flow (0.9035) between the populations E5 and E6 is quite understandable as they are similar in phenotypic traits, such as cocoon color (greenish blue). The high phenotypic and genetic similarity and gene flow observed within these two populations suggests its common origin and later succession into different populations by adapting to the varying climatic conditions. Since the E3 (Titabar) population is genetically and

geographically distanced from all other populations and seems to be an isolated one, detailed molecular analysis of this population is required to decipher its origin. The E3 population exhibited comparatively higher G_{ST} (0.657) combined with low gene flow ($N_m = 0.2762$) that has contributed significantly to genetic differentiation between populations (Wright 1978; Hartl and Clark 1997). The genetic distance of E3 from all other populations was found to be positively correlated with the geographic distance between Titabar and the natural habitats of other populations. This confirms an independent and slow isolated genetic differentiation of E3 at the high altitude of upper Assam where within population heterozygosity and gene flow was found to be low, and the possibility of genetic mixing seemed negligible. The independent genetic differentiation of this population is attributed to the strong effect of isolation by geographical distance. In fact, geographical isolation, mutation and selection are the most likely forces of population differentiation. This kind of association between geographic components and genetic differentiation was observed in natural populations of *Drosophila* (Frydenberg et al. 2003; Anderson et al. 2005). *S. cynthia ricini* silkmoths are poor flyers with a short life span, which is likely to reduce chances of genetic mixing among geographically close populations. This could account for the genetic differentiation among the geographically close *S. cynthia ricini* populations of Lower Assam. It is a known fact that high gene flow between populations precludes local adaptation that results from fixation of alleles favoured by local climatic conditions, and will prevent speciation (Barton and Hewitt 1985). At the same time gene flow generates new polymorphisms and new gene combinations in populations, upon which selection can act.

Thus from this study it is clear that the populations evaluated not only differ in their phenotypic traits but also in their genetic make up. Since *S. cynthia ricini* is believed to have originated in the Brahmaputra river valley of north-eastern India and this is the region where maximum genetic diversity is expected, the populations identified as genetically divergent have to be conserved *in situ*. The high economic value of the silk industry further emphasizes the need for urgent measures in this direction.

Acknowledgements

The authors are thankful to the anonymous reviewers for valuable criticism and to Central Silk Board, Government of India, Bangalore for financial support. The authors express deep regards to Professor Henry Hagedorn for his critical reading of the manuscript.

References

- Abbot P. 2001. Individual and population variation in invertebrates revealed by inter-simple sequence repeats (ISSRs). *Journal of Insect Science* 1.8 available online: insectscience.org/1.8
- Anderson AR, Hoffmann AA, McKechnie SW, Umina PA, Weeks AR. 2005. The latitudinal cline in the In(3R) Payne inversion polymorphism has shifted in the last 20 years in Australian *Drosophila melanogaster* populations. *Molecular Ecology* 14: 851-858.
- Barton NH, Hewitt GM. 1985. Analysis of hybrid zones. *Annual Review of Ecology and Systematics* 16: 113-148.
- Biology analysis group 2004. A draft Sequence for the genome of the domesticated silkworm (*Bombyx mori*). *Science* 306: 1937-1940.
- Chatterjee SN, Pradeep AR. 2003. Molecular markers (RAPD) associated with growth, yield and origin of the silkworm, *Bombyx mori* in India. *Russian Journal of Genetics* 39: 1612-1624.
- Chatterjee SN, Vijayan K, Roy GC, Nair CV. 2004. ISSR profiling of genetic variability in the ecotypes of *Antheraea mylitta* Drury, the tropical tasar silkworm. *Russian Journal of Genetics* 40: 152-159.
- Chowdhury SN. 1983. *Eri silk Industry*, Directorate of Sericulture & Weaving, Government of Assam, India.
- Ciolfi S, de Filippis T, Torti C, Malacrida AR, Dallai R. 2005. Molecular characterization and chromosomal localization of female-specific genes from the Mediterranean fruit fly *Ceratitis capitata* (Diptera: Tephritidae). *Genome* 48: 139-144.
- Deshpande KU, Apte GS, Bahulikar RA, Lagu MD, Kulkarni BO, Suresh HS, Singh NP, Rao MK, Gupta VS, Pant A, Renjekar PK. 2001. Genetic diversity across natural populations of montane plant species from the Western Ghats, India revealed by inter-simple sequence repeats. *Molecular Ecology* 10: 2397-2408.
- Estoup A, Solignac M, Harry M, Cornuet JM. 1993. Characterization of (GT)_n and (CT)_n microsatellites in two insect species: *Apis mellifera* and *Bombus terrestris*. *Nucleic Acids Research* 21: 1427-1431.
- Ehtesham NZ, Bentur JS, Bennett J. 1995. Characterization of a DNA sequence that detects repetitive DNA elements in the Asian gall midge (*Orseolia oryzae*) genome: Potential use in DNA fingerprinting of biotypes. *Gene* 153: 179-183.
- Frydenberg J, Hoffmann AA, Loeschcke V. 2003. DNA sequence variation and latitudinal associations in hsp23, hsp26 and hsp27 from natural populations of *Drosophila melanogaster*. *Molecular Ecology* 12: 2025-32.
- Goldstein DB, Schlötterer C. 1999. *Microsatellites: Evolution and Applications*. Oxford University Press, Oxford.
- Hartl DL, Clark AG. 1997. *Principles of population genetics*, 3rd edition. Sinauer Associates, Inc.
- Ichimura S, Mita K. 1992. Essential role of duplications of short motif sequences in the genomic evolution of *Bombyx mori*. *Journal of Molecular Evolution* 35: 123-130.
- Jolly MS, Sen SK, Sonwalker TN, Prasad GK. 1979. Non-mulberry silks. In: *Manual on Sericulture*, G Rangaswami, MN, Narasimhanna, K, Kashivishwanathan, CR, Sastri, MS Jolly, editors. , pp.1-178. Food and Agriculture Organization of the United Nations, Rome.
- Kar PK, Vijayan K, Nair CV, Mohandas TP, Saratchandra B, Thangavelu K. 2005. Genetic variability and genetic structure of wild and semi-domestic populations of tasar silkworm (*Antheraea mylitta*) ecorace Daba as revealed through ISSR markers. *Genetica* 125: 173-183.
- Katiyar SK, Chandel G, Tan Y, Zhang Y, Huang B, Nugaliyaddar L, Fernando K, Bentur JS, Inthavong S, Constantino S, Bennett J. 2000. Biodiversity of Asian rice gall midge (*Orseolia oryzae* Wood Mason) from five countries examined by AFLP analysis. *Genome* 43: 322-332.
- Kimpton CP, Gill P, Walton A, Urquhart A, Millican ES, Adams M. 1993. Automated DNA profiling employing multiplex amplification of short tandem repeat loci. *PCR methods and applications* 3: 13-22.
- Mita K, Kasahara M, Sasaki S, Nagayasu Y, Yamada T. 2004. The genome sequence of silkworm, *Bombyx mori*. *DNA Research* 11: 27-35.
- Nagaraja GM, Nagaraju J. 1995. Genome fingerprinting in silkworm, *Bombyx mori*, using random arbitrary primers. *Electrophoresis* 16: 1633-1638.
- Nei M. 1972. Genetic distance between populations. *American Naturalist* 106: 283-292.
- Nei M. 1973. Analysis of gene diversity in subdivided populations. *Proceedings of National Academy of Sciences, USA* 70: 3321-3323.
- Neupane FP, Thapa RB, Parajulee MN. 1990. Life and seasonal histories of the eri silkworm, *Samia cynthia ricini* Hutt. (Lepidoptera: Saturniidae), in Chitwan, Nepal. *Journal of Institute of Agriculture: Animal Science* 11: 113-120.
- Peigler RS. 1993. Wild silks of the world. *American Entomologist* 39: 151-161.
- Pradeep AR, Chatterjee SN, Nair CV. 2005. Genetic differentiation induced by selection in an inbred population of the silkworm *Bombyx mori*, revealed by RAPD and ISSR marker systems. *Journal of Applied Genetics* 46: 291-298.

- Prasad MD, Nurminsky DL, Nagaraju J. 2002. Characterization and molecular phylogenetic analysis of mariner elements from wild and domesticated species of silkmoids. *Molecular Phylogenetics and Evolution* 25 : 210-217.
- Prasad MD, Muthulakshmi M, Madhu M, Archak S, Mita K, Nagaraju J. 2005. Survey and analysis of microsatellites in the silkworm, *Bombyx mori*: Frequency, distribution, mutation, marker potential and their conservation in heterologous species. *Genetics* 169: 197-214.
- Reddy KD, Nagaraju J, Abraham EG. 1999a. Genetic characterization of silkworm *Bombyx mori* by simple sequence repeats (SSR) – anchored PCR. *Heredity* 83: 681-687.
- Reddy KD, Abraham EG, Nagaraju J. 1999b. Microsatellites of the silkworm, *Bombyx mori*: abundance, polymorphism and strain characterization. *Genome* 42: 1057-1065.
- Reineke A, Karlovsky P, Zebitz CPW. 1998. Preparation and purification of DNA from insects for AFLP analysis. *Insect Molecular Biology* 7: 95-99.
- Siddiqui AA, Saha LM, Das PK. 2000. Genetic variability and correlation studies of some quantitative traits in Eri silkworm. *International Journal of wild silkworm and silk* 5: 234-237.
- Singh KC, Benchamin KV. 2002. Biology and ecology of the eri silkworm *Samia ricini* (Donovan) (Saturniidae): a review. *Bulletin of Indian Academy of Sericulture* 6: 20 -33.
- Sneath PH, Sokal RR. 1973. *Numerical taxonomy*. W. H. Freeman, San Francisco.
- Suryanarayana N, Sarmah MC, Sahu AK, Kumar A, Das PK. 2002. Status paper on promotion of eri culture in India. *Proceedings of National Workshop on Sericulture Germplasm Management & Utilization*, pp. 50–53, CSGRC, Hosur, India.
- Suzuki Y, Gage L, Brown DD. 1972. The genes for silk fibroin in *Bombyx mori*. *Journal of Molecular Biology* 70: 637-649.
- Swofford DL. 1998. *PAUP*: Phylogenetic analysis using parsimony* (*and other Methods), Version 4. Sinauer Associates, Sunderland, MA.
- Thangavelu K. 1991. Wild sericigenous insects of India: A need for conservation. In: *Wild Silkmoids'91*. 71–77. International Society for wild silkmoids: Japan.
- Vlachou D, Konsolaki M, Tolias PP, Kafatos FC, Komitopoulou K. 1997. The autosomal chorion locus of the medfly, *Ceratitis capitata*. I. Conserved synten, amplification and tissue specificity but sequence divergence and altered temporal regulation. *Genetics* 147: 1829-1842.
- Vogel JM, Scolnik PA. 1997. Direct amplification from microsatellites: detection of simple sequence repeat-based polymorphisms without cloning. In: Caetano-Anolles G, Gresshoff PM, editors. *DNA markers Protocols, Applications, and Overviews*, 133–150. Wiley-VCH: New York.
- Vos P, Hogers R, Bleeker M, Reijans M, van de Lee T, Hornes M, Frijters A, Pot J, Peleman J, Kuiper M, Zabeau M. 1995. AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Research* 23: 4407-4414.
- Wright S .1978. Evolution and Genetics of population variability within and among natural populations. The University of Chicago press, Chicago.
- Williams JGK, Kubelik AR, Livak KJ, Rafalski JA, Tingey SV. 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Research* 18: 6531-6535.
- Yeh FC. 1998. *POPGENE16*. Version 1.31. (<http://www.ualberta.ca/~fyeh>) Agriculture and Forestry Molecular Biology and Biotechnology center, University of Alberta and Center for International Forestry Research, Canada.
- Zietkiewicz E, Rafalski A, Labuda D. 1994. Genome fingerprinting by simple sequence repeat (SSR) - anchored polymerase chain reaction amplification. *Genomics* 20: 176-183.