

In situ preserved equisetophyte stems in the Upper Bajocian Parkinsoni Zone (Middle Jurassic) of the Northern Caucasus (Russia)

Authors: Naugolnykh, Serge V., and Mitta, Vasily V.

Source: Palaeodiversity, 9(1) : 113-120

Published By: Stuttgart State Museum of Natural History

URL: <https://doi.org/10.18476/pale.v9.a8>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

In situ preserved equisetophyte stems in the Upper Bajocian Parkinsoni Zone (Middle Jurassic) of the Northern Caucasus (Russia)

SERGE V. NAUGOLNYKH AND VASILY V. MITTA

Abstract

This paper focuses on the first record of equisetophyte stems preserved *in situ* in the dark siltstones of the Djangura Formation of the Late Bajocian (Middle Jurassic) age in the Bolshoi Zelenchuk River Basin (Zelenchuk district of the Karachaevo-Cherkessk Republic, Northern Caucasus, Russia). The stems are long and narrow, with long internodes, vertically orientated in growth upright position. The outer surface of the stems bears thin, prolonged ribs. There are well-formed diaphragms at the stem nodes. The diaphragms are covered with fine radial striations. The stems possess a central internal cavity of subcylindrical shape. The sedimentary conditions of forming the layer with equisetophyte stems can be interpreted as tidal plain or littoral.

Key words: Middle Jurassic, equisetophytes, Equisetaceae, *Neocalamites*, *Equisetites*, Northern Caucasus, Upper Bajocian, palaeoecology.

1. Introduction

The palaeoecological significance of fossil equisetophytes is related to their stable environmental preferences, allowing the use of these plants as markers of certain palaeoenvironments. Most of the ancient and recent equisetophytes were and still are inhabitants of near-shore humid environments. Moreover, many equisetophytes can be regarded as pioneering plants which are adapted to fast colonization of clastic substrates under stressed ecological conditions. This is why the equisetophytes s. l. can be considered as a useful tool for general palaeoecological reconstructions, especially if these plants are preserved *in situ* (TAYLOR et al. 2009; THOMAS 2014).

This paper is devoted to the first record of equisetophytes preserved *in situ* in the Middle Jurassic Djangura Formation (BESNOV 1967) of the Northern Caucasus. The exact stratigraphic position of this find is based on the rich marine fauna (including those found at the same level). This observation is helpful for a palaeogeographic reconstruction of that region for the Late Bajocian.

Acknowledgements

The authors express their deep gratitude to the reviewers MIKHAI POPA (University of Bucharest, Faculty of Geology and Geophysics) and AGATA JARZYŃKA (W. Szafer Institute of Botany, Polish Academy of Sciences) for their helpful and constructive remarks and advices. This work was funded by the subsidy of the Russian Government to support the Program of Competitive Growth of Kazan Federal University among World's Leading Academic Centers and the State Program no. 01201459177 of the Geological Institute of Russian Academy of Sciences, and Program no. 28 of Presidium of the Russian Academy of Sciences.

2. Material

The material described was collected by one of the present authors (VM) during the field seasons in 2015–2016, which were aimed for studying of the Upper Member of the Djangura Formation (BESNOV 1967) outcropping in the Kuban River Basin (Zelenchuk District of the Karachaevo-Cherkessk Republic of Russia, Northern Caucasus; Fig. 1). This member is represented by clastic terrigenous deposits, e.g., clays, siltstones, alevrolites, with siderite concretions forming layer-like aggregations. All layers strike at 5–7° towards North to North-East. The general sedimentary environment was related to a deep to shallow shelf with low hydrodynamic conditions (BESNOV & MITTA 1993).

The specimens described and discussed in the present paper come from the locality No. 25. The section is disposed on the left bank of the Kyafar River, 1.8 km upstream from its mouth (Figs. 1, 2). The length of the section is ca. 150 m. The dense grey and dark-grey clays and siltstones with siderite concretions, 5–7 cm in diameter, containing ammonites: *Parkinsonia* ex gr. *djanelidzei* KAKHADZE, a single *Rarecostites donezianus* (BORISSJAK), and numerous *Dinolytoceras zhivagoi* BESNOV. The ammonites most commonly are represented by fragmentary preserved casts. Small belemnite rostra (*Megateuthis* sp.) and rare bivalves (*Entolium* sp.) were also found in this section. The subvertically preserved stems of equisetophytes have been recorded in the upper part of the section, one meter above the level with ammonites. The general thickness of the deposits outcropping in the section is more than five meters.




Fig. 1. Geographical position of locality No. 25 (marked by an asterisk). **A:** General map; **B:** detailed map.

Without any doubt the clays with the equisetophyte macrofossils can be assigned to the *Parkinsonia parkinsoni* Chronozone of the Late Bajocian. Older deposits of Bajocian age crop out along both banks of the Kyafar River upstream of the locality No. 25. The equivalents of the standard subdivisions of the Upper Substage, e.g. the lower subzone of the *Parkinsonia parkinsoni* Zone and the *Garantiana garantiana* and *Strenoceras niortense* zones occur upstream, up to the Storozhevaya village (MITTA & SHERSTYUKOV 2014; MITTA & SCHWEIGERT 2016). The Lower Bajocian outcrops upstream of the Storozhevaya village on the Kyafar River banks and its incomings just above the water level. The Lower Bathonian rocks, which are time-equivalent to the *Zigzagiceras zigzag* Chronozone, can be observed downstream of the Kyafar River mouth along the banks of the Bolshoy Zelenchuk River (MITTA 2015).

The deposits observed in the section No. 25 belong to the middle part of the *Parkinsoni* Zone and are described as the Beds with *Parkinsonia djanelidzei* (MITTA et al. 2017, in press), probably corresponding to the lowest part the Truellei Subzone of the Standard Time Scale).

The material studied is stored in the Geological Institute of Russian Academy of Sciences (GIN RAS), Moscow, under the collection numbers 4914/29–4914/33.

3. Observations

The material includes two well-preserved stems (Figs. 3A–C, 4A, B, E), both with basal parts, and seven fragments of stems preserved in different extent. The best preserved stem fragment is 94 mm long and 17 mm wide.




Fig. 2. Section of locality No. 25. Middle Jurassic, Upper Bajocian; the middle part of the *Parkinsonia parkinsoni* Zone, Beds with *Parkinsonia djanelidzei*. Karachaevo-Cherkessk Republic, Russia; Zelenchuk District, Kyafar River. Legend: 1 – clays and siltstones; 2 – concretions; 3 – ammonite shells; 4 – belemnite rostra; 5 – equisetophyte stems, preserved *in situ*.

The second well-preserved fragment is 87 mm long and 11 to 20 mm wide. The widest part of the stem occurs towards its upper (distal) part, so the stem width is gradually decreasing towards the stem base. Judging from the general shape of the stem fragments we can suppose that the whole length of the plant was not less than 300 mm.

The shape of the stems is more or less cylindrical, with conical to rounded basal parts (Fig. 4D). The superficial surface of the stems is rather smooth, but actually it bears very fine prolonged ribs (Fig. 4D, E). The nodes are very indistinct. Only one stem fragment shows two nodes and one complete internode (specimen GIN 4914/30). The length of the internode is 89 mm, and the width of the prolonged ribs is 0.3 mm. There is one unclear evidence (weakly preserved lateral branch scar) that the main vertical stem had lateral branches of smaller size, which were attached to the node line along the main axis.

Two fragments of stem, which are broken in nodes, clearly show nodal diaphragms typical of all representatives of equisetophytes. The diaphragms bear a fine radial striation (Fig. 4A, B). All the specimens in the collection have a distinct inner cavity, cylindrical in shape (Fig. 4C, E), which is also characteristic for all representatives of equisetophytes. The cavity is filled with secondary minerals (mostly calcite).

The field observations (Fig. 3C) showed that the stems were attached to subhorizontal stolons or rhizomes, but these rhizomes are represented by relatively short fragments. This can be explained by the horizontal rhizomes as results of a fragmentation process, which is typical of the vegetative reproduction mode of equisetophytes. One lateral branch, which has been deformed, is observed in the basal part of the stem (Fig. 3A, B).

Although the features of the fossils are relatively poor, the general habit and the morphological details allow us to suggest that these remains belonged to an equisetophyte closely related to the genera *Equisetites* STERNBERG or *Neocalamites* HALLE, and on this basis the Kyafar plants can be preliminarily attributed to the family Equisetaceae.

Concerning the *in situ* preservation of the stems and rhizomes of the Kyafar equisetophytes, it is obvious that these plants started their life on the bottom of a very shallow basin near zero depth or on the littoral, but after that they were submerged and buried by slow but permanent sedimentation of clastic material (Fig. 5, S).

4. Discussion

Stems of equisetophytes preserved *in situ* in palaeo-soils or FPS-profiles (abbreviation made after “fossil-palaeo-soil” term) as well as in the more or less common sediments, are recorded from different deposits, mostly representing near-shore or shallow-water environments (e.g., WATSON & BATTEN 1990; KELBER & VAN KONIJNENBURG-VAN CITTERT 1998; TVERDOKHLEBOV 2004; APPLETON et al. 2011; THOMAS 2014).

A specific feature of the Kyafar equisetophytes is their very long and relatively narrow internode, but this morphological character is also reported for some other equisetophytes beginning from the Permian up to the Recent. For instance, similar exceptionally long internodes are typical of *Neocalamites tubulatus* NAUGOLNYKH from the Upper Permian of the northern part of Russia (NAUGOLNYKH 2009, figs. 3, 5). The stems of the Kyafar equisetophytes have many features in common with some species of the genus *Equisetites* STERNBERG (*sensu lato*) from the Middle Jurassic of Turkmenistan. For example, the Kyafar equisetophytes are similar to *Equisetites ketovae* BURAKOVA in having very long internodes and a smooth to finely ribbed surface of the stems (BURAKOVA 1960, pl. 14, fig. 1), and they are especially similar to underground organs (stolones or rhizomes) of this species (BURAKOVA 1960, fig. 3), which were misinterpreted by that author as “roots” (BURAKOVA 1960). Very fine ribs as well as long internodes and the shape of the node diaphragms of the Kyafar equisetophytes are similar to the Middle Triassic equisetophyte described as *Zonulamites nymboidensis* HOLMES from Australia (HOLMES 2000; especially specimens illustrated on




Fig. 3. A–C: The equisetophyte stems preserved *in situ*, field photographs. Scale bars equal 10 mm. Abbreviation: HS – horizontal stolon (rhizome). The node is marked by arrows in Fig. 3B.




Fig. 4. The equisetophyte stems preserved *in situ*. **A, B:** Nodal part with the well-discernible diaphragm, specimen GIN RAS No. 4914/30; **C:** cross section with the distinct central cavity filled with secondary minerals, specimen GIN RAS No. 4914/33; **D:** basal part of rounded shape, specimen GIN RAS No. 4914/32; **E:** longitudinal ribs covering outer surface of the stem, specimen GIN RAS No. 4914/30. Scale bars equal 10 mm.




Fig. 5. Morphological interpretation of the fossils studied (see Figs. 3, 4) and the environmental conditions of growing of the Kyafar equisetophytes. Scale bar equals 10 mm. Abbreviations: BS – branch scar; R – rhizome; S – direction of sedimentation.

his figure 6A, C, E, F), which, according to our viewpoint, can be assigned to the genus *Equisetites* s. l. as well.

The evolutionary history of the equisetophytes s. str. (e.g., the family Equisetaceae) now can be traced back as deep as to the Carboniferous and Permian (DiMICHELE et al. 2005; NAUGOLNYKH 2009; YANG et al. 2011; SINGH et al. 2011). The direct phylogenetic predecessors of Equisetaceae are representatives of the family Tchernoviaceae, which is characteristic for the Carboniferous and Permian of Angaraland (MEYEN 1971; MEYEN & MENSHIKOVA 1983; NAUGOLNYKH 2002, 2004). Equisetophytes s. l. closely related to Tchernoviaceae are also recorded from the Permian of the Southern Hemisphere, e.g., Gondwanaland (HOLMES 1996; DURAN et al. 1997; ESCAPA & CUNEO 2005; CUNEO & ESCAPA 2006).

During their evolution equisetophytes show a strict adaptation for growing in near-shore environments (TAYLOR et al. 2009). The new find of *in situ* stems of equisetophytes in the Middle Jurassic of the Kyafar locality is in good agreement with that tendency and strongly supports the possibility of using equisetophytes as important ecological markers of wet environments.

In evolutionary terms, the genus *Equisetum* was in stasis at least from the Paleocene (McIVER & BASINGER 1989), or even from earlier times (GOULD 1968; WATSON & BATTEN 1990; MARKOVICH 1993). Most probably this stasis was corresponding with constant ecological preferences of the equisetophytes as more or less permanent hydro- or hygrophils.

5. References

- APPLETON, P., MALPAS, J., THOMAS, B. A. & CLEAL, C. J. (2011): The Brymbo Fossil Forest. – *Geology Today*, **27**: 107–113.
- BESNOSOV, N.V. (1967): Bajocian and Bathonian deposits of Northern Caucasus. – 179 pp.; Moscow (Nedra).
- BESNOSOV, N. V. & MITTA, V. V. (1993): Late Bajocian – Bathonian ammonites from Northern Caucasus and Central Asia. – 347 pp.; Moscow (Nedra). [In Russian]
- BURAKOVA, A. T. (1960): New species of Equisetales from the Middle Jurassic of the Western Turkmenistan. – *Paleontological Journal*, **2**: 149–157.
- CUNEO, N. R. & ESCAPA, I. (2006): The equisetalean genus *Cruciatheca* nov. from the Lower Permian of Patagonia, Argentina. – *International Journal of Plant Sciences*, **167** (1): 167–177.
- DiMICHELE, W. A., VAN KONIJNENBURG-VAN CITTERT, J. H. A., LOOY, C. V. & CHANEY, D. S. (2005): *Equisetites* from the Early Permian of North-Central Texas – The Nonmarine Permian. – *Bulletin of the New Mexico Museum of Natural History and Science*, **30**: 56–59.
- DURAN, M., HÜNICKEN, M. A. & ANTON, A. M. (1997): Nove-dos hallazgos de Sphenopsida en la formacion Bajo de Veliz, provincia de San Luis, Argentina. – *Ameghiniana*, **34** (3): 259–264.
- ESCAPA, I. & CUNEO, R. (2005): A new equisetalean plant from the Early Permian of Patagonia, Argentina. – *Review of Palaeobotany and Palynology*, **137** (1–2): 1–14.
- GOULD, R. E. (1968): Morphology of *Equisetum laterale* PHILLIPS, 1829, and *E. bryanii* sp. nov. from the Mesozoic of South-eastern Queensland. – *Australian Journal of Botany*, **16**: 153–176.
- HOLMES, W. B. K. (1996): A fossil plant organ with unusual internal structure from the Late Carboniferous of New South Wales. – *Alcheringa*, **20**: 69–72.
- HOLMES, W. B. K. (2000): The Middle Triassic megafossil flora of the Basin Creek Formation, Nymboida Coal Measures, NSW, Australia. Part 1: Bryophyta, Sphenophyta. – *Proceedings of the Linnean Society of New South Wales*, **122**: 43–68.
- KELBER, K. P. & VAN KONIJNENBURG-VAN CITTERT, J. H. A. (1998): *Equisetites arenaceus* from the Upper Triassic of Germany with evidence for reproductive strategies. – *Review of Palaeobotany and Palynology*, **100** (1): 1–26.
- MARKOVICH, E. V. (1993): Jurassic and Early Cretaceous Equisetales from the southeast of the Siberian Platform. – *Paleontological Journal*, **1**: 110–119.
- McIVER, E. E. & BASINGER, J. F. (1989): The morphology and relationships of *Equisetum fluviatoides* sp. nov. from the Paleocene Ravenscrag Formation of Saskatchewan, Canada. – *Canadian Journal of Botany*, **67**: 2937–2943.
- MEYEN, S. V. (1971): *Phyllothea*-like plants from the Upper Palaeozoic flora of Angaraland. – *Palaeontographica*, (B), **133**: 1–33.
- MEYEN, S. V. & MENSHIKOVA, L. V. (1983): Systematics of the Upper Palaeozoic articulates of the family Tchernoviaceae. – *Botanical Journal*, **68**: 721–729. [In Russian]
- MITTA, V. V. (2015): Genus *Oraniceras* (Parkinsoniidae, Ammonoidea) from the Lower Bathonian of Southern European Russia. – *Palaeontologicheskii Zhurnal*, **6**: 38–42 [in Russian]. English Translation: *Paleontological Journal*, **49** (6): 595–601.
- MITTA, V. V. & SCHWEIGERT, G. (2016): A new morphotype of lower jaw associated with *Calliphylloceras* (Cephalopoda: Ammonoidea) from the Middle Jurassic of the Northern Caucasus. – *Paläontologische Zeitschrift*, **90**: 293–297.
- MITTA, V. V. & SHERSTYUKOV, M. P. (2014): On the Bajocian and Bathonian of Bolshoy Zelenchuk River basin (Northern Caucasus). – In: IVANOV, A. V. (ed.): *Problems of paleoecology and historical geoecology*, 74–81; Saratov (Saratov State Technical University).
- MITTA, V. V., SAVELIEVA, J. N., FEODOROVA, A. A. & SHUREKOVA, O. V. (2017, in press): Biostratigraphy of the Bajocian-Bathonian boundary beds of Bolshoy Zelenchuk River basin (Northern Caucasus). – *Stratigraphy and Geological Correlation*.
- NAUGOLNYKH, S. V. (2002): *Paracalamitina striata* – a newly reconstructed equisetophyte from the Permian of Angaraland. – *Journal of Paleontology*, **76** (2): 377–385.
- NAUGOLNYKH, S. V. (2004): On some aberrations of extant horse-tails (*Equisetum* L.) and the origin of the family Equisetaceae. – *Paleontological Journal*, **38** (3): 335–342.
- NAUGOLNYKH, S. V. (2009): A new fertile *Neocalamites* from the Upper Permian of Russia and equisetophyte evolution. – *Geobios*, **42**: 513–523.
- SINGH, K. J., CHANDRA, S. & SAXENA, A. (2011): *Tatapania* gen. nov., a possible cone of *Schizoneura gondwanensis* FEISTMANTEL from the Late Permian in the Tatapani-Ramkola Coalfield, India. – *The Palaeobotanist*, **60**: 251–263.
- TAYLOR T. N., TAYLOR E. L. & KRINGS M. (2009): *Paleobotany*. – 1230 pp.; Amsterdam & Boston (Elsevier).

- THOMAS, B. A. (2014): *In situ* stems: preservation and growth habits of the Pennsylvanian (Carboniferous) Calamitaleans based upon new studies of *Calamites* STERNBERG 1820 in the Duckmantian at Brymbo, North Wales, UK. – *Palaeontology*, **57**: 21–36.
- TVERDOKHLEBOV, V. P. (2004): Buried *Equisetites* thickets from the Middle Triassic of the South Cis-Urals, Russia. – *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, **2004** (10): 595–604.
- WATSON, J. & BATTEN, D. J. (1990): A revision of the English Wealden Flora, II. Equisetales. – *Bulletin of British Museum of Natural History (Geology)*, **46** (1): 37–60.
- YANG TAO, NAUGOLNYKH, S. V. & SUN GE (2011): A new representative of *Neocalamites* HALLE from the Upper Permian of Northeastern China (Jiefangcun Formation). – *Paleontological Journal*, **45** (3): 335–346.

Addresses of the authors:

SERGE V. NAUGOLNYKH, Geological Institute, Russian Academy of Sciences, Pyzhevsky per. 7, 119017 Moscow, Russia; Contract affiliation: Kazan Federal University.

VASILY V. MITTA, Borissiak Paleontological Institute, Russian Academy of Sciences, Profsoyuznaya str. 123, 117997 Moscow, Russia.
E-mails: naugolnykh@rambler.ru; naugolnykh@list.ru; mitta@paleo.ru

Manuscript received: 17 July 2016, revised version accepted: 20 October 2016.