



Scat-Detection Dogs Seek Out New Locations of *Priodontes maximus* and *Myrmecophaga tridactyla* in Central Brazil

Authors: Vynne, Carly, Machado, Ricardo B., Marinho-Filho, Jader, and Wasser, Samuel K.

Source: *Edentata*, 2009(10) : 13-14

Published By: IUCN/SSC Anteater, Sloth and Armadillo Specialist Group

URL: <https://doi.org/10.1896/020.010.0114>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Scat-Detection Dogs Seek Out New Locations of *Priodontes maximus* and *Myrmecophaga tridactyla* in Central Brazil

Carly Vynne
Ricardo B. Machado
Jader Marinho-Filho
Samuel K. Wasser

The use of scat-detection dogs is increasingly recognized as a valuable wildlife assessment and monitoring tool (Long *et al.*, 2007a). Chosen for their drive for play-reward with a tennis ball, these dogs enable researchers to seek out scat samples of rare and otherwise difficult-to-study species. The dogs are able to cover large areas, are unbiased in their sampling of gender, and have demonstrated accuracy in their ability to home in on their targets while ignoring non-target species (Smith *et al.*, 2003). Studies comparing detection dogs with camera-traps and hairsnag surveys have demonstrated that detection dogs are superior both at locating the presence of target species as well as number of individuals (Wasser *et al.*, 2004; Harrison, 2006; Long *et al.*, 2007b). Once located, the scat samples may be used to understand wildlife movements, for diet and disease studies, and for DNA and hormone analyses (Wasser *et al.*, 2004).

In 2004 we employed three teams of scat-detection dogs in a successful pilot study to survey carnivores (maned wolf *Chrysocyon brachyurus*, jaguar *Panthera onca*, and puma *Puma concolor*) in the Cerrado of central Brazil (Vynne *et al.*, 2005). Given this success, we decided to try training these seasoned dogs to also find two species of xenarthrans, *Priodontes maximus* and *Myrmecophaga tridactyla*. Subsequent field seasons, beginning in May 2006, have yielded valuable information on the occurrence and habitat use of these two species (Vynne *et al.*, in prep.; Silveira *et al.*, 2009) in and around Emas National Park, a 133,000-hectare reserve on the common border of the states of Goiás and Mato Grosso do Sul. To our knowledge, this is the first study using scat-detection dogs outside of North America, and the first to survey for species in the Xenarthra.

Results thus far include 67 point localities of giant armadillo and 560 of giant anteater in and around Emas National Park. These locations have been the first to show extensive use of non-Park areas by the two species and to demonstrate how the two species are using the system of legal reserves and agricultural lands. Giant anteater samples are being used to monitor changes in the density of the population, which

has been susceptible to large-scale fires in the Park, and giant armadillo samples will be analyzed to learn how these localities correspond to movement and home range.

Of particular importance has been that both species are still found to occur outside of Emas National Park. While Emas is a grassland island, almost entirely surrounded by intensive agriculture, current Federal legislation requires landowners to set aside 20–30% of their farm as protected land. Our initial assessment is that this mosaic of habitat fragments is extremely important for these two species, providing corridors for movement and critical protection to individuals and their food sources. Of concern, however, is the small amount of natural grasslands, the habitat preferred by the two species inside the Park, under protection beyond Park borders. As these grasslands support the highest density of ant and termite mounds, priority should be given to protecting and restoring natural, open habitats that support these critical food sources in the landscape surrounding Emas National Park.

Our field surveys concluded in May of 2008 and spatial analyses are underway. In the meantime, we are looking for collaborators who may be interested in using the physical samples that we have been collecting. Samples have been divided for DNA and diet/hormone/disease analyses and are stored in a 20% DMSO (dimethylsulfoxide) salt solution (a preservative for DNA) and frozen. Each sample has associated information on habitat, spatial location and sample quality that we would make available. The samples are currently in Brazil at the Universidade de Brasília and within-country requests will receive first priority, although it is possible to arrange for export permits if necessary. For more information, please contact Carly Vynne at <cvynne@u.washington.edu>.

This work is a partnership between the University of Washington, Conservation International, the University of Brasília and the Jaguar Conservation Fund; financial support was provided by the TEAM Network of Conservation International, funded by the Gordon and Betty Moore Foundation and the Brazil Program of Conservation International. Licensing for the project provided by IBAMA no. 02001.00215/07-21.

Carly Vynne and Samuel K. Wasser, Center for Conservation Biology, Department of Biology, University of Washington, Seattle, WA 98115-1800, USA, e-mail <cvynne@u.washington.edu> and <wassers@u.washington.edu>, Ricardo B. Machado, Department of Zoology, Institute, of Biological Science, University

of Brasilia, Campus Darcy Ribeiro, 70910-900, Brasilia DF, Brazil, e-mail:<rbmac@uol.com.br> and **Jader Soares Marinho-Filho**, Universidade de Brasilia, Instituto de Ciências Biológicas, Departamento de Zoologia, Campus UnB, Asa Norte, 70910900 - Brasilia, DF – Brazil, e-mail: <jmarinho@unb.br>.

References

- Harrison, R. L. 2006. A comparison of survey methods for detecting bobcats. *Wildl. Soc. Bull.* 34(2): 548–55.
- Long, R. A., Donovan, T. M., Mackay, P., Zielinski W. J. and Buzas, J. S. 2007a. Effectiveness of scat detection dogs for detecting forest carnivores. *J. Wildl. Manage.* 71: 2007–2017.
- Long, R. A., Donovan, T. M., Mackay, P., Zielinski, W. J. and Buzas, J. S. 2007b. Comparing scat detection dogs, cameras, and hair snares for surveying carnivores. *J. Wildl. Manage.* 71(6): 2018–2025.
- Silveira, L., Jácomo, A. T. A., Furtado, M. M., Torres, N. M., Sollmann, R., Vynne, C. 2009. Ecology of the giant armadillo (*Priodontes maximus*) in the grasslands of central Brazil. *Edentata* 8–10: 27–36.
- Smith, D. A., Ralls, K., Hurt, A., Adams, B., Parker, M., Davenport, B., Smith, M. C. and Maldonado, J. E. 2003. Detection and accuracy rates of dogs trained to find scats of San Joaquin kit foxes (*Vulpes macrotis mutica*). *Anim. Cons.* 6: 339–346.
- Vynne, C., Silveira, L., Groom, M., and Wasser, S. 2005. Matrix composition affects presence and abundance of maned wolf, puma, and jaguar in a Cerrado ecosystem. *19th Annual Meeting of the Society for Conservation Biology: Book of Abstracts*. Brasilia, Brazil. P. 221.
- Wasser, S. K., Davenport, B., Ramage, E. R., Hunt, K. E., Parker, M., Clarke, C. and Stenhouse, G. 2004. Scat detection dogs in wildlife research and management: Applications to grizzly and black bears in the Yellowhead Ecosystem, Alberta, Canada. *Can. J. Zool.* 82: 475–492.