

Hunting Strategy of the Margay (*Leopardus wiedii*) to Attract the Wild Pied Tamarin (*Saguinus bicolor*)

Authors: de Oliveira Calleia, Fabiano, Rohe, Fabio, and Gordo, Marcelo

Source: Neotropical Primates, 16(1) : 32-34

Published By: Conservation International

URL: <https://doi.org/10.1896/044.016.0107>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

HUNTING STRATEGY OF THE MARGAY (*LEOPARDUS WIEDII*) TO ATTRACT THE WILD PIED TAMARIN (*SAGUINUS BICOLOR*)

Fabiano de Oliveira Calleia
Fabio Rohe
Marcelo Gordo

Introduction

Wild primate predation has been widely reported for various Neotropical cat species. Apparently, prey body size and predator body size are related, with large cats preying mainly on large primates. Remains from different species of primates have been observed in scats of different Neotropical cat species (Table 1). According to Cabrera and Yepes (1940), primates are the favorite prey of *Puma yaguaroundi* in some regions of Central America. More recently, Miranda *et al.* (2005), found fingers and nails from *Alouatta guariba clamitans* in two fecal samples from *Leopardus pardalis* and suggests that the ocelot may be a potential predator of all Neotropical primates.

In this study, we focus on a hunting technique by the margay, *Leopardus wiedii*. Morphologically, margays have

arboreal adaptations, but there are no published reports of the predation strategy of wild margays. The few studies on the margay suggest that its diet is mainly composed of arboreal mammals. Mondolfi (1986) analyzed the stomach contents of margay from Venezuela and found remains of squirrel (*Sciurus granatensis*) and the wedge-capped capuchin monkey (*Cebus olivaceus*-cited as *nigrivittatus*, a junior synonym). Margay prey species in Guyana were also arboreal mammals (Beebe 1925). In captivity, margays were observed preying on *Saguinus niger* (Oliveira, 1998).

In the course of our field research on felids, we interviewed local Amazon jungle inhabitants (woodsmen and mestizo indians) in different regions of central Amazonia to learn about the biodiversity of local habitats, and in particular, the natural history of Neotropical cat species, including their prey capture techniques. Interestingly, several of the interviewees described a common predation strategy by Neotropical cats as attracting their prey by mimicking the prey species' vocalizations. More than a dozen reports of *Puma concolor*, *Panthera onca* and *Leopardus pardalis* mimicking vocalizations of agoutis (*Dasyprocta* spp.), tinamous or nambús (*Crypturellus* sp.) and solitary tinamous or macucos (*Tinamus* sp.) were made in different river basins (Madeira, Juruá and Purus) (Table 2). Until now, no scientific observations of this type of behavior have been published

Table 1. Review of primates predated by Neotropical cat species.

Predator	Prey	Location	Citation
<i>Panthera onca</i>	<i>Ateles belzebuth</i>	La Macarena, Colombia	Matsuda and Izawa (2008)
	<i>Alouatta seniculus</i>	Venezuela	Peetz <i>et al.</i> (1992)
	<i>Brachyteles arachnoides</i>	Intervales State Park, Southeast Brazil	Olmos (1994)
	<i>Ateles [paniscus] chamek</i>	Perú	Emmons (1987)
<i>Puma concolor</i>	<i>Ateles geoffroyi</i>	Corcovado National Park, Costa Rica	Chinchilla (1997)
	<i>Ateles geoffroyi</i>	Maya Biosphere Reserve, Guatemala	Novack <i>et al.</i> (2005)
	<i>Ateles Belzebuth</i>	La Macarena, Colombia	Matsuda and Izawa (2008)
	<i>Alouatta pigra</i>	Maya Biosphere Reserve, Guatemala	Novack <i>et al.</i> (2005)
	<i>Alouatta caraya</i>	Mutum Island, Southern Brazil	Ludwig <i>et al.</i> (2007)
	<i>Ateles chamek</i>	Perú	Emmons (1987)
<i>Leopardus pardalis</i>	<i>Saguinus</i> spp.		Goldizen (1987)
	<i>Saguinus nigricollis</i>	Colombia	Izawa (1978)
	<i>Alouatta guariba</i>	Caratinga Biological Station, Southeast Brazil	Bianchi & Mendes (2007)
	<i>Brachyteles hypoxanthus</i>	Caratinga Biological Station, Southeast Brazil	Bianchi & Mendes (2007)
	<i>Cebus apella nigritus</i>	Caratinga Biological Station, Southeast Brazil	Bianchi & Mendes (2007)
	<i>Alouatta g. clamitans</i>	Chácara Payquere, Southern Brazil	Miranda et al (2005)
	<i>Saguinus fuscicollis</i>	Perú	Emmons (1987)
	<i>Saimiri sciureus</i>	Perú	Emmons (1987)
<i>Puma yaguaroundi</i>	Primates		Cabrera & Yepes (1940)
	<i>Callithrix jacchus</i>	Paraíba State, Northeast Brazil	Ximenes (1982)
<i>Leopardus wiedii</i>	<i>Cebus olivaceus</i>	Venezuela	Mondolfi (1986)
	<i>Saguinus niger</i>	(in captivity)	Oliveira (1998)
	<i>Cebus apella</i>	British Guiana	Beebe (1925)

for Neotropical felids. Here we report the first field observation of margay mimicking behavior, recorded during field research on the primate pied tamarin (*Saguinus bicolor*) at the Reserva Florestal Adolpho Ducke (59 56' 15,71556" W, 02 56' 25,75037" S) in Manaus, Brazil (for a description of the area, see Ribeiro *et al.*, 1999). In this brief report we suggest that *L. wiedii* uses a mimicking strategy to capture its prey. Our record confirms the reliability of the information provided by the local Amazonian inhabitants.

On October 12, 2005, at 9:13 am, a group of eight pied tamarins monitored by telemetry was feeding in a Moraceae (*Ficus* sp.). A large vine at 15 meters height connected the surrounding trees to the fig tree. At 9:18 am, a margay attracted the attention of a tamarin sentinel (Gordo *et al.*, 2005) by producing calls similar to those emitted by pied tamarin pups. The adult male sentinel climbed up and down the tree to investigate the calls coming from behind the liana tangles. It assumed a surveillance position and, using specific calls, warned the group about the foreign calls. At 9:22 am we observed movements in the vine and keep hearing the call imitations. At 9:29 am three pied tamarin individuals were feeding on *Ficus* sp. while the tamarin sentinel was keeping surveillance. At 9:40 am, four pied tamarins climbed up and down the Moraceae in response to the repeated aggressive calls from the tamarin sentinel. At that moment, was observed a cat with small body but big feet, huge eyes and a long tail walking down the trunk of a tree (like a squirrel); it quickly jumped to a liana that was connected to the fig tree and moved toward where the tamarins were feeding, about 15 meters away. At this moment, the sentinel emitted a high scream as the predator approached the group; and the group fled immediately.

In our observations, the strategy used by *Leopardus wiedii* to imitate its prey was not effective in catching *Saguinus bicolor*. However, we suggest that this strategy is very effective in attracting prey, facilitating the attack and reducing energy expenditure during a possible pursuit. Curiously, all the potential prey (agoutis, macucos, and nambus) cited by

the Amazonian inhabitants produce extremely acute vocalizations, which possibly match the potential repertoire of felines. In addition, all the aforementioned potential prey species use vocalizations in intra-specific territorial demarcation. This increases the cats' chance of success in attracting prey by imitation.

Acknowledgments

We thank INPA (Instituto Nacional de Pesquisas da Amazônia) for permission to carry out fieldwork at the Reserva Florestal Adolpho Ducke. The observations reported here were made during a study supported by grant from PROBIO – MMA but we also wish to thank CNPq to provided financial support to Fabiano Calleia and the Wildlife Conservation Society (WCS) and Conservation Leadership Program (CLP) for provided financial support to Fabio Röhe. We are deeply indebted to several collaborators for their inestimable help in the fieldwork (Projeto Sauim-de-Coleira) supported by PROBIO/MMA, FNMA/MMA, WCS, CI, Durrell Wildlife Conservation Trust Ap-enheul Primate Conservation Trust, Shaldon Wildlife Trust, La Palmyre Zoo, Newquay Zoo, and Philadelphia Zoo. We thank M. Benchimol for comments in the manuscript and B. G. Luize for the interview in Purus River.

Fabiano de Oliveira Calleia, Projeto Sauim-de-Coleira/UFAM <fabianocalleia@gmail.com>, **Fabio Rohe**, WCS – Wildlife Conservation Society <fabiorohe@gmail.com>, **Marcelo Gordo**, UFAM/Projeto Sauim-de-Coleira <mgordo@ufam.edu.br>.

References

Beebe, W. 1925. Studies of a tropical jungle: one quarter of a square mile of jungle at Kartabo, British Guiana. *Zoologica* 6:1–193.

Bianchi, R. C. and Mendes, S. L. 2007. Ocelot (*Leopardus pardalis*) predation on primates in Caratinga Biological Station, Southeast Brazil. *Am. J. Primatol.* 69:1–6.

Table 2. Reports of mimicking vocalizations of *Puma concolor*, *Panthera onca* and *Leopardus pardalis*.

Location	Cat Species	Prey Species Imitated
Rio Madeira/2005*	<i>Panthera onca</i>	<i>Dasyprocta fuliginosa</i>
Rio Aripuaná/2005*	<i>Panthera onca</i>	<i>Crypturellus</i> sp.
Rio Aripuaná/2005*	<i>Puma concolor</i>	<i>Dasyprocta</i> sp.
Rio Aripuaná/2005*	<i>Leopardus pardalis</i>	<i>Crypturellus</i> sp.
Rio Juruá/2004	<i>Leopardus pardalis</i>	<i>Crypturellus</i> sp.
Rio Juruá/2004	<i>Puma concolor</i>	<i>Crypturellus</i> sp.; <i>Tinamus</i> sp.
Rio Javari/2009	<i>Panthera onca</i>	<i>Crypturellus</i> sp.
Rio Purus/2009	<i>Leopardus</i> sp.	<i>Crypturellus undulatus</i>
Rio Urubu/2006	<i>Puma concolor</i>	<i>Dasyprocta leporina</i>
Atlantic Forest/2003**	<i>Puma concolor</i>	<i>Crypturellus obsoletus</i>

*see Röhe 2007 for a description of the area.

** Atlantic Forest location is Serra de Paranapiacaba, São Paulo State, Brazil (see Röhe *et al.* 2003; Tófoli *et al.* 2009).

- Cabrera, A. and Yepes, J. 1940. *Mamíferos Sul-Americanos (Vida, Costumbres y Descripción)*. Historia natural Ediar. Compañía Argentina de Editores.
- Chinchilla F. A. 1997. La dieta del jaguar (*Panthera onca*), el puma (*Felis concolor*) y el manigordo (*Felis pardalis*) en el Parque Nacional Corcovado, Costa Rica. *Revista de Biología Tropical* 45: 1223–1229.
- Emmons, L. H. (1987). Comparative feeding ecology of felids in a neotropical rainforest. *Behav. Ecol. Sociobiol.* 20: 271–283.
- Goldizen, A. W. 1987. Tamarins and Marmosets: communal care of offspring. In: *Primates Societies*, Smuts, B. B.; Cheney, D. L., Seyfert, R. M., Wrangham, R. W. and Strusaker, T. S. (eds.), Pp. (34–43). The University of Chicago Press.
- Gordo, M., Calleia, F. O., Moreira, A. L. B. and Leite, J. J. F. 2005. Estrat gia de fuga e vigil ncia de grupos selvagens de *Saguinus bicolor*. *XI Congresso brasileiro de primatologia*, Porto Alegre pp. 107.
- Izawa, K. 1978. A field study of the ecology and behavior of the Black-mantle Tamarin (*Saguinus nigricollis*). *Primates* 19(2): 241–274.
- Konecny, M. J. 1989. Movement patterns and food habits of four sympatric carnivore species in Belize, Central America. In: *Advances in Neotropical Mammalogy* (Redford, K. H. and Eisenberg, J. F. (eds.), Pp. 243–264. Sandhill Crane Press, Florida.
- Matsuda, I. and Izawa, K. 2008. Predation of wild spider monkeys at La Macarena, Colombia. *Primates* 49: 65–68.
- Miranda, J. M. D., Vernardi, I. P., Abreu K. C. and Passos F.C. 2005. Predation on *Alouatta guariba clamitans* Cabrera (Primates, Atelidae) by *Leopardus pardalis* (Linnaeus) (Carnivora, Felidae). *Rev. Bras. Zool.* 22:793–795.
- Mondolfi, E. 1986. Notes on the biology and status of the small wild cats in Venezuela. in *Cats of the world: biology, conservation, and management*, Miller, S. D. and Everett, D. D. (eds.), Pp. 125–146. *National Wildlife Federation*, Washington, District of Columbia.
- Novack A. J., Main M. B., Sunquist M. E., Labisky R. F. 2005. Foraging ecology of jaguar (*Panthera onca*) and puma (*Puma concolor*) in hunted and non-hunted sites within the Maya Biosphere Reserve, Guatemala. *J. of Zool.* 267: 167–178.
- Oliveira, T. G. 1998. *Leopardus wiedii*. *Mammalian species* 579: 1–6.
- Oliveira, T. G. 1994. *Neotropical cats: ecology and conservation*. Edufma, S o Lu s, Brazil, 220 pp.
- Olmos, F. 1994. Jaguar predation on murequi, *Brachyteles arachnoides*. *Neotrop. Primates* 2: 16.
- Peetz, A., Norconk, M. A. and Kinzey, W. G. 1992. Predation by jaguar on howler monkeys (*Alouatta seniculus*) in Venezuela. *Am. J. Primatol.* 28: 223–228.
- Ribeiro, J. E. L. S., Hopkins, M. J. C., Vicentini, A., Sothters, C. A., Costa, M. A. S., Brito, J. M., Souza, M. A., Martins, L. H., Lohmam, L. G., Pereira, E. C., Silva, C. F., Mesquita, M. R. and Proc pio, L. C. 1999. *Flora da Reserva Ducke: guia de identifica  o das plantas vasculares de uma floresta de terra firme na Amaz nia Central*. INPA, Manaus, 816 pp.
- R he, F., Antunes, A. P. and T foli, C. F. 2003. The Discovery of a new sub-population of the black lion tamarins (*Leontopithecus chrysopygus*) at Serra de Paranapiacaba, S o Paulo, Brazil. *Neotrop. Primates*, 11(2): 75–76.
- R he, F. 2007. *Mam feros de m dio e grande porte do m dio Rio Madeira*. In: Py-Daniel, L. R. et al. (Org.). *Biodiversidade do M dio Madeira*. INPA/MMA.
- T foli, C. F.; R he, F.; Setz, E. Z. F. 2009. Jaguarundi (*Puma yagouaroundi*) food habits in mosaic of Atlantic Rainforest and eucalypt plantations of southeastern Brazil. *Braz. J. Biol.* 69(3): 631–637.
- Ximenes A. 1982. Notas sobre f lidos neotropicales VIII: Observaciones sobre el contenido estomacal y el comportamiento alimentar de diversas especies de felinos. *Rev. Nord. de Biol.* 5(1): 89–91.

PERUVIAN RED UAKARI MONKEYS (*CACAJAO CALVUS UCAYALII*) IN THE PACAYA-SAMIRIA NATIONAL RESERVE — A RANGE EXTENSION ACROSS A MAJOR RIVER BARRIER

Mark Bowler

Javier Noriega Murrieta

Maribel Recharte

Pablo Puertas

Richard Bodmer

According to Hershkovitz (1987) *Cacajao calvus ucayalii*, listed as Vulnerable by the IUCN, (Veiga & Bowler, 2008) is distributed from the east bank of the Rio Ucayali in an easterly direction to the Rio Yavar  and from the Rio Amazonas in the north to the Rio Urubamba in the south. Hershkovitz (1987) also includes the east bank of the lower Yavar  in Brazil, but its presence there has not been confirmed and it is possible that museum specimens marked as collected on the Brazilian bank of the Yavar  actually came from the Peruvian side where this primate is locally abundant. Surveys conducted between 1979 and 1986 (Aquino 1988) showed that the range was much reduced, hunting having exterminated the species in several areas. Aquino (1988) suggested that the southern limit is now probably the Rio Sheshea and that populations close to the Rios Ucayali and Amazonas have also been reduced and in some areas populations have been exterminated (Fig. 1). Populations of *Cacajao calvus* observed by Peres (1997) on the upper Rio Juru  and unconfirmed reports by Fernandes (1990) in the Brazilian state of Acre on the upper Juru  and Purus are either of *Cacajao calvus novaesi* or *Cacajao calvus ucayalii*, which would extend the known ranges of either of these subspecies.

The Rio Ucayali is the largest tributary of the Rio Amazonas and at 400–1,200m wide presents a significant barrier to primate populations. However, the constantly-changing course of the river means that very large islands of forest