

English Common Names for Subspecies and Species of African Primates

Author: Grubb, Peter

Source: Primate Conservation, 2006(20) : 65-73

Published By: Conservation International

URL: <https://doi.org/10.1896/0898-6207.20.1.65>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

English Common Names for Subspecies and Species of African Primates

Peter Grubb

35, Downhills Park Road, London N17 6PE, UK

Abstract: Approximately 1,000 English-language names have been used for African primates. Grubb *et al.* (2003) chose a single common name for each species (with a few exceptions) and for each subspecies. The present paper provides the opportunity to compare these preferred names with others published in the literature. The aim is to encourage primatologists to evaluate the choice of names, to assess the principles adopted in compiling the selective list, to amend this list where they see fit, preferably in appropriate publications, and to comment on the whole exercise.

Key Words: Common names, African primates, species, subspecies

Introduction

This paper lists published English-language common names for species and subspecies of African primates in a systematic format. The aim is to provide primatologists and zoologists with the opportunity to decide whether a particular name should be chosen for each taxon, and whether the list of names previously selected (Grubb *et al.* 2003) should be accepted or modified. Readers may question the principles adopted in compiling that list, the merits of making lists of common names at all, or the selection of what are supposed to be the best of these names.

The Compilation

English-language common names for species and subspecies of African primates were found in the references listed at the end of this paper. In Table 1, the names are listed as published, except for the following alterations:

- Even if the whole name was capitalized in the original, only proper nouns (and adjectives) are capitalized here, and then with some exceptions. Anubis, Diana, Magot, Malbrouck, Mangabey, Pluto, Satan, Tantalus, and Thoth are not capitalized as (parts of) primate names. In general English usage, the common names of animal species are not proper names and do not have capital initials. There are contexts in which it may be appropriate to regard species as individuals, but when a common name can be used in the plural, one cannot justify treating it as a proper name that therefore requires it to be capitalized. This is not to deny that species names are often capitalized in titles or headings. Some authors prefer to capitalize common names, and some serial publications require this to be done—no doubt for sound reasons.
- Corrections are made to misspelled surnames such as Bate, de Brazzae, Preussis, and Vleeschower (i.e., Bates, de Brazza, Preuss, and Vleeschowers).
- Possessive forms of personal names are standardized—Peters’s, Pousargues’s, or Sykes’s instead of Peters’, etc. Some authors avoid possessives in vernacular animal names, though I have found few instances among names of African primates, “Foa red colobus” (instead of Foa’s) being an example.
- Gordons’ instead of Gordon’s for red or bay colobus, *Procolobus gordonorum*, because it was named after the brothers Von Gordon.
- Fernando Po instead of Fernando Poo as the old name of Bioko.
- Bush-baby instead of bushbaby or bush baby.
- Night-ape instead of night ape or nightape.
- Moholi galago instead of mohol galago for *Galago moholi*. The assumption is that “moholi” is a genitive form of a nominative “mohol,” whereas it is actually a noun in apposition—a version of the Tswana moHwele or mogwêlê.

Names for each taxon are in alphabetical order and are followed by any names applied solely to what are now synonyms (with the synonym in parentheses). Such common names may come back into use if these synonymized taxa are restored to validity. Synonyms for which no common name has been proposed are not cited. Words or letters in parentheses are used in a name by some authors but not others. Taxonomy follows that proposed by Grubb *et al.* (2003). Some subspecies are regarded as full species in other publications. For example, Groves (2001) treated *Cercocebus galeritus sanjei* as *C. sanjei* and here I list it as *C. (galeritus) sanjei* to indicate the different opinions concerning its rank. The species *C. galeritus*, therefore, has a more extensive compass or *sensu lato*, and a more restricted compass or *sensu stricto*, for which authors have assigned different names (Table 1). I draw attention to names used for more than one taxon (other than species and their nominate subspecies) and names that imply occurrence in an area where the taxon has not been found. Names selected by Grubb *et al.* (2003) are in bold.

In compiling their list, Grubb *et al.* (2003) adopted principles and made decisions that were not recorded at the time but are listed here, as follows:

- Two or more names—not a single one—were provided for each of the following species: *Macaca sylvanus*, *Cercocebus atys*, *Cercopithecus aethiops*, *C. diana*, *C. campbelli*, *C. pogonias*, *C. mitis*, *Pan troglodytes*, and *P. paniscus*. Choice of a preferred name is not always easy. “Common” in animal names—such as “common chimpanzee”—may be interpreted as “abundant,” particularly by those for whom English is not their first language, according to Duckworth and Pine (2003), who would avoid it. If instead “chimpanzee” were the preferred name for *Pan troglodytes*, and *P. paniscus* was called “pygmy chimpanzee,” “chimpanzee” would refer as well to all *Pan* species. This ambiguity is also to be avoided, but if *P. paniscus* were called “bonobo,” the difficulty would not arise. However, A. Kortlandt (in litt.) found objections to the use of “bonobo” as a common name—but “pygmy chimpanzee” is not literally descriptive, so need not be acceptable either. If a decision must be made, choice of “chimpanzee” and “bonobo” as preferred names may be the least objectionable option.
- Separate names for species and nominate subspecies were provided. A nominate subspecies has often been given the same common name as the species but while its status as a subspecies can still be identifiable—for instance, *Procolobus pennantii pennantii* could be “Pennant’s red colobus (nominate race),” a separate name can be less ambiguous. Where a single polytypic species is partitioned into several species, the original common name may be abandoned because it becomes ambiguous and a new name is required—an erstwhile subspecies name. Thus when the gorilla was partitioned into two species, these became “western gorilla” and “eastern gorilla,” and

similarly the angwantibo became “Calabar angwantibo” and “golden angwantibo.”

- Surnames of people commemorated in primate nomenclature were used in the possessive form (for example, Stuhlmann’s blue monkey, not Stuhlmann blue monkey).
- Where appropriate, locality names were identified topographically (Bale Mountain grivet, not Bale grivet; or Omo River guereza, not Omo guereza).
- Adjectival forms of places were avoided (Angola black-and-white colobus, not Angolan black-and-white colobus).
- “Galago” was used in preference to “bush-baby.”
- “Hamadryas” was chosen because it is now widely used, although the spelling in the Oxford English Dictionary is “hamadryad.”
- “Monkey” was used in preference to “guenon.”
- *Cercopithecus dryas* was called the dryad monkey. Dryad—not dryas—is the accepted spelling (Oxford English Dictionary).
- “Guereza”—an Ethiopian name for *Colobus guereza*—had been used as if it were a synonym of “colobus monkey” by Forbes (1894), Elliot (1913–1914) and Sanderson (1957), but nowadays is once again applied only to *C. guereza*.

A few changes from the original list (Grubb *et al.* 2003) are made here, as follows:

- The Uganda lesser galago (*Galago senegalensis sotikae*) is known only from the type locality (the Telek River, Sotik, in Kenya) so “Sotik lesser galago” (not “Sotik River galago”) would be preferable.
- The Ibean yellow baboon (*Papio cyncephalus ibeanus*) is named after IBEA, an acronym of the short-lived Imperial British East Africa Company, with which this taxon has no particular association. “Northern yellow baboon” would be preferable.
- The western gelada (*Theropithecus gelada gelada*) and eastern gelada (*T. g. obscurus*) occur respectively in the northern and southern sectors of their species’ range. “Northern gelada” and “southern gelada” would be preferable names.
- *Colobus guereza percivali* is called the Mt Uarges guereza, but Uaraguess is an approved spelling, so “Mt Uaraguess guereza” would be preferable.

Altogether there are approximately 1,000 English names for 174 species and subspecies of the African primate fauna.

Discussion

The prolixity of common names for African primates (Table 1) does not appear to arise from strong preferences or differences in opinion among naturalists but rather by a failure to follow precedent, leading to a clutter of permutations

Table 1. List of scientific and vernacular names for species and subspecies of African primates. The names listed in Grubb *et al.* (2003) are in bold type. See “The Compilation” for further explanation.

Scientific names of species and subspecies	Vernacular names
Galagos and lorises (b. = bush-baby; g. = galago)	
<i>Galagoides demidovii</i>	Demidoff's b. or g., Demidoff's dwarf g. , dwarf b. or g., Prince Demidoff's b. or g.; Fernando Po g. (<i>poensis</i>); mouse g., Murray's g. (<i>murinus</i>)
<i>G. thomasi</i>	Thomas's b. or g., Thomas's dwarf g.
<i>G. orinus</i>	Amani dwarf g., mountain g., mountain dwarf g. , Southern dwarf g., Uluguru b. or g., Usambara g.
<i>G. rondoensis</i>	Rondo b. or g., Rondo dwarf g.
<i>G. zanzibaricus</i>	eastern lesser g., Zanzibar dwarf g., Zanzibar g. or b., Zanzibar lesser b.; Matundu (dwarf) g., Udzungwa dwarf g., Uzungwa b. (<i>udzungwensis</i>)
<i>G. cocos</i>	Diani g., Diani small g., Kenya coast g. , Kenya coast dwarf g.
<i>G. granti</i>	Grant's b. or g., Grant's dwarf g., Grant's lesser b., Grant's night-ape, Mozambique g. , Newala small g.
<i>G. nyasae</i>	Malawi g. or b., Malawi lesser g.
<i>G. sp. nov. 1</i>	Kalwe g. , Kalwe dwarf g.
<i>G. sp. nov. 2</i>	Mt Thyolo g. , Mt Thyolo dwarf g.
<i>G. sp. nov. 3</i>	Ukinga g. , Ukinga dwarf g.
<i>Galago senegalensis</i>	bushbaby, lesser b. or g., northern lesser g. or b., Senegal b. or g.
<i>G. s. senegalensis</i>	Senegal b. or g., Senegal lesser g. ; white-footed b. or g. (<i>albipes</i>)
<i>G. s. dunni</i>	Abyssinian g., Ethiopian b. or g., Ethiopia lesser g. , Ethiopian lesser g.
<i>G. s. sotikae</i>	Sotik River g., Uganda lesser g. [rejected; replace by Sotik lesser g.], Ugandan lesser g.
<i>G. s. braccatus</i>	Kenya lesser g. , Kenyan lesser g., Mt Kilimanjaro g. [based on erroneous supposition that this was the type locality], yellow-thighed b. or g.
<i>G. gallarum</i>	Boran g., Somali b. or g., Somali lesser g.
<i>G. moholi</i>	bush-baby [also for <i>Otolemur crassicaudatus</i>], lesser b., maholi g., moholi b. or g., moholi lemur, night-ape, South African g., southern lesser g. or b.
<i>G. m. moholi</i>	moholi g., moholi lesser g. ; Mozambique g. [also for <i>Galagoides granti</i>] (<i>mossambicus</i>)
<i>G. m. bradfieldi</i>	Namibia lesser g. , Namibian lesser g.
<i>G. matschiei</i>	dusky b., east needle-clawed b., eastern needle-clawed b. or g., eastern needle-nailed b., Matschie's b., needle-clawed g., spectacled g., spectacled lesser g.
<i>Euoticus elegantulus</i>	needle-clawed b. or g., southern needle-clawed g. or b., elegant needle-clawed g., western needle-clawed g.; Du Chaillu's g. (<i>apicalis</i>)
<i>E. pallidus</i>	northern needle-clawed g. or b., pale-colored g., pallid needle-clawed g.
<i>E. p. pallidus</i>	Bioko needle-clawed g. , pale g.
<i>E. p. talboti</i>	Nigeria needle-clawed g.
<i>Sciurocheirus alleni</i>	<i>sensu lato</i> : Allen's g. , Allen's squirrel g. <i>sensu stricto</i> : Allen's squirrel g., Bioko Allen's g. or b.
<i>S. (alleni) cameronensis</i>	Cameroon g., Cross River Allen's b. , Cross River b., Cross River squirrel g.; Bates's g. (<i>batesi</i>)
<i>S. gabonensis</i>	Gabon Allen's g. or b., Gabon squirrel g., Gabon b. or g.
<i>S. sp. nov.</i>	Makande Allen's g.
<i>Otolemur garnettii</i>	black g., Garnett's b. or g., Garnett's greater b. or g., greater b. [also used for <i>O. crassicaudatus</i>], northern greater g., small-eared g., small-eared greater g.
<i>O. g. garnettii</i>	black g., Garnett's b. or g., Zanzibar small-eared g. , Zanzibar Garnett's greater g.; dark-tailed g., komba g. (<i>agisymbanus</i>)
<i>O. g. lasiotis</i>	white-tailed Garnett's greater g., white-tailed small-eared g. or b., woolly-eared g.
<i>O. g. panganiensis</i>	Pangani b. or g., Pangani Garnett's greater g., Pangani small-eared g.
<i>O. g. kikuyuensis</i>	Kikuyu b. or g., Kikuyu Garnett's greater g., Kikuyu small-eared g.
<i>O. crassicaudatus</i>	brown greater g., bush-baby, fat-tailed b., grand b. or g., great g., greater b. or g., large-eared greater b. or g., large grey night-ape, thick-tailed b. or g., thick-tailed greater g.
<i>O. c. crassicaudatus</i>	grand b. or g., great g., greater b. or g., South African thick-tailed g. , South African thick-tailed greater g., thick-tailed b. or g.; dusky g. (<i>umbrosus</i>); Zulu g. (<i>zuluensis</i>)
<i>O. c. kirkii</i>	Kirk's g., Tanganyika thick-tailed g. , Tanganyika thick-tailed greater g.; bay g. (<i>badius</i>), Lönnberg's g. (<i>loemnergi</i>)
<i>O. monteiroi</i>	Monteiro's g., silver g., silver greater g. , silvery greater g.
<i>O. m. monteiroi</i>	miombo silver g. , miombo silver greater g., Monteiro's b. or g.
<i>O. m. argentatus</i>	northern silver g. , northern silver greater g., silvery b. or g.
<i>O. sp. nov.</i>	Mwera greater g. [status questionable]
<i>Arctocebus calabarensis</i>	angwantibo, awantibo, Calabar angwantibo , Calabar potto, golden potto [name also used for <i>A. aureus</i>]
<i>A. aureus</i>	golden angwantibo , (southern) golden potto
<i>Perodicticus potto</i>	Bosman's potto, potto
<i>P. p. potto</i>	(van) Bosman's potto, western potto ; (south) Nigerian potto (<i>juju</i>)
<i>P. p. edwardsi</i>	Cameroon potto, central potto , Central African potto, Milne-Edwards's potto

continued on next page

Table 1. continued from previous page

Scientific names of species and subspecies	Vernacular names
<i>P. p. ibeanus</i>	East African potto, eastern potto
<i>Pseudopotto martini</i>	false potto, long-tailed potto, Martin's false potto [taken to be a synonym of <i>Perodicticus potto edwardsi</i>]
Monkeys of the tribe Papionini (b. = baboon; ma. = mangabey)	
<i>Macaca sylvanus</i>	Barbary ape, Barbary macaque , magot, rock ape
<i>Cercocebus galeritus</i>	<i>sensu lato</i> : agile ma., crested ma., plain-headed ma., river ma. <i>sensu stricto</i> : crested ma., Peters' ma., Tana ma., Tana River ma.
<i>C. (g.) agilis</i>	agile ma. , olive ma.; Hagenbeck's ma. (<i>hagenbecki</i>)
<i>C. (g.) sanjei</i>	Sanje (crested) ma., Sanje river ma.
<i>C. (g.) chrysogaster</i>	golden-bellied ma.
<i>C. atys</i>	sooty ma., sooty ma. or white-naped ma. , spectacled ma.
<i>C. a. atys</i>	smoky ma., sooty ma. , Western sooty ma.
<i>C. a. lumulatus</i>	crescent-naped sooty ma., crowned ma., white-crowned ma., white-naped ma.
<i>C. torquatus</i>	capped ma., cherry-capped ma., collared ma., red-capped ma. , smoky ma., white ma., white-collared ma.
<i>Mandrillus sphinx</i>	mandrill ; Fernando Po mandrill (<i>insularis</i>) [the species does not occur on Bioko = Fernando Po]; Gabonese mandrill, southern mandrill (<i>adarogaster</i>); northern mandrill (supposed nominate ssp. <i>sphinx</i>)
<i>M. leucophaeus</i>	Drill
<i>M. l. leucophaeus</i>	mainland drill , southern drill; Cameroon drill, northern drill (<i>mundamensis</i>)
<i>M. l. poensis</i>	Bioko drill , Fernando Po drill
<i>Lophocebus albigena</i>	Black ma. [name more usually given to <i>L. aterrimus</i>], crested ma. [also for <i>Cercocebus galeritus</i> , <i>L. aterrimus</i>], grey-cheeked ma. , grey-cheeked crested ma., mantled ma., white-cheeked ma.; Johnston's ma., Johnston's grey-cheeked crested ma., (<i>johnstoni</i>), Osman Hill's grey cheked crested ma. (<i>osmani</i>); western grey-cheeked crested ma. (nominate <i>albigena</i>); Zenker's ma. (<i>zenkeri</i>)
<i>L. aterrimus</i>	<i>sensu lato</i> : black ma. , black crested ma., crested ma., peaked ma. <i>sensu stricto</i> : black ma., black crested ma. [also for <i>Cercocebus galeritus</i> , <i>L. albigena</i>], Congo ma., northern black ma. , northern black crested ma., tufted ma., typical black ma.
<i>L. (a.) opdenboschi</i>	black-cheeked black ma., Opdenbosch's black crested ma., Opdenbosch's (crested) ma., Southern black ma.
<i>Papio cynocephalus</i>	<i>sensu lato</i> or <i>sensu stricto</i> : yellow b.
<i>P. c. cynocephalus</i>	eastern yellow b., central yellow b. , typical yellow b.; baboon (<i>babouin</i>); dwarf chacma, Zambian dwarf chacma b. (<i>jubilaeus</i>); Langheld's b. (<i>langheldi</i>); Nyasa yellow b., southern yellow b. (<i>streptus</i>); thoth (b.) (<i>thoth</i>)
<i>P. c. ibeanus</i>	northern yellow b., Ibean b., Ibean yellow b. [rejected; replace by northern yellow b.]
<i>P. (c.) kindae</i>	western yellow b., Kinda b., Kinda yellow b. , Rhodesian long-armed b.
<i>P. ursinus</i>	<i>sensu lato</i> : chacma , chacma b., pig-tailed b., South African b. <i>sensu stricto</i> : Cape chacma (b.), chacma b., southern chacma ; eastern Cape b., eastern chacma b. (<i>orientalis</i>); Transvaal chacma b. (<i>occidentalis</i>); Kalahari b. or chacma, Shortridge's chacma b., western chacma b. (<i>ruacana</i>)
<i>P. (u.) griseipes</i>	grey-footed chacma (b.), Rhodesia b., Rhodesian chacma (b.); Botswana chacma b. (<i>chobiensis</i>)
<i>P. anubis</i>	anubis b., doguera b., olive b. ; Abyssinian b., Ethiopian olive b., doguera b. (<i>doguera</i>); Baringo b., Rift Valley b., western Kenya olive b. (<i>furax</i>); Congolese olive b. (<i>tesselatus</i>); East African olive b., Neumann's (olive) b. (<i>neumanni</i>); Nigeria b. (<i>nigeriae</i>); West African anubis b. (nominate <i>anubis</i>); western olive b. Heuglin's (olive) b., Sudanese b. (<i>heuglini</i>)
<i>P. papio</i>	Guinea b. , papion, western b.
<i>P. hamadryas</i>	Arabian b., hamadryad, hamadryas, hamadryas b. , mantled b., sacred (Anubis) b.
<i>Theropithecus gelada</i>	gelada (b.)
<i>T. g. gelada</i>	common gelada, northern gelada, western gelada [rejected; replace by northern gelada]
<i>T. g. obscurus</i>	dusky gelada, eastern gelada [rejected; replace by southern gelada] , southern gelada
Monkeys of the tribe Cercopithecini (g. = guenon; m. = monkey)	
<i>Allenopithecus nigroviridis</i>	Allen's baboon-like m., Allen's m., Allen's swamp m. , blackish-green g., swamp g.
<i>Miopithecus talapoin</i>	Angola(n) talapoin, dwarf g., mangrove m., southern talapoin, southern talapoin m. ; Angola talapoin, Ansoerge's talapoin (<i>ansorgei</i>); common talapoin, Gabon talapoin [does not occur in Gabon] (supposed nominate ssp. <i>talapoin</i>); Ruwenzori talapoin (<i>pilettei</i>) [does not occur on the Ruwenzori Mountains]; Vleeschouwers's talapoin, Zaire talapoin (<i>vleeschouwersi</i>)
<i>M. ogouensis</i>	Gabon talapoin, northern talapoin, northern talapoin m.
<i>Erythrocebus patas</i>	hussar m., military m., patas, patas m. , nisanas m., red m.; Air patas (m.), Air red m. (<i>villiersi</i>); black-nosed patas, patas g. or m., West African patas (m.), West African red m. (nominate <i>patas</i>); Blue Nile hussar m., dancing red m., Nile patas (m.), nisanas (g.), nisanas m., (<i>pyrrhonotus</i>); Ikoma patas (m.) (<i>baumstarki</i>)
<i>Cercopithecus aethiops</i>	<i>sensu lato</i> : green m., grey vervet, grivet, savanna m., vervet (m.); including green, malbrouck, and tantalus monkeys, grivets, and vervets <i>sensu stricto</i> : common grivet m., Ethiopian grivet, grivet (g. or m.), tota; Abyssinian grivet, Abyssinian vervet m., Hilgert's grivet (<i>hilgerti</i>); Matschie's grivet or g., Matschie's grivet m., (<i>matschiei</i> [or <i>ellenbecki</i>])
<i>C. (a.) djamdjamensis</i>	Bale (Mountains) m., Bale Mountains grivet or vervet, Djamdjam grivet or g.
<i>C. (a.) tantalus</i>	tantalus m. or g.; Jebel Marra tantalus (m.) (<i>marrensis</i>); Budgett's tantalus (m.) (<i>budgetti</i>)
<i>C. (a.) sabaesus</i>	callithrix, green m. or g.; Werner's g. (<i>wernerii</i>)

continued on next page

Table 1. continued from previous page

Scientific names of species and subspecies	Vernacular names
<i>C. (a.) pygerythrus</i>	common vervet, vervet , vervet g. or m.; black-chinned vervet, Cuvier's vervet, South African vervet, southern vervet m., vervet (g.) (nominate <i>pygerythrus</i>); black-faced vervet, Bukoba green m. (<i>centralis</i>); Cloet's vervet (<i>cloeti</i>); de Beaux's grivet (<i>zavattarii</i>); desert tumbuli m., Heller's vervet, northeastern vervet (<i>arenarius</i>); East African vervet, yellow m. (<i>johnstoni</i>); Isle of Patta vervet, Manda vervet (m.) (<i>excubitor</i>); Mozambique g. or m. [same name for <i>C. mitis erythrarchus</i>], reddish-green g., russet-green or russety-green vervet (<i>rufoviridis</i>); Naivasha vervet (<i>callidus</i>); Namaqualand vervet (<i>marjoriae</i>); Okavango vervet (<i>ngamiensis</i>); Pemba vervet (m.), Zanzibar vervet (<i>nesiotes</i>); Whyte's vervet (<i>whytei</i>)
<i>C. (a.) cynosuroides</i>	malbrouck (g.), malbrouck m. ; Cunene vervet (<i>helvescens</i>);
<i>C. lhoesti</i>	east Congo l'Hoest's m., l'Hoest's g., l'Hoest's m. , mountain g. or m.
<i>C. preussi</i>	Cross's m., Preuss's m. or g.
<i>C. p. preussi</i>	mainland Preuss's g., Preuss's m.
<i>C. p. insularis</i>	Bioko Preuss's m. or g., Fernando Po l'Hoest's m. [but not a subspecies of <i>C. lhoesti</i>]
<i>C. solatus</i>	sun-tailed m. , sun-tailed g.
<i>C. diana</i>	<i>sensu lato</i> : diana g. or m., diana and roloway monkeys <i>sensu stricto</i> : diana (g.), diana m. , typical diana m.
<i>C. (d.) roloway</i>	roloway m. , roloway diana m.; palatine g. (<i>palatinus</i>)
<i>C. dryas</i>	Congo diana m., dryad m. , dryas g., dryas m.; salongo g. or m., Zaire diana g. or m. (<i>salongo</i>)
<i>C. neglectus</i>	chestnut-browed g., neglectus m., Schlegel's g., Brazza's m., de Brazza's m. or g. (<i>brazzae</i>)
<i>C. campbelli</i>	<i>sensu lato</i> : Campbell's m., Campbell's and Lowe's monkeys <i>sensu stricto</i> : Campbell's mona, g., or m.
<i>C. (c.) lowei</i>	Lowe's mona or g., Lowe's mona m., Lowe's m.
<i>C. mona</i>	mona g., mona m. , mone, typical mona m.
<i>C. pogonias</i>	<i>sensu lato</i> : crowned and Wolf's monkeys <i>sensu stricto</i> : bearded g. or m., crested mona (m.), crowned m. or g., golden bellied g.
<i>C. p. pogonias</i>	bearded g. or m., crowned g., crowned mona m., golden-bellied crowned m. or g.
<i>C. p. nigripes</i>	black-footed crowned m. or g., black-footed g., black-footed mona (m.)
<i>C. p. grayi</i>	Erleben's g., Gray's crowned m. or g., Gray's g., Gray's mona (m.); white-crowned g. (<i>petronellae</i>)
<i>C. (p.) denti</i>	Dent's mona or g., Dent's mona m., Dent's m.
<i>C. (p.) wolffi</i>	Wolf's mona or g., Wolf's mona m., Wolf's m.
<i>C. p. or w. wolffi</i>	Wolf's mona or g., Wolf's mona m., Congo Basin Wolf's m. or g.
<i>C. p. or w. elegans</i>	elegant mona (m.), Lomami River Wolf's m. or g.
<i>C. p. or w. pyrogaster</i>	fire-bellied g., fire-bellied Wolf's m. or g., red-bellied mona or g.
<i>C. hamlyni</i>	Hamlyn's g. or m., Hamlyn's owl-faced m., owl-faced g., owl-faced m.
<i>C. h. hamlyni</i>	Hamlyn's owl-faced g., nose-stripe owl-faced m.
<i>C. h. kahuziensis</i>	Mt Kahuzi owl-faced m. or g.
<i>C. petaurista</i>	lesser spot-nosed m. or g., lesser white-nosed g. or m., spot-nosed m., white-cheeked hocheur
<i>C. p. petaurista</i>	lesser white-nosed g. or m., eastern lesser spot-nosed m. or g., typical lesser white-nosed g., white-cheeked hocheur
<i>C. p. buettikoferi</i>	Büttikofer's g. or m., Büttikofer's white-nosed g., Büttikofer's lesser white-nosed g., western lesser spot-nosed m. or g.
<i>C. signatus</i>	Jentink's g., Jentink's white-nosed m. [supposedly of hybrid origin; it has been thought—perhaps wrongly—that <i>C. erythrogaster</i> is one of the parental forms]
<i>C. erythrogaster</i>	red-bellied g. or m., white-throated g., white-throated m. [names also used for <i>albogularis</i> section of <i>C. mitis</i>]
<i>C. e. erythrogaster</i>	red-bellied m. or g., red-bellied white-throated g.
<i>C. e. pococki</i>	Nigeria white-throated m. , Nigerian white-throated g.
<i>C. sclateri</i>	Sclater's m. or g., Sclater's white-nosed g. or m.
<i>C. erythrotis</i>	red-eared g., red-eared m. , red-eared nose-spotted g. or m., russet-eared g., russet-eared nose-spotted g.
<i>C. e. erythrotis</i>	Bioko red-eared m. , Bioko russet-eared g., Fernando Po red-eared g., Fernando Po russet-eared g.
<i>C. e. camerunensis</i>	Cameroon red-eared m. or g., Cameroon russet-eared g.
<i>C. cephus</i>	moustach m., moustache, moustached g., (true) moustached m. , mustached m.
<i>C. c. cephus</i>	blue-faced m., moustached m., red-tailed moustached m. or g.
<i>C. c. cephodes</i>	grey-tailed moustached m. or g.
<i>C. c. ngottoensis</i>	Ngotto moustached m. or g., white-nosed moustached m.
<i>C. ascanius</i>	black-cheeked white-nosed m., coppertail m., redtail (m.), red-tailed m. or g., white-nosed m.; black-cheeked g. or hocheur (<i>melanogenys</i>)
<i>C. a. ascanius</i>	black-cheeked red-tailed m. or g., black-cheeked white-nosed m.
<i>C. a. schmidti</i>	Schmidt's g. or m., Schmidt's red-tailed m. or g., Uganda red-tailed g. or m.; montane red-tailed g. (<i>montanus</i>)
<i>C. a. whitesidei</i>	Whiteside's g. or m., yellow-nosed m., yellow-nosed red-tailed m. or g.
<i>C. a. katangae</i>	Katanga black-cheeked white-nosed m., Katanga red-tailed m. or g.

continued on next page

Table 1. continued from previous page

Scientific names of species and subspecies	Vernacular names
<i>C. a. atrinatus</i>	black-nosed m., black-nosed red-tailed m. or g.
<i>C. nictitans</i>	greater spot-nosed m. or g., greater white-nosed g. or m., hocheur (g or m.), putty-nosed m. or g., spot-nosed m., white-nosed g. or m.
<i>C. n. nictitans</i>	Eastern putty-nosed m. , Eastern greater spot-nosed g., greater white-nosed m., hocheur (g.)
<i>C. m. martini</i>	blackish hocheur, Martin's g., Martin's putty-nosed m., Martin's white-nosed g. or m.; ludio g. or m., white-chested hocheur (<i>ludio</i>); Stampfli's putty-nosed m. , Stampfli's greater spot-nosed g., Stampfli's g. (<i>stampflii</i>)
<i>C. (mitis) mitis</i>	<i>sensu lato</i> : blue m., diademed g. or m., gentle m., mitis m., samango m.; including blue m., Sykes's m., golden m., and samango <i>sensu stricto</i> : blue m., diademed g.; Carruthers's g. (<i>carruthersi</i>)
<i>C. m. heymansi</i>	Lomami River blue m. , Lomami River diademed g., Zaire Basin gentle m.
<i>C. m. mitis</i>	Angolan diademed g.; diadem (g. or m.), diademed g. or m. (<i>leucampyx</i>); black-bellied m., pluto diademed g., pluto m. (<i>pluto</i>)
<i>C. m. opisthostictus</i>	rump-spotted blue m. , rump-spotted (diademed) g.
<i>C. m. boutourlinii</i>	Boutourlini's blue m. , Boutourlini's (diademed) g.
<i>C. m. stuhlmanni</i>	blue m., Stuhlmann's diademed g., Stuhlmann's g. or m., Stuhlmann's blue m. ; Congo diademed g., Maes's g. (<i>maesi</i>); Mt Elgon diademed g. (<i>elgonis</i>)
<i>C. m. schoutedeni</i>	Schouteden's blue m. , Schouteden's diademed g., Schouteden's g. or m.
<i>C. (mitis) doggetti</i>	Ankole blue m., Doggett's g., silver m. or g.
<i>C. (mitis) kandti</i>	bamboo m., Congo red m., golden m. or g., red Congo m.; orange-colored g. (<i>insignis</i>)
<i>C. (mitis) albogularis</i>	<i>sensu lato</i> (<i>albogularis</i> section of <i>C. mitis</i>): samango, Sykes's m. or g., white-throated g. <i>sensu stricto</i> , as a ssp.: blue m. [also the name for other sections of the species, <i>sensu lato</i>], Sykes's m., white-throated g., Zanzibar Sykes's m. or g., Zanzibar white-throated g.; Kilimanjaro blue m., Kilimanjaro Sykes's g. (<i>kibonotensis</i>);
<i>C. m. or a. albotorquatus</i>	Pousargues's white-throated g., Pousargues's white-collared m. , Tana River Sykes's g.; Patta Island Sykes's g., Patta Island white-throated g. (<i>phylax</i>); Somali Sykes's g., Zammarano's white-throated g. (<i>zammaranoi</i>)
<i>C. m. or a. kolbi</i>	Kolb's g. or m., Kolb's white-collared m. , Mt Kenya Sykes's m. or g., Mt Kenya white-throated g., Sykes's m.
<i>C. m. or a. monoides</i>	maritime white-throated g. or m., Tanganyika Sykes's m., Tanzania Sykes's m. , Tanzanian Sykes's g.
<i>C. m. or a. francescae</i>	northern Malawi blue m., red-eared Sykes's m. or g.
<i>C. m. or a. erythrarcus</i>	Mozambique m., red-rumped Sykes's g.; Beira g. or m. (<i>beirensis</i>); Nyasa white-throated g. (<i>nyasae</i>); Stairs' g. or m., Stairs's white-collared m. (<i>stairsi</i>); Transvaal Sykes's g., Transvaal white-throated g. (<i>schwarzi</i>)
<i>C. m. or a. labiatus</i>	white-lipped g., white-lipped Sykes's g.; samango , samango g. or m. (<i>samango</i>)
<i>C. (mitis) moloneyi</i>	Lake Malawi Sykes's g., Moloney's g. or m., Moloney's white-collared m.
Monkeys of the subfamily Colobinae (c = colobus [or colob — archaic]; c.m. = colobus monkey; m = monkey; r. c. = red colobus)	
<i>Procolobus verus</i>	olive c. or c.m., Van Beneden's c. or guereza
<i>P. badius</i>	bay c. or c. m., Guinea Forest r. c., red c., western r. c.
<i>P. b. badius</i>	bay c. or guereza, upper Guinea r. c.
<i>P. b. temminckii</i>	Temminck's r. c. , Temminck's c. or guereza; fuliginous guereza (<i>fuliginosus</i>)
<i>P. b. waldroni</i>	Miss Waldron's r. c. , Miss Waldron's (bay) c.
<i>P. pennantii</i>	Pennant's r. c. , Pennant's c. m. or guereza, West Central African r. c.
<i>P. p. pennantii</i>	Bioko r. c. , Pennant's c. or r. c.
<i>P. p. bouvieri</i>	Bouvier's r. c. , Bouvier's c.
<i>P. p. epieni</i>	Delta r. c., Niger Delta r. c.
<i>P. (pennantii) preussi</i>	Cameroon r. c., Preuss c., Preuss's c., Preuss's r. c.
<i>P. foai</i>	<i>sensu lato</i> , including the following eight taxa, or <i>sensu stricto</i> including only the following six taxa: Central African r. c. [systematic status uncertain]
<i>P. f.? oustaleti</i>	Oustalet's r. c. or guereza; Powell-Cotton's c. (<i>powelli</i>) [almost the same name used for <i>Colobus angolensis cottoni</i>]
<i>P. f.? parmentieri</i>	Lomami River r. c. , Lomami r. c.
<i>P. f.? foai</i>	Foa's c. or r. c., Kivu r. c.
<i>P. f.? ellioti</i>	Elliot's c. or c.m. or r. c.; (<i>semlikiensis</i>) Semliki r. c.
<i>P. f.? lulindicus</i>	Lulindi River r. c.
<i>P. f.? langi</i>	Kisangani r. c.
<i>P. (f.?) tholloni</i>	Thollon c., Thollon's r. c., Tshuapa r. c.
<i>P. (f.?) tephrosceles</i>	ashy r. c. , bay Uganda r. c., Ugandan c., Ugandan r. c.
<i>P. rufomitratu</i>	red-crested guereza, Tana River r. c. [eastern r. c., if it includes <i>P. foai sensu lato</i>]
<i>P. gordonorum</i>	Gordons' r. c. or bay c., Iringa r. c., Udzungwa r. c. , Uhehe r. c., Uzungwa r. c.
<i>P. kirkii</i>	Kirk's c. or guereza, Kirk's r. c. , Zanzibar r. c.
<i>Colobus satanas</i>	black c. or guereza, satan c., satanic c., satanic black c.
<i>C. s. satanas</i>	Bioko black c.
<i>C. s. anthracinus</i>	Gabon black c.
<i>C. angolensis</i>	Angola(n) c. , Angola black-and-white c., Angolan c. or guereza, Angola pied c., Angolan black c., Angolan black-and-white c. (m.), white-epauletted black c.

continued on next page

Table 1. continued from previous page

Scientific names of species and subspecies	Vernacular names
<i>C. a. angolensis</i>	Angolan c. or guereza, mantled c., Sclater's Angola(n) c.
<i>C. a. palliatus</i>	Peters's Angola(n) c. , Tanzanian black c.; Sharpe's guereza (<i>sharpei</i>)
<i>C. a. ruwenzorii</i>	Adolf Friedrich's Angola(n) c. , Kivu black c., Ruwenzori c.
<i>C. a. cottoni</i>	Cotton's c., Powell-Cotton's Angola(n) c. , Powell-Cotton's black c.
<i>C. a. cordieri</i>	Cordier's Angola(n) c. , Cordier's black c., Cordier's c.
<i>C. a. prigoginei</i>	Prigogine's Angola(n) c. , Prigogine's black c., Prigogine's c.
<i>C. a. ssp. nov.</i>	Mahali Mountain Angola(n) c.
<i>C. polykomos</i>	full-bottom m., king c. or m., Western black-and-white c., Western pied c.; ursine c. or guereza (<i>ursinus</i>)
<i>C. vellerosus</i>	Geoffroy's black-and-white c., Geoffroy's pied c., ursine c. [more usually applied to <i>C. polykomos</i> ?], white-thighed c. or guereza, white-thighed black-and-white c., white-throated guereza; Dollman's (black-and-white) c. is a hybrid with <i>C. polykomos</i>
<i>C. guereza</i>	Abyssinian black-and-white c. (m.), Abyssinian c., (eastern) black-and-white c., guereza , guereza c., guereza black-and-white c., magistrate c., magistrate black c., mantled guereza, white-mantled c.
<i>C. g. occidentalis</i>	Congo guereza, magistrate c., Western guereza
<i>C. g. dodingae</i>	Dodinga Hills guereza
<i>C. g. matschiei</i>	Matschie's c., Mau Forest guereza
<i>C. g. guereza</i>	Abyssinian guereza, guereza, Omo River guereza , Rüppell's guereza, typical guereza; greyish-white-tailed guereza (<i>poliurus</i>)
<i>C. g. gallarum</i>	Neumann's (black) c., Djaffa Mountains guereza
<i>C. g. percivali</i>	Mt Uargess guereza [rejected; replace by Mt Uaraguess guereza]
<i>C. g. caudatus</i>	Kilimanjaro guereza or c., Mt Kilimanjaro guereza , white-tailed c. or guereza
<i>C. g. kikuyuensis</i>	Mt Kenya guereza or c.
Apes, Hominidae	
<i>Gorilla gorilla</i>	western gorilla
<i>G. g. gorilla</i>	coast gorilla, lowland gorilla, western gorilla, western lowland gorilla
<i>G. g. diehli</i>	Cross River gorilla
<i>G. beringei</i>	eastern gorilla
<i>G. b. beringei</i>	eastern gorilla, eastern highland gorilla, mountain gorilla
<i>G. b. graueri</i>	eastern lowland gorilla, Grauer's gorilla
<i>Pan troglodytes</i>	chimpanzee, common chimpanzee, robust or common chimpanzee
<i>P. t. verus</i>	common chimpanzee, masked chimpanzee, pale-faced chimpanzee, true chimpanzee, Upper Guinea chimpanzee, western chimpanzee
<i>P. t. vellerosus</i>	Nigeria chimpanzee , Nigerian chimpanzee
<i>P. t. troglodytes</i>	black chimpanzee, black-faced chimpanzee, central chimpanzee , Central African chimpanzee, choga, common chimpanzee, Lower Guinea chimpanzee, tschego, western chimpanzee; bald chimpanzee (<i>calvus</i>); gorilla-like chimpanzee, koolakamba (<i>koolokamba</i>)
<i>P. t. schweinfurthii</i>	long-haired chimpanzee, eastern chimpanzee , Schweinfurth's chimpanzee
<i>P. t. marungensis</i>	Marunga Mountains chimpanzee
<i>P. paniscus</i>	bonobo, bonobo, gracile or pygmy chimpanzee ; dwarf chimpanzee, lesser chimpanzee, pygmy chimpanzee

and combinations. Some authors have treated nouns attributively so as to inform us of the group of animals to which they refer (e.g., samango monkey or gelada baboon) while others see no need for this (e.g., samango or gelada). Other sources of diversity include alternative stem-words (galago or bush-baby, guenon or monkey), forms of qualifiers (Angola or Angolan, moustache or moustached), neologisms without a long history in the literature and often cited as alternative names (mitis monkey, neglectus monkey), abandoned names (colob), and possible unawareness that common names have already been provided (fire-bellied Wolf's monkey or red-bellied mona for the same taxon).

Zoologists have rejected some common names for animals as misleading or unsuitable because they do not conform to their chosen principles. Where it has seemed appropriate, they have provided new names. Primatologists could follow these precedents. Acting alone, collectively, or institutionally—for instance in a committee—they could recommend

which common names are to be quoted in publications on primates. From published sources they could select a leading common name for each species and recommend that it take precedence where a range of names is cited or that it is to be the sole name cited. Corbet and Hill (1991), R. W. Hayman's translation of Haltenorth and Diller (1980), Wilson and Cole (2000), Groves (2001, 2005), and Duff and Lawson (2004) have already selected single names for African primate species.

Common names for African primate subspecies have not been used very frequently. Authors have not published common names for all the subspecies to which they refer in systematic compilations (for example, Elliot 1913–1914; R. W. Hayman's translation of Haltenorth and Diller 1980; Napier 1981). Grubb *et al.* (2003) provided a complete list of subspecies names (Table 1). Some subspecies names are modified species names (*western potto*, *Kinda* yellow baboon), but if all were formed in this way, they could become too long (e.g.,

white-tailed small-eared greater galago). Other subspecies names in general use are not modified species names (e.g., white-naped mangabey, Moloney's monkey, roloway monkey). If subspecies names were altered so that they were all modified versions of species names, there would be adverse consequences. The numerous changes required would further burden the stock of published common names, and every alteration in taxonomic rank would require yet another name change.

Some authors (e.g., Duckworth and Pine 2003) prefer common names to be applied only to species, not to subspecies, on the grounds that the public may be misled or may draw false inferences about systematics, hampering their appreciation of taxonomy—subspecies could be thought to be species. However, there are positive reasons for using common names for subspecies. They could contribute to the conservation of these taxa by helping to make them more widely recognized by the public in popular articles, posters and films—as in the case of Miss Waldron's red colobus (see McGraw 2005). Using appropriate common names would more precisely identify primate taxa in contexts where scientific names of subspecies would not be appropriate. It is useful to have names for other subspecies (Table 1) should they be required for publicity material, for legends to illustrations, or for cases in which subspecies are raised to species status. If species were to be recognized as the smallest diagnosable assemblages of interbreeding organisms (Groves 2001), according to the phylogenetic-species definition (“phylogenetic species-concept”), many common names for subspecies would be needed as names for species.

Conclusions

The preferred names of species and subspecies in Table 1 and Grubb *et al.* (2003) could be regarded as a recommended list. However, conformity with this or any other list—if desired—can only be achieved through consensus. I do not wish to assert which names should be selected. Readers can see what common names have been used for African primates and, if they wish, decide which names they would prefer to use. They may not agree with some of the choices of recommended names or with some of the principles employed in their selection. They may choose different names out of the alternatives listed in Table 1 or they may wish to create some entirely new names. If the latter, it would be desirable if they explained their reasons and remembered that new names increase the load of vernacular nomenclature in the literature. More discussion on the common names to be adopted for primates is probably desirable. It would be interesting to know of the diversity of common names of primates in French, German, and other languages. To the extent that English is an international language of science, it would be desirable to achieve more stability for English-language common names.

Literature cited, including sources of primate names in Table 1.

- Anonymous. 1933. *Primates*. Trustees of the British Museum (Natural History), London.
- Alden, P. C., R. D. Estes, D. Schlitter and B. McBride. 1996. *Collins Guide to African Wildlife*. Harper Collins, London.
- Allen, G. M. 1939. A checklist of African mammals. *Bull. Mus. Comp. Zool. Harvard College* 83: 1–763.
- Ansell, W. F. H. 1978. *The Mammals of Zambia*. The National Parks and Wildlife Service, Chilanga, Zambia.
- Burgess, N. D., D. Kock, A. Cockle, C. FitzGibbon, P. Jenkins and P. Honess. 2000. Mammals. In: *Coastal Forests of Eastern Africa*, N. D. Burgess and G. P. Clarke (eds.), pp.173–190. IUCN – The World Conservation Union, Gland, Switzerland.
- Corbet, G. B. and J. E. Hill. 1991. *A World List of Mammalian Species*, 3rd edition. Natural History Museum Publications and Oxford University Press, London and Oxford.
- Dandelot, P. 1974. Order Primates, main text. In: *The Mammals of Africa: An Identification Manual*, Part 3, J. A. J. Meester and H. W. Setzer (eds.), pp.1–35. Smithsonian Institution Press, Washington, DC.
- Dorst, J. and P. Dandelot. 1970. *A Field Guide to the Larger Mammals of Africa*. Collins, London.
- Duckworth, J. W. and R. H. Pine. 2003. English names for a world list of mammals, exemplified by species of Indochina. *Mammal Rev.* 33: 151–173.
- Duff, A. and A. Lawson. 2004. *Mammals of the World. A Checklist*. A. & C. Black, London.
- Elliot, D. G. 1913–1914. *A Review of the Primates*. American Museum of Natural History, New York.
- Forbes, H. O. 1894. *A Hand-book to the Primates*. Edward Lloyd Ltd., London.
- Gray, J. E. 1870. *Catalogue of Monkeys, Lemurs, and Fruit-eating Bats in the Collection of the British Museum*. Trustees of the British Museum, London.
- Groves, C. P. 2001. *Primate Taxonomy*. Smithsonian Institution Press, Washington, DC.
- Groves, C. P. 2005. Order Primates. In: *Mammal Species of the World. A Taxonomic and Geographic Reference*, D. E. Wilson and D. M. Reeder (eds.), pp.111–184. Third Edition. Johns Hopkins University Press, Baltimore.
- Grubb, P., T. M. Butynski, J. F. Oates, S. K. Bearder, T. R. Disotell, C. P. Groves and T. T. Struhsaker. 2003. Assessment of the diversity of African primates. *Int. J. Primatol.* 24: 1301–1357.
- Haltenorth, T. and H. Diller [translated from the German by R. W. Hayman]. 1980. *A Field Guide to the Mammals of Africa Including Madagascar*. Collins, London.
- Hill, W. C. O. 1953. *Primates. Comparative Anatomy and Taxonomy. Strepsirhini*. University Press, Edinburgh.
- Hill, W. C. O. 1966. *Primates. Comparative Anatomy and Taxonomy. Catarrhini, Cercopithecidae, Cercopithecinae*. Edinburgh University Press, Edinburgh.

- Hill, W. C. O. 1969. The nomenclature, taxonomy and distribution of chimpanzees. In: *The Chimpanzee. Anatomy, Behavior and Diseases of Chimpanzees*, G. A. Bourne (ed.), pp. 22–49. Karger, Basel.
- Hill, W. C. O. 1970. *Primates. Comparative Anatomy and Taxonomy. Cynopithecinae*. Papio, Mandrillus, Theropithecus. University Press, Edinburgh.
- Hill, W. C. O. 1974. *Primates. Comparative Anatomy and Taxonomy. Cynopithecinae*. Cercocebus, Macaca, Cynopithecus. University Press, Edinburgh.
- Jenkins, P. D. 1987. *Catalogue of Primates in the British Museum (Natural History) and Elsewhere in the British Isles. Part IV: Suborder Strepsirrhini, Including the Subfossil Madagascan Lemurs, and Family Tarsiidae*. British Museum (Natural History), London.
- Kavanagh, M. 1984. *A Complete Guide to Monkeys, Apes and Other Primates*. Viking, New York.
- Kingdon, J. 1971. *East African Mammals. An Atlas of Evolution in Africa*. Vol. 1. Academic Press, London.
- Kingdon, J. 1997. *The Kingdon Field Guide to African Mammals*. Academic Press, London.
- Lee, P. C., J. Thornback and E. L. Bennett (compilers). 1988. *Threatened Primates of Africa. The IUCN Red Data Book*. IUCN – The World Conservation Union, Gland.
- Lydekker, R. 1893. *The Royal Natural History*. Vol. 1. Warne, London.
- Macdonald, D. (ed.). 1984. *The Encyclopaedia of Mammals*. George, Allen and Unwin, London.
- McGraw, W. S. 2005. Update on the search for Miss Waldron's red colobus monkey. *Int. J. Primatol.* 26: 605–619.
- Napier, P. H. 1981. *Catalogue of Primates in the British Museum (Natural History) and Elsewhere in the British Isles. Part II: Family Cercopithecidae, Subfamily Cercopithecinae*. British Museum (Natural History), London.
- Napier, P. H. 1985. *Catalogue of Primates in the British Museum (Natural History) and Elsewhere in the British Isles. Part III: Family Cercopithecidae, Subfamily Colobinae*. British Museum (Natural History), London.
- Napier, J. R. and P. H. Napier. 1967. *A Handbook of Living Primates*. Academic Press, London.
- Napier, J. R. and P. H. Napier. 1985. *The Natural History of the Primates*. British Museum (Natural History), London.
- Oates, J. F. 1996. *African Primates. Status Survey and Conservation Action Plan*. Revised Edition. IUCN, Gland, Switzerland.
- Rowe, N. 1996. *The Pictorial Guide to the Living Primates*. Pogonias Press, East Hampton, New York.
- Rylands, A. B., R. A. Mittermeier and W. R. Konstant. '2001' [2003]. Species and subspecies of primates described since 1990. *African Primates* 5: 51–54.
- Sanderson, I. T. 1957. *The Monkey Kingdom. An Introduction to the Primates*. Hamish Hamilton, London.
- Shortridge, G. C. 1934. *The Mammals of South West Africa*. William Heinemann, London.
- Smithers, R. H. N. and J. L. P. Lobão Tello. 1976. *Check List and Atlas of the Mammals of Mocambique*. Trustees of the National Museums and Monuments, Salisbury, Zimbabwe.
- Smithers, R. H. N. and V. J. Wilson. 1979. *Check List and Atlas of the Mammals of Zimbabwe Rhodesia*. Trustees of the National Museums and Monuments, Salisbury, Zimbabwe.
- Stuart, C. and T. Stuart. 1988. *Chris and Tylde Stuart's Field Guide to the Mammals of Southern Africa*. New Holland, London.
- Stuart, C. and T. Stuart. 1997. *Field Guide to the Larger Mammals of Africa*. Struik, Cape Town.
- Walker, E. P. 1964. *Mammals of the World*. Vol. 1. Johns Hopkins, Baltimore.
- Wilson, D. E. and F. R. Cole. 2000. *Common Names of Mammals of the World*. Smithsonian Institution Press, Washington, DC.
- Wolfheim, J. H. 1983. *Primates of the World. Distribution, Abundance, and Conservation*. University of Washington Press, Seattle.

Author's address:

Peter Grubb, 35, Downhills Park Road, London N17 6PE, United Kingdom. E-mail: <Pgrubb@aol.com>.

Received for publication: December 2004

Revised: March 2006