



## Corrigenda

Source: Zoological Science, 36(4) : 354-355

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/0289-0003-36.4.354>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Corrigenda

In the article “Interpopulation Comparison of Sex-Biased Mortality and Sexual Size Dimorphism in Sea-Run Masu Salmon, *Oncorhynchus masou*” by Tsuyoshi Tamate, which was published in *Zoological Science* 32(4): 364–371 (2015), a number of errors appeared in Tables 1 and 3.

Table 1. The sex ratios (female proportions) of age 1+ and age-combined smolts in the Moshosanbetsu 2010 cohort should be reversed. The correct values are 76.5(%) for age 1+ and 74.3(%) for age-combined.

Table 3. The fork length S.D.s of age-combined female adults for Moshosanbetsu 2010 and 2012 cohorts are incorrect. The correct values are 26 (mm) for 2010 and 24 (mm) for 2012.

The corrected Tables are shown below.

**Table 1.** Sex ratios (female proportions) of smolts and adults for each emigrating year (cohort) in each population. ‘Age-combined’ smolts include age 1+ and 2+ years smolts and ‘age-combined’ adults are composed of age 2+ and 3+ years adults and those with unknown age.

Population	Cohort	Smolts			Adults		
		Age	<i>n</i>	%	Age	<i>n</i>	%
Moshosanbetsu	2010	1+	34	76.5	2+	28	64.3
		age-combined	35	74.3	age-combined	30	66.7
	2011	1+	41	53.7	2+	21	85.7
		age-combined			age-combined	22	86.4
	2012	1+	77	63.6	2+	27	88.9
		age-combined			age-combined	28	89.3
Betsubetsu	2008	1+	86	70.9	2+	56	50.0
		age-combined	87	71.3	age-combined	57	50.9
	2009	1+	46	73.9	2+	53	60.4
		age-combined	48	75.0	age-combined	58	62.1
Okutsunai	2008	1+	48	93.8	2+	17	70.6
		age-combined	59	94.9	age-combined	21	76.2
	2009	1+	88	89.8	2+	17	70.6
		age-combined	89	89.9			

**Table 3.** Body sizes (fork length, FL) of female and male adults for each cohort in each population. 'Cohort' and 'age-combined' are defined in the caption in Table 1. Note, male FLs indicate 'adjusted FLs' considering their elongated snouts as a secondary sexual character (see text).

Population	Cohort	Age	Sex	<i>n</i>	Mean FL ± S.D. (mm)	
Moshosanbetsu	2010	2+	female	18	492 ± 28	
			male	10	407 ± 38	
	2011	2+	female	2	500, 504*	
			age-combined	female	20	493 ± 26
		2+	female	18	439 ± 38	
			male	3	393 ± 37	
	2012	2+	unknown	female	1	431
			age-combined	female	19	439 ± 37
		2+	female	24	466 ± 25	
			male	3	419 ± 18	
	2012	2+	unknown	female	1	483
			age-combined	female	25	467 ± 24
Betsubetsu		2008	2+	female	28	488 ± 38
				male	28	428 ± 42
	2008	2+	unknown	female	1	488
			age-combined	female	29	488 ± 37
2009	2+	female	32	494 ± 35		
		male	21	431 ± 45		
	2+	female	1	434		
		unknown	female	3	468 ± 43	
2009	2+	male	1	557		
		age-combined	female	36	491 ± 37	
	2+	male	22	437 ± 52		
		age-combined	female	36	491 ± 37	
Okutsunai	2008	2+	female	12	499 ± 25	
			male	5	446 ± 60	
		2+	female	2	527, 535*	
			unknown	female	2	477, 585*
	2009	2+	age-combined	female	16	507 ± 33
			female	12	490 ± 25	
		2+	male	5	401 ± 33	
			male	5	401 ± 33	

\*Individual data.