



Fish 3 β -Hydroxysteroid Dehydrogenase/ Δ 5- Δ 4Isomerase: Antibody Production and Their Use for the Immunohistochemical Detection of Fish Steroidogenic Tissues

Authors: Kobayashi, Tohru, Chang, X-T., Nakamura, Masaru, Kajiura, Hiroko, and Nagahama, Yoshitaka

Source: Zoological Science, 13(6) : 909-914

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.13.909>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Fish 3β -Hydroxysteroid Dehydrogenase/ Δ^5 - Δ^4 Isomerase: Antibody Production and Their Use for the Immunohistochemical Detection of Fish Steroidogenic Tissues

Tohru Kobayashi¹, X.-T. Chang^{1,2}, Masaru Nakamura³,
Hiroko Kajjura¹ and Yoshitaka Nagahama^{1,*}

¹Laboratory of Reproductive Biology, National Institute for Basic Biology,
Okazaki 444, Japan

²Molecular Biomechanics, School of Life Science,

The Graduate University for Advanced Studies, Okazaki 444, Japan

³Department of Biology, Teikyo University, Hachioji, Tokyo 192-03, Japan

ABSTRACT—We have produced polyclonal antibodies against two oligopeptides corresponding to middle and c-terminal regions of amino acid sequences predicted from rainbow trout (*Oncorhynchus mykiss*) 3β -hydroxysteroid dehydrogenase (3β -HSD) cDNA (Sakai *et al.*, 1994, FEBS Letters 350, 309-313). Both antibodies (α -tr 3β -M and α -tr 3β -C) recognized recombinant rainbow trout 3β -HSD protein derived from rabbit reticulocyte lysate system and non-steroidogenic mammalian COS-1 cell lysate. Immunoblot analysis of rainbow trout ovarian follicle homogenates revealed specific recognition of 3β -HSD protein. In rainbow trout testis, furthermore, immunoreactive 3β -HSD localized in Leydig cells in the interstitium of immature testes and interrenal cells in the head kidney. These results indicate that both α - 3β -HSD antibodies recognized rainbow trout 3β -HSD protein at the level of immunoblot and immunohistochemical analyses. Furthermore, both antibodies also recognized immunohistochemically 3β -HSD in various steroidogenic organs (ovary, testis, and interrenal glands) of several teleost fishes.

INTRODUCTION

The enzyme 3β -hydroxysteroid dehydrogenase/isomerase (3β -HSD) is essential for the biosynthesis of most steroid hormones. 3β -HSD catalyzes pregnenolone, 17α -hydroxypregnenolone, dehydroepiandrosterone and androstendiol to progesterone, 17α -hydroxyprogesterone, androstenedione and testosterone, respectively. Recently, recombinant 3β -HSD proteins from several mammalian species have been characterized (human, Lorence *et al.*, 1990a, b; mouse, Bain *et al.*, 1990; rat, Lorence *et al.*, 1991). Dynamics of 3β -HSD protein expression were examined using 3β -HSD antibody against purified human placental 3β -HSD protein (Clarke *et al.*, 1993a, b).

Although numerous studies of steroid metabolism have been reported (see Nagahama, 1987), analysis of 3β -HSD protein in lower vertebrates has little attention due to the lack of the specific 3β -HSD antibodies. Recently, Sakai *et al.* (1994) cloned rainbow trout 3β -HSD cDNA and consequently the derived amino acid sequence allows production of specific antibodies to help clarify the role and function of 3β -HSD in

fish. This study determines the specificity of 3β -HSD antibodies produced and their use for the immunohistochemical detection of fish steroidogenic tissues.

MATERIALS AND METHODS

Animals

Rainbow trout (*Oncorhynchus mykiss*) were obtained from the Aichi Prefectural Fisheries Station, Toyokawa, Japan. These animals were maintained in the laboratory until use.

Production of polyclonal antibodies

The middle portion sequence (CTCALRPMYIYGEC: M) with an additional cysteine in the N- and C-terminus and C-terminal sequence (CTMDWVASQLPKERERIKV: C) in amino acid sequence with an additional cysteine in the N-terminal sequence predicted from rainbow trout 3β -HSD cDNA (Sakai *et al.*, 1994) were synthesized by the Fmoc protocol on an Applied Biosystem model 431A peptide synthesizer, and purified by reversed-phase HPLC using a ODS-5 column (Develosil). To increase antigenicity, these peptides were coupled to bovine serum albumin (BSA: Fraction V, Sigma) or Keyhole limpet hemocyanin (KLM: Calbiochem), using EMCS (N-(ϵ -Maleimidocaproyloxy)succinimide) (Dojindo), following the cleavage of disulfide bonding within the molecules of BSA and KLM with dithiothreitol (DTT).

Female rabbits were immunized at 2-week intervals by four

* To whom all correspondence should be addressed.

subcutaneous injections of peptide-linked protein (1 mg/rabbit/injection). These antigens were emulsified in Freund's complete adjuvant at the first injection and in Freund's incomplete adjuvant after the first injection. One week after the last injection, whole blood from rabbit was collected. From collected blood, serum was separated and then purified by affinity chromatography using synthetic peptides used as antigens.

Preparation of recombinant 3 β -HSD protein

To obtain 3 β -HSD protein, we tried to produce recombinant 3 β -HSD protein using rabbit reticulocyte lysate system (Promega) and COS-1 cells transfected with rainbow trout 3 β -HSD cDNA.

For rabbit reticulocyte lysate system, we used rainbow trout 3 β -HSD cDNA ligated to pBluescriptII SK(-) as template. According to the instruction manual, lysate containing recombinant trout 3 β -HSD protein was obtained and then this lysate was treated with Laemmli's SDS sample buffer (Laemmli, 1970) for immunoblot analysis. For expression of rainbow trout 3 β -HSD in COS-1 cells, the rainbow trout 3 β -HSD expression vector was constructed by ligating the blunt-ended cDNA fragment into the *Sma*I site of pSVL (Pharmacia LKB). Transfection of rainbow trout 3 β -HSD cDNA construct to COS-1 cells was carried out as described previously (Sakai *et al.*, 1994). After this, transfected COS-1 cells were recovered and then homogenized in 0.25 M sucrose, 20 mM Hepes (pH 7.5). To demonstrate whether 3 β -HSD activity was in these homogenates, a part of these homogenates was applied to steroid metabolism experiments as described previously (Kobayashi *et al.*, 1993), and the other was treated with SDS-sample buffer for immunoblot analysis.

Protein extraction and electrophoresis

To obtain native 3 β -HSD proteins, proteins from testis and interrenal glands were extracted as a mitochondria and microsomal fraction after ultracentrifugation (100,000 g, 1 hr, 4°C), then frozen in liquid nitrogen and stored at -80°C until use. Also ovarian follicles were frozen in liquid nitrogen and stored at -80°C. For electrophoresis, proteins from testis and interrenal glands and ovarian follicles were treated with Laemmli's SDS-sample buffer containing 10% β -mercaptoethanol, for 3 min at 100°C, and analyzed by SDS-PAGE with 12.5% gel (Laemmli, 1970).

Immunoblotting

Proteins separated by SDS-PAGE were transferred to Immobilon membrane (Millipore) by electroblotting (Towbin *et al.*, 1979). The membrane was rinsed in Tris-buffered saline (TBS: 20 mM Tris-HCl, 150 mM NaCl, pH 7.5), blocked with non-fat dry milk in TBS containing 0.1% Tween 20 (TTBS). After washed three times (5 min each) with TTBS, the membrane was incubated with a 1:1000 dilution of serum for 2 hr. After washing three times (5 min each) with TTBS, the membrane was incubated with a 1:1000 dilution of alkaline phosphatase-conjugated goat anti-rabbit IgG (Tago). Following further three washes with TTBS, phosphatase activity was visualized by treating the membrane with 0.2 mM 5-bromo-4-chloro-3-indolylphosphate p-toluidine salt and nitroblue tetrazolium (Sigma) in 100 mM diethanolamine buffer (pH 9.5) containing 5 mM MgCl₂. All incubations were performed at room temperature.

Immunohistochemistry

Ovary, testis and interrenal glands were dissected from rainbow trout, tilapia (*Oreochromis niloticus*), goldfish (*Carassius auratus*) and Japanese eel (*Anguilla japonica*), then fixed in Bouin's fixative solution and embedded in paraffin. Serial cross sections were cut at 6 μ m. The antibodies were used at 1:1600 dilution. The procedure of this immunohistochemistry in details was described in a previous report (Kobayashi and Iwasawa, 1992).

RESULTS

Immunoblotting with anti-tr3 β -M and tr3 β -C antibodies

To characterize anti-tr3 β -M and tr3 β -C antibodies, first, proteins extracted from rainbow trout ovarian follicles were immunoblotted. As shown in Fig. 1, both antibodies recognized specifically 45 kDa band. This immunoreactive 45 kDa band was disappeared after adsorption of each peptide coupled with carrier protein (data not shown). Next, we examined whether these antibodies could detect recombinant 3 β -HSD protein. Rabbit reticulocyte lysate system using rainbow trout 3 β -HSD cDNA as complement produced only 45 kDa protein (Fig. 2a). Immunoblot analysis demonstrated that recombinant 3 β -HSD protein was recognized by both anti-tr3 β -M and anti-tr3 β -C antibodies (Fig. 2b). COS-1 cell lysates transfected with rainbow trout 3 β -HSD cDNA produced a bioactive 3 β -HSD protein which was also detected by these antibodies (Fig. 3a, b). These results indicated that the anti-tr3 β -M and tr3 β -C antibodies specifically recognized 3 β -HSD protein from rainbow trout.

To demonstrate whether the multiple forms of 3 β -HSD are present, immunoblot analysis of several steroidogenic organs (i.e., testis, ovary and interrenal glands) from rainbow trout has completed. As shown in Fig. 4, a single and immunoreactive protein of equivalent size was detected in all organs tested.

Immunohistochemistry with anti-tr3 β -M and tr3 β -C antibodies

To examine the availability of anti-tr3 β -M and tr3 β -C antibodies for immunohistochemistry, we applied these antibodies to immature and mature testes, immature ovaries, and interrenal glands of several teleost fishes including rainbow trout (Fig. 5a-f). In rainbow trout, immunoreactive 3 β -HSD




Fig. 1. Immunoblot analysis of ovarian follicle homogenates from rainbow trout with α -tr3 β -M and tr3 β -C antibodies. For SDS-PAGE, 5 μ g of protein from each sample was applied. Both antibodies recognized specifically 45 kDa band.




Fig. 2. Immunoblot analysis of recombinant rainbow trout 3 β -HSD protein with α -tr3 β -M and tr3 β -C antibodies. Rabbit reticulocyte lysate system was used for the production of 3 β -HSD protein. a, Incorporation of 35 S-methionine at production of rainbow trout 3 β -HSD protein. SK-tr3 β , rainbow trout 3 β -HSD cDNA ligated to pBluescriptII SK(-). b, Immunoblot analysis of recombinant rainbow trout 3 β -HSD protein with α -tr3 β -M and tr3 β -C antibodies. For SDS-PAGE, 5 μ g of protein from each sample was applied.

localized in Leydig cells in the interstitium of immature testes (Fig. 5a) and interrenal cells in the head kidney (Fig. 5d). These immunoreactive cells have typical steroidogenic features such as smooth endoplasmic reticulum, mitochondria with tubular cristae and lipid droplets (data not shown). Immunoreactive 3 β -HSD was also found in Leydig cells of the mature testis (Fig. 5b) and interstitial cells of the immature ovary of tilapia (Fig. 5c); a weak immunoreaction was found in the cytoplasm of immature oocytes. Positive staining for 3 β -HSD was also detected in interrenal cells of goldfish (Fig. 5e) and Japanese eel (Fig. 5f). There was no immunostaining associated with non-steroidogenic cell types such as chromaffin cells, endothelial cells and blood cells.

DISCUSSION

We described the characteristics of anti-tr3 β -M and tr3 β -C antibodies that recognized 3 β -HSD proteins and the localization of 3 β -HSD proteins in ovary, testis and interrenal glands. To our knowledge, this report is the first examination on the localization of 3 β -HSD using homologous 3 β -HSD antibodies in nonmammalian vertebrates.

Previous reports indicated that multiple forms of 3 β -HSD proteins are present in mammals. Although immunoblot analyses were performed by two distinct antibodies, the present study indicated a single immunoreactive 3 β -HSD protein. However, other forms of 3 β -HSD protein in rainbow trout may not be recognized by these antibodies. In contrast to the multiple related 3 β -HSD isoenzymes in human (Lorence




Fig. 3. Immunoblot analysis of recombinant rainbow trout 3 β -HSD protein with α -tr3 β -M and tr3 β -C antibodies. COS-1 cell lysates transfected with rainbow trout 3 β -HSD cDNA. a, Bioactivity of COS-1 cell lysates transfected with rainbow trout 3 β -HSD cDNA. To determine 3 β -HSD activity of transfected COS-1 cell lysates, dehydroepiandrosterone (DHEA) was used as substrate. The extracted steroid metabolites were applied on thin layer chromatography with benzene : acetone (4 : 1). And, Androstenedione. b, Immunoblot analysis of COS-1 cell lysates transfected with rainbow trout 3 β -HSD cDNA. For SDS-PAGE, 5 μ g of protein from each sample was applied.




Fig. 4. Immunoblot analysis of several steroidogenic tissues from rainbow trout with α -tr3 β -M and tr3 β -C antibodies. Testis, lysates of mitochondria and microsome-rich fractions from testis; Ovary, homogenates of ovarian follicles from ovary; Interrenal, lysates of mitochondria and microsome-rich fractions from head kidney. For SDS-PAGE, 5 μ g of protein from each sample was applied.

et al., 1999a, b), rat (Naville *et al.*, 1991; Zhao *et al.*, 1990, 1991) and mouse (Bain *et al.*, 1990), Sakai *et al.* (1994) indicated that a single gene encoded 3 β -HSD in rainbow trout from Southern hybridization analysis of genomic DNA in rainbow trout. Thus, these results suggest that rainbow trout 3 β -HSD protein is not multiform.

Although numerous studies on the histochemical detection of 3 β -HSD and 3 β -HSD activity at the level of steroid metabolism were reported (Lofts and Bern, 1972; Guraya, 1976; Nagahama, 1987), there is a little information in lower vertebrates on the localization of 3 β -HSD protein in steroidogenic tissues at the level of immunohistochemistry and immunoblotting. In the present study, both 3 β -HSD antibodies recognized immunohistochemically 3 β -HSD in immature ovaries, immature and mature testes, and interrenal glands of several teleost fishes. In the immature ovary of tilapia, strong immunostaining for 3 β -HSD was observed in the interstitial cells which had previously been shown to be steroidogenic by ultrastructural observations (Nakamura *et al.*, 1993). A weak 3 β -HSD immunoreaction was also seen in the cytoplasm of immature oocytes of tilapia. Further biochemical studies are necessary to determine whether tilapia oocytes are steroidogenic. Leydig cells in both immature and mature testes were positive to 3 β -HSD antibodies. These findings are in agreement with previous histochemical and ultrastructural observations on the Leydig cells of several teleost fishes (Nagahama *et al.*, 1978; Nagahama, 1987). Immunoreactive 3 β -HSD in Sertoli cells was not detected in the present study, though some previous reports suggested the presence of 3 β -HSD activity in Sertoli cells (*Cymatogaster aggregata*, Wiebe, 1969; *Fundulus heteroclitus*, Bara, 1969; *Salmo gairdneri*, van den Hurk *et al.*, 1978a, b). Further studies are necessary whether Sertoli cells are steroidogenic in fish. In the present

study the immunohistochemical localization of 3 β -HSD in the head kidney was confined only to the interrenal cells; immunoreaction was not observed in either chromaffin cells or blood cells. These findings are consistent with previous results for the restricted distribution of 3 β -HSD in interrenal cells in the teleost head kidney (Hanke and Chester Jones, 1966; Lofts and Bern, 1972; Kagawa and Nagahama, 1989).

In the present study, we demonstrated that rainbow trout anti-tr3 β -M and tr3 β -C antibodies which could detect the 3 β -HSD protein specifically were available for immunoblot analysis and immunohistochemistry. Recently we obtained the results that the tr3 β -M-antibody was able to recognize the 3 β -HSD protein in many animals including mammals, birds, reptiles and amphibians (Kobayashi *et al.*, unpublished). Thus it seems that these antibodies are also available for detection of 3 β -HSD protein in steroidogenic tissues throughout vertebrates.

ACKNOWLEDGEMENTS

We thank Dr. M. Yamashita, Hokkaido University for technical advice to produce antibodies. This work was supported in part by Grant-in-Aid for Scientific Research on Priority Areas from the Ministry of Education, Science, Sports and Culture of Japan (07283104), and by a Grant-in-Aid (Bio Media Program) from the Ministry of Agriculture, Forestry and Fisheries (BMP 95-II-2-6). Research support by Grants from the Mitsubishi Foundation and the Uehara Memorial Foundation is also acknowledged.

REFERENCES

- Bain PA, Yoo M, Clarke T, Hammond SH, Payne AH (1990) Multiple forms of mouse 3 β -hydroxysteroid dehydrogenase/ Δ^5 - Δ^4 isomerase and differential expression in gonads, adrenal glands, liver, kidney of both sexes. *Proc Natl Acad Sci USA* 88: 8870–8874
- Bara G (1969) Histochemical demonstration of 3 β -, 3 α -, 11 β -, 17 β -hydroxysteroid dehydrogenases in the testis of *Fundulus heteroclitus*. *Gen Comp Endocrinol* 13: 189–200
- Clarke TR, Bain PA, Sha L, Payne AH (1993a) Enzyme characteristics of two distinct forms of mouse 3 β -hydroxysteroid dehydrogenase/ Δ^5 - Δ^4 -isomerase complementary deoxyribonucleic acids expressed in COS-1 cells. *Endocrinology* 132: 1971–1976
- Clarke TR, Bain PA, Greco TL, Payne AH (1993b) A novel mouse kidney 3 β -hydroxysteroid dehydrogenase complementary DNA encodes a 3-ketosteroid reductase instead of a 3 β -hydroxysteroid dehydrogenase/ Δ^5 - Δ^4 -isomerase. *Mol Endocrinol* 7: 1569–1578
- Guraya SS (1976) Recent advances in the morphology, histochemistry, biochemistry of steroid-synthesizing cellular sites in the nonmammalian vertebrate ovary. *Int Rev Cytol* 44: 365–409
- Hanke W, Chester Jones I (1966) Histological and histochemical studies on the adrenal cortex and the corpuscles of Stannius of the European eel (*Anguilla anguilla*). *Gen Comp Endocrinol* 7: 166–178
- Kagawa H, Nagahama Y (1980) Ultrastructural localization of Δ^5 -3 β -hydroxysteroid dehydrogenase in the interrenal cells of the goldfish (*Carassius auratus*). *Cell Tissue Res* 212: 225–231
- Kobayashi T, Iwasawa H (1992) Timing of proliferation of spermatogonia and spermatogonia-supporting Sertoli cells in the newt *Cynopus pyrrhogaster*. *Biomed Res* 13: 167–172
- Kobayashi T, Sakai N, Adachi S, Asahina K, Iwasawa H, Nagahama Y (1993) 17 α -, 20 α -dihydroxy-4-pregnen-3-one is the naturally occurring spermiation-inducing hormone in the testis of a frog, *Rana nigromaculata*. *Endocrinology* 133: 321–327




Fig. 5. Immunohistochemistry of several steroidogenic tissues with 3 β -HSD antibodies. a, Immature testis from young rainbow trout was stained with α -tr3 β -M antibody. Seminiferous tubules were filled with spermatogonia and poorly-developed Leydig cells. b, Mature testis from tilapia was stained with α -tr3 β -M antibody. Note strong immunoreaction for 3 β -HSD in only Leydig cells (a and b). ED, intratesticular efferent duct. c, Immature ovary from young tilapia was stained with tr3 β -C antibody. 3 β -HSD positive cells are found in the interstitium. d, Interrenal gland in the head kidney from female rainbow trout was stained with tr3 β -C antibody. e, Interrenal gland in the head kidney from female goldfish was stained with tr3 β -C antibody. CV, cardinal vein. f, Interrenal gland in the head kidney from male Japanese eel was stained with α -tr3 β -M antibody. Note strong immunoreactive 3 β -HSD in interrenal cells (d, e and f). \times 270.

Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680–685

Lofts B, Bern HA (1972) The functional morphology of steroidogenic tissues. In "Steroids in Nonmammalian Vertebrates" Ed by Idler DR, Academic Press, New York, pp 37–126

Lorence MC, Murry BA, Trant JM, Mason JI (1990a) Human 3 β -hydroxysteroid dehydrogenase/ Δ^5 - Δ^4 -isomerase from placenta: expression in non-steroidogenic cells of a protein that catalyzes

the dehydrogenation/isomerization of C21 and C19 steroids. *Endocrinology* 126: 2493–2498

Lorence MC, Corbin CJ, Kamimura N, Mahendroo MS, Mason JI (1990b) Structural analysis of the gene encoding human 3 β -hydroxysteroid dehydrogenase/ Δ^5 - Δ^4 -isomerase expression in rat and characterization of the testis isoform. *Mol Cell Endocrinol* 80: 21–31

Lorence MC, Naville D, Graham-Lorence SE, Mack SO, Murry BA, Trant JM, Mason JI (1991) 3 β -hydroxysteroid dehydrogenase/

- Δ^5 - Δ^4 isomerase expression in rat and characterization of the testis isoform. *Mol Cell Endocrinol* 80: 21–32
- Nagahama Y, Clarke WC, Hoar WS (1978) Ultrastructure of putative steroid-producing cells in the gonads of coho (*Oncorhynchus kisutch*) and pink salmon (*Oncorhynchus gorbuscha*). *Can J Zool* 56: 2508–2519
- Nagahama Y (1987) Testis. In "Vertebrate Endocrinology: Fundamentals and Biochemical Implications Vol 1" Ed by Pang PKT, Schreibman MP, Gorbman A, Academic Press, New York, pp 399–438
- Nakamura M, Specker JL, Nagahama Y (1993) Ultrastructural analysis of the developing follicle during early vitellogenesis in tilapia, *Oreochromis niloticus*, with special reference to the steroid-producing cells. *Cell Tissue Res* 272: 33–39
- Naville D, Keeney DS, Jenkin G, Murry BA, Head JR, Mason JI (1991) Regulation of expression of male-specific rat liver microsomal 3 beta-hydroxysteroid dehydrogenase. *Mol Endocrinol* 5: 1090–1100
- Sakai N, Tanaka M, Takahashi M, Fukada S, Mason JI, Nagahama Y (1994) Ovarian 3 β -hydroxysteroid dehydrogenase/ Δ^5 - Δ^4 -isomerase of rainbow trout: its cDNA cloning and properties of the enzyme expressed in a mammalian cell. *FEBS Lett* 350: 309–313
- Towbin H, Staehelin T, Gordon J (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: Procedure and some applications. *Proc Natl Acad Sci USA* 76: 4350–4354
- van Den Hurk R, Peute J, Vermeij JAJ (1978a) Morphological and enzyme cytochemical aspects of the testis and vas deferens of the rainbow trout, *Salmo gairdneri*. *Cell Tissue Res* 186: 309–325
- van Den Hurk R, Vermeij JAJ, Stegenga J, Peute J, van Oordt PGWJ (1978b) Cyclic changes in the testis and vas deferens of the rainbow trout *Salmo gairdneri* with special reference to sites of steroidogenesis. *Ann Biol Anim Biochim Biophys* 18: 899–904
- Wiebe JP (1969) Endocrine controls of spermatogenesis and oogenesis in the viviparous sea perch *Cymatogaster aggregata* Gibbons. *Gen Comp Endocrinol* 12: 267–275
- Zhao H-F, Rheaume E, Trudel C, Couet J, Labrie F, Labrie F (1990) Structure and sexual dimorphic expression of a liver-specific rat 3 β -hydroxysteroid dehydrogenase/isomerase. *Endocrinology* 127: 3237–3239
- Zhao H-F, Labrie C, Simard J, de Lanuoi Y, Trudel C, Martel C, Rheaume E, Dupont E, Luu-The V, Pelletier G, Labrie F (1991) Characterization of rat 3 β -hydroxysteroid dehydrogenase/ Δ^5 - Δ^4 isomerase and different expression of the corresponding mRNAs in steroidogenic and peripheral tissues. *J Biol Chem* 266: 583–593

(Received February 29, 1996 / Accepted August 4, 1996)