



GnRH Antagonist Effects on Follicle Number and Size in Rat Neonates and Infants

Authors: Muir, Tristi W., Leach, Richard E., Roche, Patrick C., Gaffey, Thomas A., Kuehl, Thomas J., et al.

Source: Zoological Science, 16(2) : 299-302

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.16.299>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

GnRH Antagonist Effects on Follicle Number and Size in Rat Neonates and Infants

Tristi W. Muir^{1,2}, Richard E. Leach³, Patrick C. Roche¹, Thomas A. Gaffey¹,
Thomas J. Kuehl^{2*}, W Richard Dukelow^{4**} and Steven J. Ory¹

¹*Department of Obstetrics and Gynecology and Pathology, Mayo Medical Center,
Rochester, Minnesota, USA,*

²*Department of Obstetrics and Gynecology, Pathology and Medical Biochemistry and Genetics,
Scott and White Clinic and Memorial Hospital, Texas A and M University Health
Science Center, College of Medicine, Temple, Texas, USA,*

³*Department of Obstetrics and Gynecology, Wayne State University, Detroit,
Michigan, USA and*

⁴*Endocrine Research Center, Michigan State University,
East Lansing, MI 48824, USA*

ABSTRACT—Increased levels of gonadotropins in the perinatal and prepubertal period may be responsible for the rapid phase of concurrent follicular atresia. This study tests the hypothesis that follicular atresia during this period can be reduced by suppressing gonadotropin release with a GnRH antagonist. Female rat litter mates were randomized to receive daily injections of GnRH_i (100 µg Detirelix® [Syntex, Palo Alto, CA] from the day of birth and were sacrificed at 5, 15, or 26 days of life. Follicular atresia was assessed by measuring number and size distribution of ovarian follicles. Serum FSH levels were assayed. GnRH_i treatment significantly depressed serum FSH and decreased the number of large antral follicles in 26 day rats, while body weight, reproductive tract weight and total follicle number per representative section were not significantly altered. Age-related changes were significant for all variables. The loss of primordial follicles is likely the result of another mechanism or combination of mechanisms. Gonadotropins do not appear to play a major role in follicular atresia in the neonatal and infant rat.

INTRODUCTION

In Mammals, the number of oocytes declines throughout life as a result of recruitment of primordial follicles and atresia. When the quantity of oocytes reaches a number that is no longer responsive to gonadotropins, ovarian cycling stops and, in menstruating species, menopause begins (Speroff *et al.*, 1989). The cycle of follicular recruitment, selection, and atresia begins in fetal life. In fetal and prepubertal stages, full follicular maturity, expressed by ovulation, does not occur. However, the decline in oocyte number by atresia is dramatic.

Atresia may occur at any point on a continuum of follicular development, from the initial recruitment of the primordial follicle to the development of the large, preovulatory follicle (Speroff *et al.*, 1989). Interest has been directed towards understanding oocyte atresia in an attempt to control follicular function for protection from toxic entities (such as chemotherapeutics) and for the prolongation of reproductive potential

(Ataya *et al.*, 1985; Jarrell, *et al.*, 1987; Ataya *et al.*, 1989).

Pulsatile secretion of GnRH from the hypothalamus leads to LH and FSH secretion from the pituitary. In the adult ovary, the number of follicles that undergo maturation is dependent upon the amount of FSH and LH that is available to the ovary and the sensitivity of the ovary to the gonadotropins. GnRH acts directly in the rat ovary to trigger meiotic maturation of the ova within preantral follicles (Banka and Erickson, 1985). GnRH therefore, plays an important role in the regulation of follicular maturation and ovulation.

GnRH antagonists act by direct occupation of GnRH receptors on the pituitary, blocking the release of gonadotropins. The GnRH agonists act by depleting hormone stores (resulting in an initial serum elevation of gonadotropins) followed by rapid internalization of the receptors and depletion of available GnRH surface receptors (Wierman *et al.*, 1989). GnRH antagonists have been found to effectively suppress LH and FSH serum levels (Wierman *et al.*, 1989; Dvorshak-Harvey *et al.*, 1985; Zeleznik and Kubik, 1986; van den Dungen *et al.*, 1989; Wierman and Wang, 1990; Sharpe *et al.* 1990; Meijs-Roelofs *et al.*, 1990; Andreyko *et al.*, 1992), while avoiding the initial elevation in gonadotropin levels before desensitization.

* Corresponding author for reprint requests

** Present address: High Meadows Enterprises 325 Spring Creek Road Somers, MT 59932, USA. Tel. (406) 857-2157, E-mail: dukelow@digisys.net

tization, which is characteristic of the GnRH agonist.

The rate of atresia of primordial follicles in the rat is greatest during the fetal through infant periods and continues at a slower rate during the adult life (Lintern-Moore, 1977). In the neonatal period of the female rat, a high serum FSH level and premature, preovulatory type LH peaks are present during days 9–21 of life (van den Dungen *et al.*, 1989; Goldman *et al.*, 1971; Ojeda and Ramirez, 1972; Doherty and Wuttke, 1976). These peaks occur during a period of substantial loss of follicles—between birth and 20 days of life (Lintern-Moore, 1977). Treatment of young cycling rats with a GnRH agonist inhibited the process of follicular recruitment, resulting in the preservation of a large number of small follicles which would have been lost in the physiologic process of recruitment and atresia (Ataya *et al.*, 1989).

A GnRH agonist was examined for effects on follicular atresia in neonatal and infant rats (Ory *et al.*, 1990). While FSH levels were found to be significantly decreased by day 15 of life, no significant difference in follicular size or number between treated and untreated animals was present at day 25. The early agonist release of gonadotropins may have obscured the effects of GnRH suppression in this study.

To test the hypothesis that the rate of follicular atresia could be slowed during the neonatal period by suppression of serum gonadotropins, administration of a GnRH antagonist to rat pups and its effects on the number and size of follicles were examined.

MATERIALS AND METHODS

Pregnant Sprague-Dawley rats were fed rat Chow[®] and water ad libitum and housed in an environmentally controlled room in a 14 hr light, 10 hr dark cycle (20–22°C). The afternoon following delivery, female pups were randomly assigned to either control or treatment groups. The pups were identified with an ear punch and remained with their littermates. Male pups were removed from each litter. Rat pups in the treatment group were injected with a GnRH antagonist (100 µg Detirelix[®], Syntex, Palo Alto, CA, in 0.1 ml of vehicle consisting of 50% propylene glycol and 50% normal saline) subcutaneously every 24 hr (Ory *et al.*, 1990) commencing within 24 hours of birth. The control rats were injected with vehicle only.

To examine the effects of GnRH antagonist on follicular atresia, rat pups were sacrificed by cervical dislocation 5, 15, or 26 days after birth. The upper reproductive tracts (oviducts and ovaries) were identified, rinsed with normal saline, weighed, fixed in 10% buffered formalin, set in paraffin, and serial step-sectioned (6 µm thickness). Sections from both ovaries were mounted on slides. The slides were stained with hematoxylin and eosin. The sections were viewed and a section representing the mid-longitudinal axis of the ovary was selected from one ovary of each animal. The follicles with a germinal vesicle visible in the oocyte were counted and classified according to grades of maturation using a Zeiss microscope. Grade I follicles were defined as primordial follicles with a small oocyte in prophase of the first meiotic division and a single layer of granulosa cells. Grade II follicles displayed an oocyte of increased diameter surrounded by a single layer of granulosa cells. Grade III follicles were defined as those with multiple layers of granulosa cells and an increased oocyte diameter. Grade IV follicles displayed an antrum. In addition, every serial section was morphologically evaluated using a previously published method (Stachecki *et al.*, 1994). Ovary sections were visualized on an Olympus BH-2 microscope at 25X magnification. Each follicle was

digitized as a video image using a VSP SC series high resolution camera Model SC505 (Applied Intelligence Systems, Ann Arbor, MI) which connected to a Dell 486 SX computer and was analyzed using MCID M1 image analysis software (Version 4.12; Imaging Research, St. Catherine's, Ontario, Canada). The image field was calibrated in micrometers with a stage micrometer at the same magnification prior to any measurements. Area, perimeter and maximum chord diameter were measured for each follicle. Measurements were obtained by tracing the outer edge of a follicle using a computer mouse (assuming spherical geometry). Areas were recorded in square microns.

To document FSH suppression in the GnRH antagonist treated rats, blood was collected by cardiac aspiration from sacrificed rats and stored at –70°C. The FSH levels on Day 5, 15, and 26 samples were determined by NIH radioimmunoassay (FSH RIA kit NIHDDK, NIH Bethesda, MD).

Data were analyzed using two-way ANOVA with treatment and age of assessment as the factors. Student-Newman-Keuls post-hoc testing was used to compare group means. Nonparametric methods, such as Friedman's analysis of ranked data, were used when necessary to meet assumptions of normality and equal variance. $P < 0.05$ was taken as significant.

RESULTS

Measuring circulating FSH levels assessed the effect of GnRH antagonist treatment on pituitary release of gonadotropins. Treatment with GnRH antagonist significantly ($P < 0.001$) decreased serum FSH levels at all age points (5, 15, 26 days) as compared to those of control animals (Fig. 1). The effect was not related to an adverse effect of GnRH antagonist on animal health and growth as assessed by comparing animal




Fig. 1. FSH blood levels (means \pm SE) in control (N = 5,4,8) and GnRH antagonist treated (N = 3, 4, 6) rats at 5, 15, and 26 days of life, respectively.

body and organ weights. There were no significant differences in upper reproductive tract weights and total body weights between the control and treated animals ($P = 0.41$ and 0.38 ,




Fig. 2. Numbers (means \pm SE) of follicles in a representative ovarian section in control and GnRH antagonist treated rats at 5, 15, and 26 days of life.




Fig. 3. Numbers (means \pm SE) of grade I to IV follicles in control and GnRH antagonist treated rats at 5, 15, and 26 days of life.

respectively). Age related differences in weights are apparent between the 5, 15, and 26 day groups. A discrepancy in sample sizes occurred between the data set for FSH levels (Fig. 1). Fewer animals at 5 and 15 days of age were assayed for FSH level due to difficulty in obtaining sufficient amounts of blood from the smaller animals.

There was not a significant difference between the control and treated animals ($P = 0.77$) in the total number of ovarian follicles in representative 6 micron sections. There were age-related changes in the number of ovarian follicles ($P < 0.001$), confirming the significance of atresia in the infant and juvenile period (Fig. 2). There were no significant differences in the number of grade I, II, and III follicles between control and treated animals (Fig. 3). The control animals exhibited more grade IV follicles ($P < 0.05$) than the treated animals (Fig. 3), likely in response to the availability of gonadotropins (Fig. 1). There is a trend towards an increase in the mean follicular area in the control animals (Fig. 4). However, due to the small number of antral follicles in comparison to primordial follicles, there is no statistical difference between the two groups.




Fig. 4. Follicular area (means \pm SE) in control and GnRH antagonist treated rats at 5, 15, and 26 days of life.

DISCUSSION

The orchestration of the process of follicular atresia in the prepubertal period is a mystery. Gonadotropin surges have been shown to occur during the neonatal and infant periods, coinciding with a dramatic decrease in follicle number in the rat (van den Dungen *et al.*, 1989; Lintern-Moore, 1977; Goldman *et al.*, 1971; Ojeda and Ramirez, 1972; Dohter and Wuttke, 1976). However, we observed that inhibition of gona-

dotropin secretion with a GnRH antagonist fails to alter the reduction in number of ovarian follicles in the prepubertal rat. Ataya *et al.* (1985) demonstrated that the process of follicular recruitment could be inhibited by a GnRH agonist in young cycling rats. Taken together, these observations suggest that either the ovarian response varies due to the differing nature of the GnRH analogues, or the process of follicular recruitment and atresia in the prepubertal rat is the result of a different mechanism than in the cycling rat.

Atresia in the neonatal and juvenile rat could result from a loss of oocytes directly through the surface epithelium of the ovary, degeneration of large follicles, or as a result of a preprogrammed event (Lunenfeld *et al.*, 1975). These events could be dependent upon other events occurring in utero and not upon neonatal factors or the postnatal environmental factors, or perhaps, as combination of these mechanisms is responsible for atresia in the rat.

During the prepubertal period follicles may grow to a large size, but the hormonal environment does not support the development of an ovulatory follicle (Lunenfeld *et al.*, 1975). These follicles likely undergo atresia through degeneration and reabsorption. Gonadotropins, therefore, may play a role in atresia as it relates to the development and degeneration of large follicles. There was a significantly greater number of antral follicles (grade IV) in the 26 day old control animals, but the number of large follicles contributes minimally to the total follicle population. Therefore, degeneration of large follicles may be reduced with suppression of gonadotropins, but this reduction is not significant in the process of atresia. The loss of primordial follicles is likely the result of another mechanism or combination of mechanisms. Gonadotropins do not appear to play a major role in follicular atresia in the neonatal and infant rat.

ACKNOWLEDGEMENTS

Financial support for this project provided by intramural sources at Mayo Medical Center. Partial support was received under NIEHS Grant ES004911. The authors acknowledge Dr. Scott Monroe's assistance in providing the GnRH antagonist, Detirelex, and Kate Kitzmann's assistance in performing FSH assays.

REFERENCES

- Andreyko IL, Monroe SE, Marshall LA, Fluker MR, Nerenberg CA, Jaffe RB (1992) Concordant suppression of serum immunoreactive luteinizing hormone (LH), follicle-stimulating hormone, alpha subunit, bioactive LH, and testosterone on postmenopausal women by a potent gonadotropin releasing hormone antagonist (Detirelex). *J Clin Endocrinol Metab* 74: 399–405
- Ataya KM, McKanna JA, Weintraub AM, Clark MR, LeMaire WJ (1985) A luteinizing hormone-releasing hormone agonist for the prevention of chemotherapy-induced ovarian follicular loss in rats. *Cancer Res* 45: 3651–3656
- Ataya KM, Tadros M, Ramahi A (1989) Gonadotropin-releasing hormone agonist inhibits physiologic ovarian follicular loss in rats. *Acta Endocrinol* 121: 55–60
- Banka CL, Erickson GF (1985) Gonadotropin-releasing hormone induces classical meiotic maturation in subpopulations of atretic preantral follicles. *Endocrinology* 117: 1500–1507
- Dohler KD, Wuttke W (1976) Circadian fluctuations of serum hormone levels in prepubertal male and female rats. *Acta Endocrinol* 83: 269–279
- Dvorshak-Harvey E, Peluso JJ, Bona-Gallo RV (1985) Effect of alterations in pulsatile luteinizing hormone release on ovarian follicular atresia and steroid secretion on diestrus 1 on the rat estrous cycle. *Biol Reprod* 33: 103–111
- Goldman BD, Grazia YR, Kamberi IA, Porter JC (1971) Serum gonadotropin concentrations in intact and castrated neonatal rats. *Endocrinol* 88: 771–776
- Jarrell J, Younglai EV, McMahon A, Barr R, O'Connell G, Belbeck L (1987) Effects of ionizing radiation and pretreatment with [D-Leu⁶,des-Gly¹⁰] luteinizing hormone-releasing hormone ethylamide on developing rat ovarian follicles. *Cancer Res* 47: 5005–5008
- Lintern-Moore S (1977) Effect of athymia on the initiation of follicular growth in the rat ovary. *Biol Reprod* 17: 155–161
- Lunenfeld B, Kraiem Z, Eshkol A (1975) The function of the growing follicle. *J Reprod Fertil* 45: 567–574
- Meijs-Roelofs HM, Kramer P, Van Cappellen WA, van Leeuwen EC (1990) Effects of a luteinizing hormone-releasing hormone antagonist in late-juvenile female rats: blockade of follicle growth and delay of first ovulation following suppression of gonadotropin concentrations. *Biol Reprod* 43: 607–613
- Ojeda SR, Ramirez VD (1972) Plasma level of LH and FSH in maturing rats: response to hemigonadectomy. *Endocrinol* 90: 466–472
- Ory SJ, Giebel SC, Leach RE, Ryan RJ (1990) The effects of LHRH agonist on follicular atresia in the neonatal and infant rat. Program of the 46th annual meeting of the Endocrine Society, Abstract P-076, Atlanta
- Peters H (1969) The development of the mouse ovary from birth to maturity. *Acta Endocrinol* 62: 98–116
- Sharpe KL, Bertero MC, Lyon BP, Muse KN, Vernon MW (1990) Follicular atresia and infertility in rats treated with a gonadotropin-releasing hormone antagonist. *Endocrinol* 127: 25–31
- Speroff L, Glass RH, Kase NG (1989) *Clinical Gynecologic Endocrinology and Infertility*. 4th ed. pp 121–128, Williams & Wilkins, Baltimore.
- Stachecki JJ, Yelian FD, Schultz JF, Leach RE, Armandt DR (1994) Blastocyst cavitation is accelerated by ethanol- or ionophore-induced elevation of intracellular calcium. *Biol Reprod* 50: 1–9
- Van den Dungen HM, van Dielen JA, Tilders FJ, van Rees GP, Shoemaker J (1989) Administration of a GnRH antagonist to immature rats affects subsequent female and male pubertal development differently. *Acta Endocrinol* 120: 778–784
- Wierman ME, Rivier JE, Wang C (1989) Gonadotropin-releasing hormone-dependent regulation of gonadotropin subunit messenger ribonucleic acid levels in the rat. *Endocrinology* 124: 272–278
- Wierman ME, Wang C (1990) Androgen selectivity stimulates follicle-stimulating hormone-beta mRNA levels after gonadotropin-releasing hormone antagonist administration. *Biol Reprod* 42: 563–571
- Zeleznik AJ, Kubik CJ (1986) Ovarian responses in macaques to pulsatile infusion of follicle-stimulating hormone (FSH) and luteinizing hormone: increased sensitivity of the maturing follicle to FSH. *Endocrinol* 19: 2025–2032

(Received October 1, 1998 / Accepted December 1, 1998)