



## **Leptogenys khammouanensis sp. nov. (Hymenoptera: Formicidae). A Possible Troglobitic Species of Laos, with a Discussion on Cave Ants**

Authors: Roncin, Eric, and Deharveng, Louis

Source: Zoological Science, 20(7) : 919-924

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.20.919>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# ***Leptogenys khammouanensis* sp. nov. (Hymenoptera: Formicidae). A Possible Troglotitic Species of Laos, with a Discussion on Cave Ants**

Eric Roncin<sup>1\*</sup> and Louis Deharveng<sup>2</sup>

<sup>1</sup>Laboratoire Dynamique de la Biodiversité, UMR 5172, Bât. 4R3, Université Paul Sabatier, 118 route de Narbonne, 31062 Toulouse Cedex 4, France

<sup>2</sup>Laboratoire d'Entomologie, ESA 8043 du CNRS, Muséum National d'Histoire Naturelle, 45 rue Buffon, 75005 Paris, France

**ABSTRACT**—The new species *Leptogenys khammouanensis* sp. nov. is described from two caves of the Khammouan karst (Laos). It is characterized by a set of striking morphological characters (reduced eyes, light pigmentation, slender body and very elongated legs and antennae), which recall the troglotitic traits of cave arthropods. Relations between caves and ants are discussed at this occasion, in the light of the recent biological explorations of caves in Southeast Asia. The classical view that ants are rare and unimportant in caves is challenged. Ants are actually major and regular components of guano assemblages in many caves of the region, but none of these guano species exhibit cave-related adaptation in its external morphology. Conversely, ants are very rare in low-resources habitats, where only accidental occurrence of outside species are reported in Southeast Asia. *Leptogenys khammouanensis* has been found only in such an oligotrophic environment, very deep in the cave and far from any guano deposits. Its presence there, together with its troglotitic traits, support the idea that *Leptogenys khammouanensis* might be the first truly troglotitic ant.

**Key words:** *Leptogenys*, taxonomy, troglotitism, cave ants, Laos

## **INTRODUCTION**

The pantropical genus *Leptogenys* presently includes 215 species of which 54 species and 28 subspecies are known from the Oriental region. African species have been revised by Bolton (1975) and Melanesian species by Wilson (1958). A revision of the Neotropical species by J. Lattke is on the way. No similar work exists for the Oriental region. Most species were described posteriorly to the only treatment of the Asian *Leptogenys* by Mayr (1879). The species of India have been keyed out by Forel (1900) and Bingham (1903), and Chinese fauna has recently been revised with several new species described (Xu 1996, 2000; Zhou 2001).

In this work, we describe *Leptogenys khammouanensis* sp. nov. collected in two caves of Laos. It exhibits the combination of morphological characters which defines troglotitism in arthropods (Christiansen, 1965): reduced eyes, elongate appendages (legs and antennae), light color. The possible troglotitic status of the new species is dis-

cussed in the context of our knowledge of Southeast Asia cave ants.

## **MEASUREMENTS AND INDICES**

All measurements were made on specimens in alcohol using a Leitz Weitzlar Stereomicroscope at magnifications of  $\times 150$ ,  $\times 100$ , and  $\times 50$  for total Length. Measurements (in millimeters) and abbreviations used here are based on Bolton (1975), Ward (1989) and Roncin (2002). Total Length (TL): Total length of an individual, from the mandibular apex to the gastral apex. Head Length (HL): Length of the head, excluding the mandibles. This measurement is taken in frontal view on the sagittal line from the foremost point of the clypeal margin (or its projection if situated laterally to the line) to the most posterior point on the occipital margin (or its projection). Head Width (HW): Maximum head width of the head measured in full-face view excluding the eyes. Scape Length (SL): Length of the first antennal segment, excluding the condylus. Cephalic index (CI):  $(HW / HL) \times 100$ . Scape Index (SI):  $(SL / HW) \times 100$ . Eye Diameter (EL): Maximal length of eye. Pronotal Width (PrW): Maximum width of pronotum in dorsal view. Mesosoma Length (ML): Diagonal length of the mesosoma, measured in lateral view from the point at which the pronotum meets the collare to the base of the propodeal lobes. Often called Weber's length of mesosoma. Petiole Height (PH): Height of the petiole measured in profile from the apex of the ventral (subpetiolar) process vertically to a line intersecting the dor-

\* Corresponding author: Tel. +33-5-61-55-64-35;  
FAX. +33-5-61-55-61-96.  
E-mail: roncin@cict.fr

salmost point of the node. Petiole length (PL): Length of the petiole from the anterior flanges to the posteriormost point of the petiole. Lateral Petiole Index (LPI):  $(PH / PL) \times 100$ . Dorsal Petiole Width (DPW): Maximum width of the petiole, measured in dorsal view. Dorsal Petiole Index (DPI):  $(DPW / PL) \times 100$ .

## DESCRIPTION

Genus *Leptogenys* Roger, 1861

*Leptogenys khammouanensis* sp. nov.

(Fig. 1: a–d)

**Description of Holotype (worker).** TL 7.6, HL 1.54, HW 1.01, CI 65, SL 1.99, SI 197, EL 0.09, PrW 0.71, ML 2.55, PH 0.46, PL 0.77, LPI 59, DPW 0.32, DPI 0.40

Head elongate, quite narrow, slightly broadened anteriorly, broadest across the eyes, the sides feebly convex. Occipital margin passing to lateral margin by a regular curve. A low nuchal carina present. Eyes small, oval, composed of 15–20 ommatidia. Maximum diameter of eye 0.093 mm, about  $0.09 \times HW$ , that is less than the maximum width of the scapes (0.127 mm). Eyes well in front of the middle of head sides, with their posterior borders situated on the same level than the posterior end of the frontal carinae in full face view. Eyes position frontal, not overlapping the outlines of head sides in full-face view. Circumocular groove strongly developed. Mandibles not striated, weakly shining, with feeble puncturations more numerous near masticatory border. Masticatory margin of mandibles with 14 teeth: the two apical ones well developed and followed by three denticles, one tooth and a series of 8 small teeth and denticles. Basal margin of mandibles close to the basal tooth of the apical margin with small crenulations. Clypeus triangular, projecting anteriorly as a distinct lobe closing tightly against the basal border of the mandibles, and with its foremost part broadly truncated. A wide longitudinal carina present on the median portion of clypeus. Anterior border of clypeus bidentate due to two massive enlarged central setae and laterally a row of normal and straight setae, all much shorter than other cephalic setae. Antennal scapes extremely long surpassing the occipital margin by almost half their length, with numerous erect to subdecumbent hairs, most of them being half the scape width. Funicular segments elongate: length of segments 1–3 ca 0.254, 0.422, 0.355 respectively (the partly visible condylar bulb of the first antennal segment is exclude from this measurement). Funicular segment 11: length 0.232; width 0.131.

Mesosoma elongate. Pronotum widest behind the middle, narrower in front than behind. Mesonotum markedly elongate (dorsal length of mesonotum 0.659 mm, taken at the level of the anterior border of metathoracic spiracles) and approximately of the same length than pronotum length (collare excluded) in dorsal view. Mesosoma shallowly depressed between mesonotum and propodeal dorsum but without any trace of the transverse metanotal suture. A transverse carina present on the posterior end of the declivitous face of propodeum surrounding insertion of petiolar peduncle. A low longitudinal median carina is present on the

mesosternum and is crossed by small wrinkles. Legs extremely long, with metafemur length (2.10 mm) exceeding gaster length.

Node of the petiole in dorsal view much longer than broad. Anterior part of the node with a carina separating it from the true anterior peduncle which is indeed very short. Dorsal outline of the anterior part of the node concave in lateral view. The node is widening progressively backward, giving to the anterior part of the node an aspect of false peduncle.

Gaster constricted between its first and second segment. Sting relatively short.

All body sculpture smooth. Body color light orange-yellow with small brown spots regularly spaced at the basis of body setae. Callow individuals yellow. Body with a dense pilosity especially on head, gaster and coxae, and with the longest hairs situated on clypeus, ventral surface of the head capsule and procoxae.

Gyne and male unknown.

**Etymology.** From “Khammouan”, a province of Laos with beautiful calcareous landscapes, uncountable caves and subterranean rivers.

**Type material.** Holotype worker, Laos: Tham (= cave) Nam Non (18.0270° N, 104.6883° E, coordinates from Brouquisse, 1999), alt. 185 m, several km inside the cave, hand collecting, 15.II.1999, (LAO-070), J. Lordon and J.F. Vacqu  .

Paratypes workers: 2 workers, same data; collected with holotype.


Other material: 4 workers, Laos: Ban Nakok: Tham Th   cave (17.9964° N, 104.4482° E, coordinates from Brouquisse, 1999), 11.II.1998, (LAO-016), collected at about 500 m from cave mouth, hand collecting, L. Deharveng and A. Bedos.

Standard measurements: workers. TL 7.0–7.8, HL 1.54–1.65, HW 0.98–1.05 CI 64–65, SL 1.94–2.12, SI 197–201, EL 0.09, PrW 0.66–0.74, ML 2.42–2.64, PH 0.45–0.50, PL 0.76–0.84, LPI 58–59, DPW 0.29–0.32, DPI 0.38–0.40

One worker from the type locality possesses on the right side an abnormal eye of circular shape reduced to a single large ommatidia of 0.041 mm diameter.

The holotype of the new species is deposited in the Museum National d'Histoire Naturelle de Paris (MNHN) and the paratypes in the author's collection.

**Remarks.** The general habitus of *Leptogenys khammouanensis* with very elongate head, mesosoma, petiole, antennae and legs is very similar to that of *Leptogenys assamensis* Forel, 1900, a species that seems to be known only from the type series collected by Long in the Gar   hills in Assam (E. India), and to *Leptogenys ergatogyna* Wheeler, 1922, a forest species known from Zaire, Uganda and Cameroun and which presents few affinities to other African species (Bolton, 1975). However, *Leptogenys assamensis* and *Lep-*



**Fig. 1.** *Leptogenys khammouanensis* sp. nov. a–d, holotype worker. a, lateral view; b, dorsal view; c, head in frontal view; d, anterior part of head with clypeal setae figured.

*togenys ergatogyna* differ from *L. khammouanensis* by the presence in both species of the metanotal suture, the absence of a pair of enlarged, straight and hard setae on the clypeal lobe border, less mandibular teeth (no mandibular teeth in *L. assamensis* and only two apical teeth in *L. ergatogyna*) and by other characters (color, eyes size) discussed in the paragraph below concerning troglbiomorphic characters in *Leptogenys*.

## DISCUSSION

The combination of reduced eyes, very pale color and elongate appendages isolates *Leptogenys khammouanensis* in its genus. Short of a revision of Oriental *Leptogenys*, it is impossible to assess its affinities, especially because these conspicuous morphological traits are adaptive to hypogean life. For this reason, however, they are of great interest, as they point to an unusual mode of life for an ant. The possibility of troglbionic life for ants has long puzzled specialists (see Wilson, 1962), and is still a matter of debate (Tinaut and López, 2001). Compared to other ant candidates to troglbionic life, *L. khammouanensis* is one step ahead in terms of morphological characters linked to cave life. This question of troglbionic and troglbiomorphic ants will be discussed below, taking into account recent literature and the rich material of cave ants we have gathered in Southeast Asia.

Assessing the troglbionic status of a taxon is not easy. However, troglomorphy (Christiansen, 1962), or better troglbiomorphy (Boutin, in press), i.e. the set of morphological traits characteristics of cave organisms, is always a strong indication of troglbionism (while the opposite is not always true). Troglbiomorphy in arthropods is defined by four morphological traits: loss of wings in winged arthropod groups, reduction of eyes, reduction of tegumentary pigment and elongation of appendages (Vandel, 1964; Christiansen, 1965; Culver, 1982; Marques and Gnaspini, 2001). All known specimens of *L. khammouanensis* are workers, which are always apterous in ants. We do not know therefore the state of the first character in the new species, but the last three traits are present.

### Troglbiomorphic characters in *Leptogenys*

The small eyes and light color of *L. khammouanensis* recall those of the group of *L. processionalis* sensu Taylor (1969) (= *fallax*-group of Andersen, 2000), which comprises three Australian species, *L. fallax* (Mayr, 1876), *L. tricolor* Taylor, 1969 and *L. fortior* Forel 1900, and the Oriental species *L. myops* (Emery, 1887), *L. crassicornis* Emery, 1895 and *L. processionalis* (Jerdon, 1851). Species of this group possess also light color and reduced eyes varying from a single (*L. tricolor*) to about fifteen facets, but antennae are short, and size is small. Such a morphology appears as a classical adaptation to endogean, not to cave life. At least several African species in the *guineensis*- and the *nitida*-group also possess small eyes, and are yellow brown to

dark brown. However their head and appendages are so not elongated.

In its slender habitus (very elongate head, mesosoma, petiole, antennae and legs), *Leptogenys khammouanensis* is similar to *Leptogenys ergatogyna* Arnold 1954 from Africa (redescribed in Bolton, 1975) and to *L. assamensis* Forel, 1900 from Assam (redescribed by Bingham, 1903). But both species differ from *L. khammouanensis* by their black color and larger eyes, and cannot be considered as troglbiomorphic.

No species of *Leptogenys* therefore approaches *L. khammouanensis* in its combination of troglbiomorphic characters.

### Troglbiomorphic ants

Only one species of Formicidae, *Aphaenogaster cardenai* Espadaler, 1981, could be associated to subterranean habitats (Decu *et al.*, 1998, Tinaut and López, 2001). This rare Spanish species has always been collected in cryptic habitats: under big rocks, galleries of rodents and caves. However in this latter habitat no nests have ever been found, which led these authors to postulate that *A. cardenai* is more probably an inhabitant of the MSS ("Milieu Souterrain Superficiel" of Juberthie *et al.*, 1980, "superficial underground compartment" in Humphreys, 2000), than a strictly cave dwelling species.

*A. cardenai* is related to *A. splendida* (Roger, 1859), *A. ovaticeps* (Emery, 1898) and *A. muelleriana* Wolf, 1915. All have reduced eyes, slender body, elongate appendages and often pale color, compared to other species of the genus. Bernard (1968: 136) stressed the peculiar morphology of the species then known and noted that they have "un faciès aphaenopsien de cavernicole" ("an *Aphaenops*-like morphology of cave species"). He supposed they inhabit hypogean habitats, probably deep cracks that they rarely leave, hence their rarity in collection. However a few captures from surface habitats (Forel, 1911; Wheeler and Mann, 1916) could indicate nocturnal activity more than hypogean life.

*Leptogenys khammouanensis* is therefore the second example in Formicidae with the *Aphaenogaster* of the *splendida*-group, and the first Ponerinae, which combines hypogean life with microphthalmia, light color, and elongated appendages, i.e. the typical adaptive characters of cave inhabiting arthropods. None of the *splendida*-group species of *Aphaenogaster* is cave-restricted. Conversely, *L. khammouanensis* was observed only in deep parts of caves and in rather significant numbers, bringing an exciting question to the fore: is *L. khammouanensis* the first truly troglbionic ant?

### Cave ants

Though frequently cited from caves, ants have not provided so far any unambiguously troglbionic species. Most records (Wilson, 1962; Tinaut and López, 2001) concern in fact accidental occurrences, generally not far from cave

entrance (Decu *et al.*, 1998). Most species supposed to be strictly cavernicolous have been later found also outside caves. Even the rare *Hypoponera ragusai* considered by Tinaut (2001) to be limited to caves in Europe has been collected outside caves in France (Bernard, 1968) and in the Mediterranean islands of Lampedusa and Linosa (Mei, 1992, 1995).

This rarity of ants in caves concerns essentially temperate regions. Though limited, available evidence suggests that Formicidae might be much more frequent in tropical caves. More than sixty species have already been collected from the dark part of various caves of Southeast Asia (Roncin *et al.*, 2001, and unpublished data). Most of these species, well-known outside, are represented by isolated specimens in cave collections. But several are regular guano inhabitants, like *Hypoponera confinis* in the Farm caves of Myanmar (Annandale *et al.*, 1913) and an unidentified species in the Mulu caves of Sarawak (Chapman, 1982). In fact, ants were present in most guano caves recently sampled in Southeast Asia, with *Hypoponera* the dominant genus (Roncin *et al.*, 2001). They were sometimes found very far from cave entrance, living in loose colonies, foraging in or around the guano piles, where they are believed to prey on a rich fauna of micro- and meso-arthropods. None of these regular guano species are troglobiomorphic, and all were also collected outside caves. Among the hundreds of caves prospected so far in all regions of Southeast Asia (see Juberthie and Decu, 2001 for an overview), troglobiomorphic species were found only in Laos with *L. khammouanensis*.

*Leptogenys khammouanensis* was encountered far from the entrance in two big caves, Tham Nam Non (22 km long, the longest cave of continental Southeast Asia, Mouret, 2001) and Tham Thê (2.2 km long, Brouquisse, 1999). In spite of being 25 km apart and in different hydrogeological systems, both caves do belong to the same, uninterrupted, huge limestone unit. In Tham Nam Non, *L. khammouanensis* was collected at about 4.5 km of the entrance. These giant caves of the Khammouan karst host a rich troglotic fauna, only recently discovered: microphthalmic crabs (*Erebusa calobates*, Yeo and Ng), various blind terrestrial Isopods, Araneids and Millipeds (Polydesmidae, Glomeridae), Campodeids, springtails, blind Nocticolid cockroaches, blind or microphthalmic *Diestrammena* sp. crickets (Besson *et al.*, 2001). As numerous species of *Leptogenys* are woodlice-hunters (for a full list of references see: Hölldobler and Wilson, 1990; Dejean 1997), the terrestrial isopods frequent in these caves could constitute a potential diet of *L. khammouanensis*, but this has to be confirmed.

The genus *Leptogenys* has already been found in caves. *Leptogenys jeanneli* Santschi, 1914 was described from a cave in Tanzania, and *Leptogenys diminuta* (Smith, 1857) was collected in the Batu Caves of peninsular Malaysia (Wilson, 1962; McClure, 1965). However, none of these species is cave-restricted, and none exhibits the combina-

tion of troglbiomorphic traits of *L. khammouanensis*.

Having analysed what was known about cave ants, Wilson (1962) hypothesized that social insects "never become truly troglbionic" because "they are unable to maintain sufficiently large cave demes" (implicitly because of food scarcity). The well-known link between troglbiomorphy and oligotrophic habitats (Deharveng and Bedos, 2000) certainly explains the extreme difficulty for ants to establish long-term colonies and to adapt to cave environment. This view is challenged today by the discovery of *L. khammouanensis* with its clear troglbiomorphic morphology. The unusually large underground voids of Laos may have given the opportunity for such an evolution to take place, by providing large food reservoirs on the long-term. To confirm this exciting hypothesis, it remains to document the peculiarities of the biology and social life of the new species (Tinaut and López, 2001).

## ACKNOWLEDGEMENTS

We thank the members of the 1998 and 1999 French caving expedition to Laos, who provided the specimens, especially Anne Bedos, Jérôme Lordon, and Jean-François Vacquie. Bernard Kaufmann and Hervé Jourdan made useful comments on the manuscript. We further thank Penelope Greenslade and Robert Taylor for style improvement of large parts of the manuscript.

## REFERENCES

- Andersen AN (2000) The ants of northern Australia. A guide to the monsoonal fauna. CSIRO Publishing, Collingwood
- Annandale N, Coggin Brown J, Gravely FH (1913) The limestone caves of Burma and the Malay Peninsula. J Proc Asiat Soc Bengal 9: 391–424
- Arnold G (1954) New Formicidae from Kenya and Uganda. Ann Mus R Congo Belg Zool 1: 291–295
- Bernard F (1968) Les fourmis (Hymenoptera : Formicidae) d'Europe occidentale et septentrionale. Masson, Paris
- Besson JP, Deharveng L, Brehier F (2001) Laos. In "Encyclopaedia Biospeologica Tome 3" Ed by C Juberthie, V Decu, Soc Biospéologie, Moulis, pp 1883–1889
- Bingham CT (1903) The fauna of British India, including Ceylon and Burma. Hymenoptera. 2. Ants and Cuckoo-wasps. Taylor and Francis, London
- Bolton B (1975) A revision of the ant genus *Leptogenys* Roger (Hymenoptera: Formicidae) in the Ethiopian Region with a review of the Malagasy species. Bull Br Mus Nat Hist (Entomol) 31: 237–305
- Boutin C Organisms classification. In "Encyclopedia of cave and karst science", Ed by J Gunn, Fitzroy Dearborn Publishers, London (in press)
- Brouquisse F (1999) Catalogue of caves in the Lao peoples's Democratic Republic. Int Caver 25: 13–16
- Chapman P (1982) The ecology of caves in the Gunung Mulu National Park, Sarawak. Trans Br Cave Res Assoc 9: 142–162
- Christiansen KA (1962) Proposition pour la classification des animaux cavernicoles. Spelunca 2: 76–78
- Christiansen KA (1965) Behavior and form in the evolution of cave Collembola. Evolution 19: 529–537
- Culver D (1982) Cave life, evolution and ecology. Harvard Univ Press, Cambridge
- Decu V, Casale A, Scaramozzino PL, López F, Tinaut A (1998)

- Hymenoptera. In "Encyclopaedia Biospeologica Tome 2" Ed by C Juberthie, V Decu, Soc Biospéologie, Moulis and Bucarest, pp 1015–1024
- Deharveng L, Bedos A (2000) The cave fauna of southeast Asia. Origin, evolution and ecology. In "Ecosystems of the world Vol 30 Subterranean ecosystems" Ed by H Wilkens, DC Culver, WF Humphreys, Elsevier, Amsterdam, pp 603–632
- Dejean A (1997) Predatory behavior in the genus *Leptogenys*: a comparative story. *J Insect Behav* 10: 177–191
- Forel A (1900) Les formicides de l'Empire des Indes et de Ceylan. Part. 7. *J Bombay Nat Hist Soc* 13: 303–332
- Forel A (1911) Fourmis nouvelles ou intéressantes. *Bull Soc Vaud Sci Nat* 47: 331–400
- Hölldobler B, Wilson EO (1990) The ants. Springer-Verlag, Berlin
- Humphreys WF (2000) Background and glossary. In "Ecosystems of the world Vol 30. Subterranean ecosystems" Ed by H Wilkens, DC Culver, WF Humphreys, Elsevier, Amsterdam pp 3–14
- Juberthie C, Decu V. Ed (2001) *Encyclopaedia Biospeologica Tome 3*, Soc Biospéologie, Moulis, pp 1375–2294
- Juberthie C, Delay B, Bouillon M (1980) Extension du milieu souterrain en zone non calcaire : description d'un nouveau milieu et de son peuplement par les Coléoptères troglobies. *Mém Biospéol* 7: 19–52
- Marques AC, Gnaspini P (2001) The problem of characters susceptible to parallel evolution in phylogenetic reconstructions: suggestion of a practical method and its application to cave animals. *Cladistics* 17: 371–381
- Mayr G (1879) Beiträge zur Ameisen-Fauna Asiens. *Verh K K Zool Bot Ges Wien* 28: 645–686
- McClure HE (1965) Microcosms of Batu caves. *Malay Nat J* 19: 65–74
- Mouret C (2001) Le karst du Khammouane au Laos central. *Spe-lunca* 84: 7–32
- Mei M (1992) Su alcune species endogee o cryptobiotiche della myrmecofauna italiana (Hymenoptera, Formicidae). *Fragm Entomol* 23: 411–422
- Mei M (1995) Arthropoda di Lampedusa, Linosa e Pantelleria (Canale di Sicilia, Mar Mediterraneo). Hymenoptera Formicidae (con diagnosi di due nuove specie). *Nat Sicil* 19 (Suppl): 753–772
- Roncin E (2002) Two new *Tetramorium* species (Hymenoptera: Formicidae) from Vietnam with a discussion of the *mixtum*, *tonganum* and *scabrosum* groups. *Sociobiology* 40: 281–292
- Roncin E, Deharveng L, Bedos A (2001) Cave ants in Southeast Asia. In "XV International Symposium of Biospeleology", Intervales, São Paulo (Brasil), 8–15 July 2001. Abstracts: 66–67
- Taylor R (1969) The identity of *Dorylozelus mjobergi* Forel (Hymenoptera: Formicidae) *J Aust Entomol Soc* 8: 131–133
- Tinaut A (2001) *Hypoponera ragusai* (Emery, 1895) a cavernicolous ant new for the Iberian peninsula (Hymenoptera, Formicidae). *Graellsia* 57: 3–8
- Tinaut A, López F (2001) Ants and caves: sociability and ecological constraints (Hymenoptera, Formicidae). *Sociobiology* 37: 651–659
- Vandel A (1964) Biospéologie. La biologie des animaux cavernicoles. Gauthier Villars, Paris
- Ward PS (1989) Systematic studies on pseudomyrmecine ants: revision of the *Pseudomyrmex oculatus* and *P. subtilissimus* species groups, with taxonomic comments on other species. *Quaest Entomol* 25: 393–468
- Wheeler WM, Mann WM (1916) The ants of the Phillips expedition to Palestine during 1914. *Bull Mus Comp Zool* 59: 167–174
- Wilson EO (1958) Studies on the ant fauna of Melanesia. Studies on the ant fauna of Melanesia. 1. The tribe Leptogenyini. 2. The tribes Amblyoponini and Platythyreini. *Bull Mus Comp Zool* 118: 101–153
- Wilson EO (1962) The trinidad cave ant *Erebomyrma* (= *Spelaemyrmex*) *urichi* (Wheeler), with a comment on cavernicolous ants in general. *Psyche* 69: 62–72
- Xu Z (1996) A taxonomic study on the ant genus *Leptogenys* (Hymenoptera: Formicidae) in China [in Chinese]. *J Yunnan Agric Univ* 11: 222–227
- Xu Z (2000) Five new species and one new record species of the ant genus *Leptogenys* Roger (Hymenoptera: Formicidae) from Yunnan Province, China. *Entomol Sinica* 7: 117–126
- Zhou S (2001) Ants of Guangxi [in Chinese]. Guangxi Normal University Press, Guilin

(Received July 12, 2002 / Accepted March 12, 2003)