



Spermiophagy in the Male Reproductive Tract of Some Passerine Birds

Authors: Chiba, Akira, Nakamura, Masahiko, and Morimoto, Gen

Source: Zoological Science, 28(9) : 689-693

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.28.689>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Spermiophagy in the Male Reproductive Tract of Some Passerine Birds

Akira Chiba^{1*}, Masahiko Nakamura² and Gen Morimoto³

¹Department of Biology, Nippon Dental University School of Life Dentistry at Niigata, Niigata 951-8580, Japan

²Laboratory of Animal Ecology, Joetsu University of Education, 1 Yamayashiki-machi, Joetsu-shi, Niigata 943-8512, Japan

³Department of Animal Behavior, Graduate School of Science, Rikkyo University, 3-34-1, Nishi-Ikebukuro, Toshima-ku, Tokyo 171-8501, Japan / Tokyo Bay Ecosystem Research Center, Toho University, 2-2-1, Miyama, Funabashi-shi, Chiba 274-8510, Japan

In order to elucidate the locus and means of spermiophagy in passerine birds, we examined histologically the entire male reproductive tract of sexually mature birds of three passerine species with different forms of sperm competition, namely, the alpine accentor (*Prunella collaris*), the red-flanked bush robin (*Tarsiger cyanurus*), and the Bengalese finch (*Lonchura striata* var. *domestica*). Spermiophagy occurred consistently and frequently in the epithelial layer of the seminal glomera and ejaculatory duct in each species, which were regularly identified by non-ciliated epithelial cells. The epithelial spermiophagy was occasional or infrequent in other portions of the seminal tract, and spermiophagy by macrophages was uncommon throughout the tract. Quantitative data in the seminal glomera and ejaculatory duct gave no clear answer concerning a possible relationship between the epithelial spermiophagy and different levels of sperm competition among these passerine species. In conclusion, the epithelial lining of the terminal region of the seminal tract is the main site for spermiophagy in the male reproductive tract of these passerine species, which activity serves to maintain the quality of semen by eliminating infertile spermatozoa as well as sperm remaining at the end of the breeding season.

Key words: epithelial cells, passerine birds, seminal glomera, spermiophagy, sperm competition

INTRODUCTION

Spermiophagy (or spermatophagy) denotes the phagocytosis of spermatozoa and their fragments by somatic cells (Holstein, 1978). This activity has been reported to occur widely in various groups of vertebrates, e.g., teleosts (Porawski et al., 2004), amphibians (Sever, 1992), reptiles (Akbarsha et al., 2007), birds (Tingari and Lake, 1972; Nakai et al., 1989; Aire, 2000), and mammals (Murakami et al., 1979; 1985; Goyal, 1982; Abou-Elmagd and Wrobel, 1990). In mammals, spermiophagy takes place in various portions of the male reproductive tract, such as the seminiferous tubules (Holstein, 1978), rete testis, efferent duct (Holstein, 1978; Goyal, 1982), ampulla of the vas deferens (Murakami et al., 1985), seminal vesicle (Murakami et al., 1978), and ejaculatory duct (Abou-Elmagd and Wrobel, 1990). In these locations, Sertoli cells, macrophages or epithelial cells lining the seminal tract are involved in this spermiophagy. According to Holstein (1978), spermiophagy is frequently seen in the gonads of older men and in certain cases of oligozo-

spermia, i.e., generally defined, in man, as a concentration of spermatozoa less than 20 million per ml. Also, in the macaque *Macaca fascicularis*, spermiophagy increases at the end of the spermatogenetic season.

In birds, previous studies on the domestic fowl *Gallus domesticus* have shown that the epithelial lining of the rete testis and epididymal efferent ducts phagocytose spermatozoa only occasionally (Tingari, 1972; Nakai et al., 1989; Kirby et al., 1990) and that macrophages are also involved in spermiophagy (Nakai et al., 1989). Later, Aire (2000) reported active spermiophagy by the non-ciliated cells in the epithelial lining of the epididymal efferent ducts of the normal chicken. In contrast, studies on wild passerines (Middleton, 1972; Chiba and Nakamura, 2003) suggested spermiophagy in the seminal glomera (or glomus). This organ is known as a characteristic feature of the male reproductive system of passerines (although it has also been reported to exist in the psittacid *Melopsittacus undulatus*; Samour et al., 1987), and is regarded as an anatomical adaptation that ensures a storage and probable maturation site for spermatozoa that are necessary for sperm competition (Bedford, 1979; Lake, 1981; Birkhead et al., 1993; Aire, 2007). In light of the above, the following subjects should be investigated and clarified in particular concerning the passerines: (1) What is the main site for spermiophagy in their male repro-

* Corresponding author. Phone: +81-25-267-1500;
Fax : +81-25-267-1134;
E-mail: chiba@ngt.ndu.ac.jp

doi:10.2108/zsj.28.689

ductive system? (2) What types of cell are involved in spermiphagy in the seminal tract? (epithelial cells, macrophages, or cells of other types?), and (3) What is the biological significance of spermiphagy in the normal healthy animal? At present, nearly nothing is known about these aspects of spermiphagy in wild birds (Aire, 2007).

In order to answer these questions, we microscopically examined spermiphagy in the male reproductive system of three species of passerines, namely, the alpine accentor (*Prunella collaris*), the red-flanked bush robin (*Tarsiger cyanurus*), and the Bengalese finch (*Lonchura striata* var. *domestica*). We also focused on its possible relation to sperm competition (Birkhead et al., 1993), because there is a background information regarding this possibility; i.e., in passerines with low levels of sperm competition, the incidence of malformed sperm is higher than in species with intense sperm competition, strongly suggesting that spermiphagy may be more intense in species in which sperm competition seems to be less intense (Calhim et al., 2007; Birkhead et al., 2007; Immler et al., 2008). We consider this prediction worth testing in the present study.

MATERIALS AND METHODS

Animals

All treatments complied with the *Guidelines for the Care and Use of Laboratory Animals in Nippon Dental University*.

Mature adult males of the alpine accentor (four birds, 44–55 g in body weight (b.w.)), red-flanked bush robin (five birds, 13–14 g in b.w.), and Bengalese finch (seven birds, 14–16 g in b.w.) were used in this study. The samples of alpine accentor and red-flanked bush robin were collected (by permission of the Ministry of the Environment of Japan and the Government of Shizuoka Prefecture) during their breeding season, from June to August, on the summit of Mt. Norikura, Nagano Pref., Japan (36°06'N, 137°33'E, 2,600–3,026 m above sea level (a.s.l.)) and in a forest of Mt. Fuji, Shizuoka Pref., Japan (35°26'N, 138°47'E, 1,500–2,000 m a.s.l.), respectively. Those of the Bengalese finch were purchased from a local dealer, but information about their mating conditions in cages was incomplete.

Light and electron microscopy

After measurement, the birds were killed by quick dislocation of their cervicales, dissected, and immersed in a solution of 5% buffered formaldehyde, 5% paraformaldehyde in 0.1 M phosphate buffer at pH 7.2 or in Bouin's mixture for a few days. The entire male reproductive system was carefully dissected out and re-fixed in freshly prepared 5% buffered formaldehyde or Bouin's fixative for light microscopy (LM) or in Karnovsky's fluid for transmission electron microscopy (TEM). The sexual organ index (SOI: the weight of the entire male reproductive system times 100 divided by the body weight) was calculated thereafter. Various sexual organs were examined under a dissection microscope, appropriately trimmed, and processed for routine histology and cytology. Paraffin sections cut at 8- μ m thickness were stained with the following stains: Mayer's hematoxylin-eosin, azan trichrome, and periodic acid Schiff's (PAS) reagent for demonstration of polysaccharide. The sections were examined under a light microscope. For TEM, the aldehyde-fixed tissue blocks were rinsed with 0.1 M phosphate buffer (pH 7.4), immersed in 1% osmium tetroxide in the same buffer for one hour, dehydrated, and embedded in Spurr resin. Ultra-thin sections were cut with a diamond knife, stained with uranyl acetate and lead citrate, and viewed with a JEOL 1200 EX electron microscope. Semi-thin sections were stained with toluidine blue in borax and observed under a light microscope.

Quantitative assessment

Spermiphagy was microscopically examined and its incidence was roughly scored as follows (Table 1): absent (–), rare or uncommon (+), occasional (++) or frequent (+++). In order to compare the incidence of epithelial spermiphagy (i.e., spermiphagy by seminal duct epithelium) among these three species, we counted the number of absorbed spermatozoa in the epithelium of the posterior region of the seminal tract, namely the seminal glomera (SG) and the ejaculatory duct (ED). For this purpose, three alpine accentors, five red-flanked bushrobins, and three Bengalese finches were examined and the number of absorbed spermatozoa per single section of the seminal tract was counted in 40 randomly selected sections per individual. Furthermore, in order to assess local differences of the epithelial spermiphagy we similarly counted the number of absorbed spermatozoa in three different portions of the seminal tract, i.e., the proximal, middle, and distal portions, respectively (Figs. 1 and 2): the proximal portion includes approximately anterior 2/5 of the SG, the middle one occupies the following (middle) 2/5 of the SG, and the distal one corresponds to the posterior 1/5 of the SG and the whole ejaculatory duct (Fig. 2 for example). In the alpine accentor, remarkably developed SG and ED (Fig. 1) are actually equivalent to the dorsal and ventral lobes of the cloacal protuberance described in a previous paper (Chiba and Nakamura, 2003). In practice, the number of absorbed spermatozoa per single section of the seminal tract was counted in randomly selected 40 sections of each portion per individual. The counting was carried out on one alpine accentor (one of four birds, tentatively named Ac-1), three red-flanked bushrobins (three of five birds: Br-1, -2 and -3), and three Bengalese finches (three of seven birds: Bg-1, -2 and -3) (Fig. 4), due to the limited number and condition of the specimens. We used the median of each individual as a representative value to minimize bias due to pseudoreplication.

RESULTS AND DISCUSSION

The male reproductive system of the birds examined showed considerable inter-specific variation in their SOI values (Table 1), mainly due to differences in the weight of the testis and SG/ED relative to that of the body mass (Fig. 1A–C), probably reflecting the degree of sperm competition related to their mating systems. For example, the polygynandrous alpine accentor, having a higher frequency of copulation than the other two species, has a remarkably voluminous testis and SG/ED (Nakamura, 1990; Chiba and Nakamura, 2003) in comparison with those of the monoga-

Table 1. Summary of spermiphagy in the male reproductive tract of three species of passerines, namely, the alpine Accentor, red-flanked bush robin, and Bengalese finch.

| Name of animal | alpine accentor | | red-flanked bush robin | | bengalese | |
|------------------------------|-----------------|----|------------------------|----|-----------|----|
| Number of animals | n = 4 | | n = 5 | | n = 7 | |
| Mean SOI (%)* | 6.8 | | 2.8 | | 0.6 | |
| Range | 4.6–8.4 | | 1.9–3.6 | | 0.5–0.7 | |
| Type of spermiphagy | SE | SM | SE | SM | SE | SM |
| Portions of seminal tract | | | | | | |
| Rete testis | – | – | – | – | – | – |
| Epididymis | ++ | + | + | – | + | – |
| Vas deferens | + | – | ++ | – | – | – |
| Seminal glomera | +++ | + | +++ | + | +++ | + |
| Ejaculatory duct and papilla | +++ | – | +++ | – | +++ | – |

*SOI, sexual organ index (weight of the entire male reproductive system relative to the body weight, %). SE, spermiphagy by epithelial cells; SM, spermiphagy by possible macrophages; –, absent; +, rare or uncommon; ++, occasional; +++, frequent.




Fig. 1. Macroscopic features of the fully developed male reproductive system (one side) of the alpine accentor (**A**), red-flanked bush robin (**B**), and Bengalese finch (**C**). Each system was isolated from visceral organs and laid cranially to the left. e, epididymis; ED, ejaculatory duct; SG, seminal glomera; t, testis; vd, vas deferens; thick arrows, terminal papilla of the ejaculatory duct. Proximal (p), middle (m), and distal (d) portions of the SG/ED are also indicated. Scale bar = 1 cm.

mous red-flanked bushrobin, which has occasional extra-pair copulation (about 50% of nests showed the extra-pair fertilization; unpublished data by Morimoto) and of the domesticated monogamous Bengalese finch, in which detailed results on the size of their testes and SG, ejaculating capacity, and sperm depletion were previously documented (Birkhead, 1991).

LM examination for spermiophagy was carried out throughout this system, i.e., the testis, epididymis, vas deferens, SG, ED, and terminal papilla. Spermiophagy was found consistently and frequently in the epithelial layer of the SG and ED in each species examined (Fig. 3A–C; Table 1). There, morphologically intact spermatozoa in the cavity had been ingested by non-ciliated epithelial cells (Fig. 3A–C). However, no clear numerical difference was found in the incidence of the epithelial spermiophagy among the 3 species examined: alpine accentor (20.7 ± 4.6 as mean \pm SD, $n = 3$); red-flanked bushrobin (19.1 ± 10.7 , $n = 5$); and Bengalese finch (21.8 ± 27.9 , $n = 3$). On the other hand, the epithelial spermiophagy in the SG/ED tended to be more active towards the posterior portion of the seminal duct system in the alpine accentor and the red-flanked bushrobin (Fig. 4), although this tendency was not necessarily clear in the Bengalese finch, in which the incidence of epithelial spermiophagy was considerably variable (Fig. 4). The numerical data in Fig. 4 were also shown: 14.6 in the proximal portion, 17.0 in the middle portion, and 25.9 in the distal por-




Fig. 2. Macroscopic view of the posterior region of the male reproductive system of the red-flanked bush robin to show the anatomical arrangement of the seminal glomera (SG), ejaculatory duct (ED), and terminal papilla (tp) to the cloacal structures (c). The same sample is carefully dissected and unfolded to be subdivided into 3 portions, namely proximal (p), middle (m), and distal (d) portions. Scale bar = 1 cm

tion for the alpine accentor ($n = 1$); 5.3 ± 23.0 (average \pm SD), 8.2 ± 12.4 , 18.1 ± 12.8 for the red-flanked bushrobin ($n = 3$); 18.8 ± 25.2 , 25.1 ± 32.2 , 16.7 ± 15.7 for the Bengalese finch ($n = 3$), respectively.

TEM examination confirmed the spermiophagy and demonstrated that the epithelial cells had a varied number of microvilli at their apical plasma membrane, empty vesicles or vacuoles of different sizes in their apical cytoplasm, lysosome-like dense bodies, and a moderate number of mitochondria (Fig. 3E), in general accordance with the previous observations made in domestic fowl (Tingari, 1972; Nakai et al., 1989; Aire, 2000). Epithelial spermiophagy was occasional or rare in the epididymal efferent ducts and vas deferens, but not found in the rete testis or ejaculatory duct or its terminal papilla (Table 1). We also observed that free cells, presumptive macrophages, had ingested spermatozoa and fragments of spermatogenetic cells in the cavities of the seminal tract (Fig. 3D, F; Table 1), although this type of spermiophagy was uncommon throughout the seminal tract.

The present study along with previous observations (Middleton, 1972; Chiba and Nakamura, 2003) demonstrated spermiophagy in the seminal glomera and some other locations of the seminal tract in the passerines. Such findings support the view that the seminal glomera, a site for storage and probable maturation of spermatozoa in passerines (Lake, 1981; Bedford, 1979), simultaneously serves as a major site for spermiophagy, contributing to maintenance of the quality of semen by removal of remaining and infertile spermatozoa, i.e., acting as a “physiological filter” (Chiba and Nakamura, 2003) or “sorting apparatus” for selection of healthy fertile sperm as well as for removing immature or aging sperm (Birkhead et al., 1995). Our findings also indicate that this epithelial spermiophagy is the main process for




Fig. 3. Spermiophagy by non-ciliated epithelial cells (ep) in the seminal glomera (A–C and E) and by macrophages (mc) in the seminal cavities (D) and (F). A–D, light microscopic photographs; E and F, electron microscopic photographs. A, alpine Accentor; B, red-flanked bush robin; C, Bengalese finch. Arrows in “A”–“E” point to ingested spermatozoa; ly, lysosome-like dense body; pl, phagolysosome; v, vacuole. Scales in “A”–“D” = 10 μ m; and in “E” and “F” = 1 μ m.




Fig. 4. Local difference of the epithelial spermiophagy in the proximal, middle, and distal portions of the seminal glomera and ejaculate duct of the 3 passerine birds, alpine accentor (Ac-1), red-flanked bush robin (Br-2, -3 and -4), and Bengalese finch (Bg-5, -6 and -7).

sperm absorption in the seminal tract of the passerines under normal healthy conditions. Concerning the site and cell type for spermiophagy in the seminal tract, the present

results are inconsistent with those found for chickens, in which spermiophagy by macrophages and/or epithelial cells occurs mainly in the epididymis (Tingari, 1972; Nakai et al., 1989; Kirby et al., 1990; Aire, 2000). In more detail concerning the spermiophagy of the chicken, different authors have reported different results. Tingari (1972) reported that three types of the epithelial cells are responsible for spermiophagy, i.e., the low cuboidal cell in the rete testis, the ciliated cell lining the efferent and connecting ducts in the epididymis, and the non-ciliated type II cell in the vas deferens, whereas Nakai et al. (1989) showed that the luminal macrophages, not epithelial cells, in the epididymis and rete testis showed active spermiophagy. On the other hand, Aire (2000) reported active spermiophagy by the non-ciliated type I epithelial cell in the efferent duct of the epididymis. Thus, the epithelial spermiophagy in the chicken appeared to occur in a wide range of the seminal tract and to be performed by various cell types, being in contrast to the case of the passerine birds studied. We consider that the non-ciliated cells in the passerines responsible for spermiophagy may be equivalent to the non-ciliated type II cell

in the chicken. This discrepancy may be due to the differences in the systematic position and breeding habits between the passerines and domestic chickens. The alpine accentor and red-flanked bushrobin are seasonal breeders and show clear seasonal changes in their reproductive activities (Nakamura, 1990; Morimoto et al., 2006), although the Bengalese finch as a domesticated passerine can breed throughout the year. Generally, the seminal glomera (and probably the ejaculatory duct) of passerines increases in size in parallel with spermatogenesis and stores plenty of sperm for copulation during the breeding period (Bailey, 1953; Nakamura, 1990); but thereafter it shrinks (Chiba and Nakamura, 2003), and a variable amount of sperm may remain un-ejaculated in the organ, and therefore needs to be absorbed prior to the next breeding season. Thus, it seems reasonable to suggest that the seminal glomera together with the ejaculate duct also serve as the main site for removing or ingesting the remaining and infertile (including immotile and inactive) sperm.

As mentioned above, the epithelial spermiophagy in the seminal tract tended to be more active towards the posterior portion of the seminal glomera and the ejaculate duct, at least in the alpine accentor and the red-flanked bush robin. The results may explain the fact that in the zebra finch (*Taeniopygia guttata*) the number of morphologically abnormal sperm in the seminal glomera decreases towards the ejaculatory duct, showing a clear pattern of sperm quality across the proximal, mid, and distal regions of the seminal glomera (Birkhead et al., 1995). On the other hand, no clear difference was found in the incidence of epithelial spermiophagy

phagy among the three species with different intensities or levels of sperm competition, although it tends to be more active towards the terminal portion of the seminal tract in the alpine accentor and the red-flanked bushrobin in contrast to varied activity in the Bengalese finch. Currently, it is premature to determine whether the present data support the prediction (see Introduction) deduced from the studies on sperm quality and sperm competition (Birkhead et al., 1995, 2007; Calhim et al., 2007; Immler et al., 2008), or not.

In spite of the biological significance of spermiophagy, its physiological mechanism remains obscure and should be analyzed in future studies by experimental approaches as well as from the viewpoints of reproductive and evolutionary biology.

ACKNOWLEDGMENTS

This research was partly supported by a Grant-in-Aid (No. 20770018) from the Grants-in-Aid for Young Scientists (B) of The Ministry of Education, Culture, Sports, Science, and Technology, (2008-2010) to G. Morimoto and by the Research Promotion Grant (NDFU-09-17) from Nippon Dental University to A. Chiba.

REFERENCES

- Abou-Elmagd A, Wrobel KH (1990) The epithelial lining of the bovine ejaculatory duct. *Acta Anat (Basel)* 139: 60–65
- Aire TA (2000) Active spermiophagy in the initial part of the proximal efferent duct of the epididymis of normal domestic fowl (*Gallus domesticus*). *Res Vet Sci* 68:135–140
- Aire TA (2007) Anatomy of the testis and male reproductive tract. In “Reproductive Biology and Physiology of Birds, Vol 6A” Ed by BGM Jamieson, Science Publisher, Enfield, NH, pp 37–113
- Akbarsha MA, Kadalmani B, Tamilarasan V (2007) Efferent ductules of the fan-throated lizard *Sitana ponticeriana* Cuvier: light and transmission electron microscopy study. *Acta Zool (Stockh)* 88: 265–274
- Bailey RE (1953) Accessory reproductive organs of male fringillid birds: seasonal variations and response to various sex hormones. *Anat Rec* 115: 1–20
- Bedford JM (1979) Evolution of the sperm maturation and sperm storage function of the epididymus. In “The Spermatozoon” Ed by DW Fawcett, JM Bedford, Urban and Schwarzenberg, Baltimore/Munich, pp 7–21
- Birkhead TR (1991) Sperm depletion in the Bengalese finch *Lonchura striata*. *Behav Ecol* 2: 267–275
- Birkhead TR, Briskie JV, Møller AP (1993) Male sperm reserves and copulation frequency in birds. *Behav Ecol Sociobiol* 32: 85–93
- Birkhead TR, Fletcher F, Pellant EJ, Staples A (1995) Ejaculate quality and the success of extra-pair copulations in the zebra finch. *Nature* 377: 422–423
- Birkhead TR, Giusti F, Immler S, Jamieson BGM (2007) Ultrastructure of the unusual spermatozoon of the Eurasian bullfinch (*Pyrrhula pyrrhula*). *Acta Zool [Stokh]* 88: 119–128
- Calhim S, Immler S, Birkhead TR (2007) Postcopulatory sexual selection is associated with reduced variation in sperm morphology. *PLoS ONE* 2(5): e413
- Chiba A, Nakamura M (2003) Anatomical and histophysiological characterization of the male cloacal protuberance of the polygynandrous alpine accentor *Prunella collaris*. *Ibis* 145: E83–E93
- Goyal HO (1982) Light microscopic and ultrastructural evidence of epithelial phagocytosis of sperm in the rete testis and ductuli efferentes in the bull. *Am J Vet Res* 43: 785–790
- Holstein AF (1978) Spermatophagy in the seminiferous tubules and excurrent ducts of the testis in rhesus monkey and in man. *Andrologia* 10: 331–352
- Immler S, Calhim S, Birkhead TR (2008) Increased postcopulatory sexual selection reduces the intra-male variation in sperm design. *Evolution* 62: 1538–1543
- Kirby JD, Froman DP, Angel HR, Bernier PE, Hess RA (1990) Decreased spermatozoal survivability associated with aberrant morphology of the ductuli efferentes proximales of the chicken (*Gallus domesticus*). *Biol Reprod* 42: 383–389
- Lake PE (1981) Male genital organs. In “Form and Function in Birds, Vol 2” Ed by AS King, J McLelland, Academic Press, New York, pp 1–61
- Middleton ALA (1972) The structure and possible function of the avian seminal sac. *Condor* 74: 185–190
- Morimoto G, Yamaguchi N, Ueda K (2006) Plumage color as a status signal in male-male interaction in the red-flanked bushrobin *Tarsiger cyanurus*. *J Ethol* 24: 261–266
- Murakami M, Sugita A, Shimada T, Yoshimura T (1978) Scanning electron microscope observation of the seminal vesicle in the Japanese monkey with special reference to intraluminal spermiophagy by macrophages. *Arch Histol Jpn* 41: 275–283
- Murakami M, Nishida T, Shiromoto M, Inokuchi T (1985) Phagocytosis of spermatozoa and latex beads by epithelial cells of the ampulla vas deferentis of the rabbit: a combined SEM and TEM study. *Arch Histol Jpn* 48: 269–277
- Nakai M, Hashimoto Y, Kitagawa H, Kon Y, Kudo N (1989) Histological study on seminal plasma absorption and spermiophagy in the epididymal region of domestic fowl. *Poult Sci* 68: 582–589
- Nakamura M (1990) Cloacal protuberance and copulatory behavior of the alpine accentor (*Prunella collaris*). *Auk* 107: 284–295
- Porawski M, Wassermann GF, Achaval M (2004) Localization of acid phosphatase activity in the testes of two teleostean species (*Oreochromis niloticus* and *Odonthestes perugiae*). *Braz J Biol* 64: 853–858
- Samour JH, Moore HDM, Bailey IT, Watson PF (1987) Annual testicular cycle in budgerigars (*Melopsittacus undulatus*). *J Zool Lond* 212: 465–473
- Sever DM (1992) Spermiophagy by the spermathecal epithelium of the salamander *Eurycea cirrigera*. *J Morphol* 212: 281–290
- Tingari MD (1972) The fine structure of the epithelial lining of the excurrent duct system of the testis of the domestic fowl (*Gallus domesticus*). *Quart J Exp Physiol* 57: 271–295
- Tingari MD, Lake PE (1972) Ultrastructural evidence for resorption of spermatozoa and testicular fluid in the excurrent ducts of the testis of the domestic fowl, *Gallus domesticus*. *J Reprod Fertil* 31: 373–381

(Received October 4, 2010 / Accepted February 9, 2011)