

Chromosome polymorphism in Polish populations of northern birch mouse *Sicista betulina* 1

Authors: Fedyk, Stanisław, Chętnicki, Włodzimierz, Ruprecht, Andrzej L., and Cichocki, Jan

Source: *Folia Zoologica*, 60(1) : 31-36

Published By: Institute of Vertebrate Biology, Czech Academy of Sciences

URL: <https://doi.org/10.25225/fozo.v60.i1.a5.2011>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Chromosome polymorphism in Polish populations of northern birch mouse *Sicista betulina*¹

Stanisław FEDYK², Włodzimierz CHĘTNICKI², Andrzej L. RUPRECHT³ and Jan CICHOCKI⁴

² Institute of Biology, University of Białystok, Świerkowa 20B, 15-950 Białystok, Poland; e-mail: fedyk@uwb.edu.pl

³ ul. Polna 12A m. 27, 87-720 Ciechocinek, Poland

⁴ Department of Zoology, University of Zielona Góra, Prof. Z. Szafrana 1, 65-516 Zielona Góra, Poland

Received 17 March 2010; Accepted 18 June 2010

Abstract. Somatic chromosomes of 17 northern birch mice, *Sicista betulina*, originating from lowland and Tatra Mountain populations were studied. In the whole studied material constant diploid number of chromosomes ($2n = 32$) was found. Polymorphism of a pair of large-sized autosomes was found; acrocentric, subtelocentric and submetacentric chromosomes in homozygous or heterozygous state form the polymorphic pair. Consequently, chromosome arm number (NF) varies within 60 and 62. In this regard Polish populations of northern birch mouse differ from those of more eastward distribution (NF = 63–64).

Key words: variability of karyotype, chromosome evolution

Introduction

Genus *Sicista* is characterised by well-differentiated karyotype. Variability in the number of chromosomes in species belonging to this genus is $2n = 16–50$ (Table 1). Chromosomes of *Sicista betulina* (Pallas, 1779) were studied in many populations in various parts of species range thus far. The first description of the northern birch mouse's karyotype ($2n = 32$) was based on the material from Białowieża Primeval Forest (Poland) (Walknowska 1960). Karyotypes with 32 chromosomes were also confirmed in various populations of northern birch mouse occurring in European part of former The Union of Soviet Socialist Republics (USSR) as well as in Siberia (Vorontsov & Malygina 1973, Sokolov et al. 1987, Baker et al. 1996). Different chromosome arm numbers (NF) were given; extreme NF or NFa values were 48 (Vorontsov & Malygina 1973) and 64 (Sokolov et al. 1987), respectively. These discrepancies were rather a consequence of a poor quality of chromosome preparations used in karyotype examination and arrangement than the reflection of the real differences in chromosomes morphology.

Chromosome polymorphism was recorded in three *Sicista* species (see Table 1). In *S. subtilis*, three pairs of autosomes have different centromere localization and homo- and heterozygotic morphs occur in populations (Sokolov et al. 1986a). In *S. severtzovi*, several heteromorphic chromosome pairs were described as well as variation in the chromosome number ($2n = 16–22$) was observed (Sokolov et al. 1986a, 1987, Zagorodniuk & Kondratenko 2000). Karyotype of *S. betulina* was for a long time considered to be monomorphic (Walknowska 1960, Vorontsov & Malygina 1973, Sokolov et al. 1987, Baker et al. 1996); although identification of homologous chromosomes made repeated problems. Sokolov et al. (1989) were the first ones that noticed the presence of distinct differences in the localization of centromeres on a pair of large-sized autosomes.

In the present work, we demonstrate karyotype structure and variation in a few Polish populations of *S. betulina*. Three specimens out of the material used in the present work were previously analysed and described by Matan (2007).

¹The work is dedicated to the memory of Professor Zdzisław Pucek

Table 1. Survey of karyotype variability in the genus *Sicista*.

Species/subspecies	Karyotype characteristics*	Remarks	Source
<i>S. severtzevi</i> Ognev, 1935	2n = 16-22; NF = 28-30; X acro (m) Y acro (s)	Originally described as ssp of <i>S. subtilis</i> ; three unpaired autosomes which differ in size and morphology; number of acroc. autosomes 6-8	1,6,7
<i>S. subtilis</i> sensu lato	2n = 17; NF = 29; X submeta (m)?; Y (d)	This karyotype form probably relate to <i>S. severtzevi</i> . One unpaired autosome	2
<i>S. subtilis subtilis</i> Pallas, 1773	2n = 24; NF = 41-44; X acro (s); Y (d)	Three pairs (8, 9, 10) polymorphic (acro and subtelo centric morphs)	1
<i>S. s. sibirica</i> Ognev, 1935	2n = 24; NF = 44-45; X acro (s); Y (d)	Karyotype similar to <i>S. subtilis subtilis</i> ; two pair (8, 9) polymorphic acro and subtelo morphs)	1
<i>S. s. vaga</i> Pallas, 1778	2n = 24; NF = 41-42; X acro (m); Y acro (s)	Karyotype similar to <i>sibirica</i> and <i>subtilis</i> ; two pair (8, 9) polymorphic	1
<i>S. s. nordmanni</i> Keyserling & Blasius, 1840	2n = 26; NF = 48; X acro (m); Y (d)	Five large-sized pairs similar to the preceding subspecies with 2n = 24	1
<i>S. kluchorica</i> Sokolov et al., 1980	2n = 24; NF = 44; X acro (m); Y acro (s)		3
<i>S. caucasica</i> Vinogradov, 1925	2n = 32; NF = 48; X acro (m); Y (d)		3
<i>S. tianshanica</i> Salensky, 1903	2n = 32; NF = 54; X acro (m); Y acro (s)		3
<i>S. betulina</i> Pallas, 1779	2n = 32; NF = 63-64; Y (d)	Different size and morphology of X chromosome were reported	see text
<i>S. armenica</i> Sokolov & Baskevich, 1988	2n = 36; NF = 52; X acro (b); Y (d)		4
<i>S. kazbegica</i> Sokolov, Baskevich & Kovalskaya, 1986	2n = 40, 42; NF = 50, 52; X acro (m); Y (d)	The two chromosome types have probably parapatric distribution in Caucasus	5
<i>S. napaea</i> Hollister, 1912	2n = 42; NF = 52; X acro (m); Y acro (s)		3
<i>S. pseudonapaea</i> Strantman, 1949	2n = 44; NF = 52; X acro (m); Y acro (s)		3
<i>S. caudata</i> Thomas, 1907	2n = 50; NF = 50; X acro (l); Y acro (s)		3
<i>S. strandi</i> Formosov, 1931	2n = 44; NF = 52; X acro (l); Y (d)	Originally described as <i>S. betulina</i> "B form"	3

* Size of chromosome: (l) large; (m) medium; (s) small; (d) dot-like chromosome.

Source: 1 – Sokolov et al. 1986a, 2 – Zagorodniuk & Kondratenko 2000, 3 – Sokolov et al. 1987, 4 – Sokolov & Baskevich 1988, 5 – Sokolov et al. 1986b, 6 – Aniskin et al. 2003, 7 – Kovalskaya et al. 2000.

Table 2. The morphology of polymorphic chromosomes of the pair 2 of *Sicista betulina*.

Geographic region	Population	Geographic coordinates	No. of animals	Sex	Chromosome morph (pair No. 2)	
Masurian Lake Region	Tylkówko	20°74' E; 53°64' N	4572	f	st/st	
			4592	f	st/st	
			4607	m	a/a	
	Michałki	20°76' E; 53°65' N	20°76' E; 53°67' N	4593	m	st/st
				4606	m	st/st
				4608	f	st/st
				4609	f	st/st
	Zgniłocha	20°34' E; 53°33' N	20°34' E; 53°31' N	4519	m	a/a
				4520	f	a/st
				4652	m	a/a
Białowieża Primeval Forest	Białowieża	23°49' E; 52°42' N	4653	m	sm/sm	
			4555	m	st/sm	
Tatra Mountains	Polana Biały Potok	19°95' E; 49°30' N	4688	m	a/sm	
			4691	m	sm/sm	
			4692	m	st/st	
			4695	f	st/sm	
			4704	f	sm/sm	




Fig. 1. The homozygous (a/a) G-banded karyotype.




Fig. 3. The heterozygous (sm/a) karyotype.




Fig. 2. The heterozygous (st/sm) karyotype.

Material and Methods

Material for chromosome studies consisted of 17 individuals originating from six lowland populations and one high mountain population (Table 2). Chromosome preparations were made from spleen and/or bone marrow by using standard method *in vivo*

(Fedyk 1980). G-bands were obtained after trypsin digestion of chromosomes and staining with Giemsa buffered solution according to Seabright (1971). No karyotype standard of *S. betulina* has been proposed thus far. Authors of previous papers arranged karyotypes based on the size of individual chromosomes (Walknowska 1960, Sokolov et al. 1982, Baker et al. 1996) or they identified chromosome groups depending on the centromere localization (Vorontsov & Malygina 1973, Sokolov et al. 1987, 1989, Baskevich & Okulova 2003). A low quality of chromosome preparations yielding poorly visible localization of the centromere can explain such huge diversity. In the present work, we recommend to use the karyotype of *S. betulina*, published by Sokolov et al. (1989) as a standard.

Results

All the studied individuals had $2n = 32$ chromosomes. Within large chromosomes, pairs 1, 4 and 5 are

metacentric chromosomes; pair 3 is submetacentric, and pair 2 is polymorphic and it is composed of uniarmed (acrocentric) or biarmed chromosomes (Figs. 1–3). Analysis of the arms ratio on biarmed chromosomes demonstrated that there are two distinct variants, which differ by centromere localization. Chromosomes with more proximal localization of centromeres have the ratio 2.13 to 3.20 between the longer and the shorter arm ($r = l/s$); however, the arms ratio on chromosomes with more distal localization of the centromere was between 4.48 and 6.92. These two chromosome variants were classified as submetacentric (sm) and subtelocentric (st), respectively. According to the formal definition (Levan et al. 1964), sm chromosomes have $r \leq 3$, and st chromosomes have $r \geq 3$. Correctness of this differentiation between observed variants is confirmed by the r values (Fig. 4), which do not overlap each other.




Fig. 4. The arm length ratio ($r = l/s$) on biarmed chromosomes of the autosomal pair 2.




Fig. 5. G-banding patterns in three variants (a, st, sm) of the polymorphic autosomal pair 2.

The second size group consists of five pairs (6–10) of submetacentric chromosomes and a pair 11 of acrocentric chromosomes. The third size group consists of four pairs (12–15) of small meta- and submetacentric chromosomes. Small differences in ratio of chromosome arms in various metaphase cells were observed within chromosomes of the second and third group. As the differences were found not only between individuals but also between metaphase spreads coming from the same individual, this differentiation was not taken into account. A subtelocentric X chromosome is of similar size as the autosomes of pairs 6 and 7. The Y chromosome

is the smallest chromosome in the karyotype. The chromosome arm numbers (NF) vary between 60 and 62 depending on the morphology of chromosomes of the pair 2.

The G-banding pattern well characterizes the individual pairs of chromosomes, including the sex chromosomes (Fig. 3). It should be noted that additional bands can occur on short arms of st and sm chromosomes from the pair 2 in comparison with acrocentric chromosomes (Fig. 5).

Discussion

The authors of all previous publications reported unanimously that *S. betulina* has $2n = 32$ chromosomes. The only exception is the Kursk area where the “form B” of *S. betulina* with $2n = 44$ was found (Sokolov et al. 1987). Further studies revealed that the “form B” is a different species, *Sicista strandi* Formosov, 1931 distributed allopatrically in respect of *S. betulina* (Sokolov et al. 1989, Zagorodniuk 2007).

No polymorphism of autosomes has been detected in the early studies of *S. betulina*. In addition, previous authors achieved only very poor quality chromosomal preparations. Therefore, not only the arrangement of chromosomal pairs but also the determination of the X chromosome was wrong in many publications. Application of the outdated “squash” method in pioneering work gave preparations with poor morphology of chromosomes (Walknowska 1960). Usage of the air-drying preparation method significantly improved the results. However, Vorontsov & Malygina (1973) did not provide a detailed morphology of *S. betulina* chromosomes as well. Some of chromosomes were described as “small, near acrocentric elements”, and as a consequence, the number of autosomal arms ($NFa = 48$) was underestimated. Poor quality of chromosome preparations also did not enable the correct determination of the X chromosome (Vorontsov & Malygina 1973). Sokolov et al. (1982) obtained better quality chromosome preparations from northern birch mice collected near Moscow, but they also arranged the karyotype incorrectly. A subtelocentric chromosome from the heteromorphic pair (pair 2 at present work) was determined as the X chromosome and only biarmed chromosomes were described in the karyotype ($2n = 32$, $NF = 64$). In a comprehensive study on the material collected from Moscow, Kurgan, Novosibirsk, Kemerowo, Tomsk and Krasnojarsk regions and from the former Buryat ASSR, Sokolov et al. (1987) gave the same characteristics ($2n = 32$, $NF = 64$) and again incorrectly determined the sex chromosomes. Chromosomes of pair 4 (according

to the nomenclature used in the present paper) were determined as the X chromosomes, however, there is no mention of polymorphism, though the heteromorphic pair (sm/st) of large autosomes is clearly visible in the published karyotype. Sokolov et al. (1989) realized that pair 2 was polymorphic and they re-analyzed material from previous publications (28 individuals from 13 populations from the European part of the former USSR and Siberia). The authors described two chromosome variants (sm and st) of the pair 2. The sm chromosomes were more common and they were found in homozygous state (sm/sm) in 15 individuals originating mostly from the European part of species range. The st chromosomes occur mostly in Siberia, where they are rare and were found in heterozygous state (sm/st) in 13 individuals. These data confirm our observations of geographic differentiation of polymorphic morph frequency; st chromosomes occur more frequently on the lowlands in Poland, whereas sm chromosomes prevail in the mountain population. In our material, the frequency of the acrocentric chromosome variant is 0.23 in the whole sample (Table 2). It is surprising that this variant was not identified in the earlier paper, as an acrocentric chromosome in heterozygous state is apparent in the complement of individuals from the former Buryat ASSR (see Sokolov et al. 1989 – Fig. 2).

The karyotype of *S. betulina* reported by Sokolov et al. (1989) was used as a standard in the present study. However, it must be emphasized that Sokolov et al. (1989) determined the X chromosome as subtelocentric, with its size similar to the smallest autosomes (pairs 13–15), whereas in the Polish populations (excluding of Walknowska 1960), X chromosome is reported as submetacentric of comparable size to pairs 6 and 7. The number of chromosome arms also differs. All the autosomes are considered biarmed in Sokolov et al. (1989), whereas pair 11 is evaluated as acrocentric in Polish populations. Baker et al. (1996) described at Chernobyl, a region close to the eastern border of Poland, a karyotype, which is the most similar to our findings. Pair 2 is heteromorphic (sm/st) and one of the autosomal pairs is acrocentric. The centromere position on the X chromosome is not clear but the

relative size of this chromosome is comparable with X chromosome from our material. Baskevich & Okulova (2003) on the basis of the G-banding spreads, determined X chromosome of similar size in the material from Moscow and Riazian environs. However, it should be mentioned that G-banding karyotype presented in their work is not similar to that presented in this paper (Fig. 1).

G-banding pattern (Fig. 5) suggests that shorter arms of st and sm chromosomes of pair 2 originated by chromatin addition rather than pericentric inversion. The presence of large heterochromatin blocks (C bands) in the centromeric region of the autosomal pair 2 (Baskevich & Okulova 2003) supports this hypothesis. It is surprising that the G-banding pattern of northern birch mice from Moscow region (Baskevich & Okulova 2003) is totally incomparable with that presented in our study (Fig. 3).

Polymorphism similar to *S. betulina* was recorded also in *S. subtilis*. Variants a, st and sm of three autosomal pairs were found in three subspecies with $2n = 24$ (*S. subtilis subtilis*, *S. s. vaga*, and *S. s. sibirica*). It was suggested that this polymorphism resulted from pericentric inversions. The fourth subspecies (*S. s. nordmanni*) differs from the above-mentioned with its chromosome number, which is $2n = 26$ (Sokolov et al. 1986a, Baskevich et al. 2005). Generally, based on the obtained results, it could be hypothesised that there is possible polymorphism of more than one pair of chromosomes that exists in *S. betulina*.

Acknowledgements

These studies were supported by the University of Białystok (BST 113). We greatly acknowledge Prof. L. Rychlik and Prof. J.R.E. Taylor for providing us S. betulina from Białowieża Primeval Forest, as well as Dr. S. Pavlova for providing us some Russian papers. We are also thankful to two anonymous reviewers for their valuable comments, as well as to Mr. A. Antol and Mr. T. Złotkowski for help in animal catches in the Zakopane region. Besides, we are thankful to Ms. K. Kwasowska and Mr. S. Mączewski for help in the laboratory works and birch mice catches in the Mazurian Lake Region.

Literature

- Aniskin V.M., Gorelov Y.K., Kovalskaya Y.M. & Shapovalov A.S. 2003: Karyotype variability of *Sicista severtzovi severtzovi* (Rodentia): toward gene pool conservation of small populations. *Abstracts of the Conference "Theriofauna of Russia and adjacent territories"*. RAS Press: 19–20. (in Russian)
- Baker R.J., Hamilton M.J., van den Bussche R.A., Wiggins L.E., Sugg D.W., Smith M.H., Lomakin M.D., Gaschak S.P., Bundova E.G., Rudenskaya G.A. & Chesser R.K. 1996: Small mammals from the most

- radioactive sites near the Chernobyl nuclear power plant. *J. Mammal.* 77: 155–170.
- Baskevich M.I. & Okulova N.M. 2003: Comparative karyology and craniology in birch mice of the group “*betulina*” (Rodentia, Dipodidae, *Sicista*). *Zool. Zh.* 82: 996–1009. (in Russian with English summary)
- Baskevich M.I., Kozlovsky A.I., Oparin M.L., Savinetskaya L.E., Cherepanov E.V., Sokolenko O.V. & Khlyap L.A. 2005: Karyological aspects of the Lower-Volga rodent fauna. *Povolzhski Ekologicheski Zhurnal* 3: 197–206. (in Russian with English summary)
- Fedyk S. 1980: Chromosome polymorphism in a population of *Sorex araneus* L. at Białowieża. *Folia Biol. (Kraków)* 28: 83–120.
- Kovalskaya Y.M., Tikhonov I.A., Tikhonova G.N., Surov A.V. & Bogomolov P.L. 2000: New geographical localities of chromosome forms of southern birch mouse (*subtilis* group) and description of *Sicista severtzovi cimlanica* subsp. n. (Mammalia, Rodentia) from the middle Don River basin. *Zool. Zh.* 79: 954–964. (in Russian with English summary)
- Levan A., Fredga K. & Sandberg A.A. 1964: Nomenclature for centromeric position on chromosomes. *Hereditas* 52: 201–220.
- Matan K. 2007: Chromosomes of *Sicista betulina*. MSc thesis, Białystok University. (in Polish)
- Seabright M. 1971: A rapid banding technique for human chromosomes. *Lancet* 2: 971–972.
- Sokolov V.E. & Baskevich M.I. 1988: A new species of birch mouse – *Sicista armenica* sp. n. (Rodentia, Dipodoidea) from lesser Caucasus. *Zool. Zh.* 67: 300–304. (in Russian with English summary)
- Sokolov V.E., Baskevich M.I. & Kovalskaya Y.M. 1986a: The karyotype variability in the southern birch mouse (*Sicista subtilis* Pallas) and substantiation of the species validity of *S. severtzovi*. *Zool. Zh.* 65: 1684–1692. (in Russian with English summary)
- Sokolov V.E., Baskevich M.I. & Kovalskaya Y.M. 1986b: *Sicista kazbegica* sp. n. (Rodentia, Dipodidae) from the basin of the Terek River upper reaches. *Zool. Zh.* 65: 949–951. (in Russian with English summary)
- Sokolov V.E., Kovalskaya Y.M. & Baskevich M.I. 1982: Taxonomy and comparative cytogenetics of some species of the genus *Sicista* (Rodentia, Dipodidae). *Zool. Zh.* 61: 102–108. (in Russian with English summary)
- Sokolov V.E., Kovalskaya Y.M. & Baskevich M.I. 1987: Review of karyological research and the problems of systematics in the genus *Sicista* (Zapodidae, Rodentia, Mammalia). *Folia Zool.* 36: 35–44.
- Sokolov V.E., Kovalskaya Y.M. & Baskevich M.I. 1989: On species status of the northern birch mouse *Sicista strandi* (Rodentia, Dipodidae). *Zool. Zh.* 68: 95–106. (in Russian with English summary)
- Walknowska J. 1960: Les chromosomes chez *Sicista betulina* Pall. *Folia Biol. (Kraków)* 8: 65–70.
- Vorontsov N.N. & Malygina N.A. 1973: Karyological studies in jerboas and birch mice (Dipodoidea, Rodentia, Mammalia). *Caryologia* 26: 193–212.
- Zagorodniuk I.V. 2007: Allospecies of rodent group *Sicista* “*betulina*”: spatial relationship from the viewpoint of limiting similarity concept. *Visnyk of the Dnipropetrovsk University, Ser. Biol. & Ecology* 15: 45–53. (in Ukrainian with English summary)
- Zagorodniuk I.V. & Kondratenko O.V. 2000: *Sicista severtzovi* and its relatives in rodent fauna of Ukraine: cytogenetic and biogeographical analysis. *Vestnik Zoologii* 15: 101–107. (in Ukrainian with English summary)