

## **Status and distribution of spined loaches (Cobitidae) and stone loaches (Nemacheilidae) in Bosnia and Herzegovina**

Authors: Tutman, Pero, Buj, Ivana, Čaleta, Marko, Hamzić, Adem, Korjenčić, Enad, et al.

Source: Folia Zoologica, 66(4) : 211-226

Published By: Institute of Vertebrate Biology, Czech Academy of Sciences

URL: <https://doi.org/10.25225/fozo.v66.i4.a2.2017>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Status and distribution of spined loaches (Cobitidae) and stone loaches (Nemacheilidae) in Bosnia and Herzegovina

Pero TUTMAN<sup>1\*</sup>, Ivana BUJ<sup>2</sup>, Marko ČALETA<sup>3</sup>, Adem HAMZIĆ<sup>4</sup>, Enad KORJENIĆ<sup>4</sup>,  
Avdul ADROVIĆ<sup>5</sup> and Branko GLAMUZINA<sup>6</sup>

<sup>1</sup> Institute of Oceanography and Fisheries, Šetalište Ivana Meštrovića 63, 21000 Split, Croatia;  
e-mail: tutman@izor.hr

<sup>2</sup> University of Zagreb, Faculty of Science, Rooseveltov trg 6, 10000 Zagreb, Croatia;  
e-mail: ivana.buj@biol.pmf.hr

<sup>3</sup> University of Zagreb, Faculty of Teacher Education, Savska cesta 77, 10000 Zagreb, Croatia;  
e-mail: marko.caleta@ufzg.hr

<sup>4</sup> University of Sarajevo, Faculty of Science, Zmaja od Bosne 33-35, 71000 Sarajevo,  
Bosnia and Herzegovina; e-mail: haadem@hotmail.com, ekorjenic@pmf.unsa.ba

<sup>5</sup> University of Tuzla, Faculty of Science, Univerzitetska 4, 75000 Tuzla, Bosnia and Herzegovina;  
e-mail: avdul.adrovic@untz.ba

<sup>6</sup> University of Dubrovnik, Department for Aquaculture, Ćira Carića 4, 20000 Dubrovnik, Croatia;  
e-mail: branko.glamuzina@unidu.hr

Received 14 July 2017; Accepted 10 January 2018

**Abstract.** This paper presents a critical assessment of the current status and geographical distribution of loaches (Cobitidae, Nemacheilidae) in Bosnia and Herzegovina. This closes a gap in research knowledge with only general distribution data currently available. In the Danube catchment four cobitid (*Cobitis elongata*, *C. elongatoides*, *Misgurnus fossilis*, *Sabanejewia balcanica*) and one nemacheilid species (*Barbatula barbatula*) were recorded, while the Adriatic Sea catchment hosts three spined loaches: *C. narentana*, *C. illyrica*, and *C. herzegoviniensis* and nemachilid *B. barbatula*. *Cobitis elongatoides* was found to be the most common and distributed throughout the northern Bosnia and Herzegovina; i.e. in the rivers Sava, Drina, Bosna, Vrbas, Una (Danube catchment). *Sabanejewia balcanica* was recorded in the same area but less frequently. Recent bibliographic evidence forces us to exclude *C. taenia* from the current checklist. On the other hand, cobitid species from the Adriatic Sea catchment are endemic with a very small distribution. Also, the extinction risk status of each taxon is mentioned following IUCN categories. This study provides geographically constrained information on the recent status of loach populations in the freshwaters of Bosnia and Herzegovina. Further data on the status and geographical distribution would improve the conservation and management of these fishes.

**Key words:** loaches, Sava River catchment, Adriatic Sea catchment, species composition, distribution area

## Introduction

Spined loaches of the family Cobitidae and stone loaches of the family Nemacheilidae are large groups of primarily freshwater fishes which are very common and widespread in waters of the Palearctic region (Nalbant et al. 2001). Their distribution and range encompasses Eurasia (including England and the Iberian Peninsula), the Japanese archipelago, Korean Peninsula and Amur River basin in the Far East, as well as northern Africa, making them one of the most widespread families of European freshwater fish (Bohlen & Ráb 2001,

Kottelat & Freyhof 2007). Members of Cobitidae and Nemacheilidae are small sized bottom-dwellers characterized by low dispersal abilities, elongated body and short lifespan. Their preferred microhabitat characteristics are open, shallow areas with slow-flowing or stagnant water (Kottelat & Freyhof 2007). Due to the specific habitat requirement and the fact that they are not commercially important, data on their distribution and ecology are relatively poor and until recently, they have been overlooked in scientific study (Mustafić et al. 2003, Šanda et al. 2008a, b).

\* Corresponding Author

In recent years, loaches have become the object of more focused investigations. Their peculiar ecology, unresolved taxonomy and unusual reproduction modes attracted the attention of ichthyologists. Recent phylogenetic studies have revealed significant phylogenetic relationships within the genus *Cobitis* and demonstrated an unexpectedly high diversity within this fish group in Europe (Perdices & Doadrio 2001, Janko et al. 2005a, b, Bohlen et al. 2006, Buj et al. 2008, 2014, 2015a, Perdices et al. 2008, 2016, Geiger et al. 2014). Genetic diversity and phylogenetic relationships have been studied on *Sabanejewia* (Perdices et al. 2003, Marešová et al. 2011) and *Barbatula* populations (Šedivá et al. 2008). However, although both loach families occur in many European waters, scientific knowledge on this species in Bosnia and Herzegovina is still sparse. Consequently, the taxonomic status and geographical distribution of Cobitidae and Nemacheilidae in Bosnia and Herzegovina is poorly understood and uncertain. The published literature only comprises a few contributions on their general distribution in the basin of the River Sava (Danube catchment) (Brusina 1892, Medić 1901, Zaplata & Taler 1932, Karaman 1963, Vuković 1963, 1977, Bogut et al. 2006, Sofradžija 2009) and a very general note from the River Neretva drainage system (Vuković 1963, 1977, Bogut et al. 2006, Sofradžija 2009) which belongs to the Adriatic Sea catchment. In addition to the sparse data on their distribution, there is a surprising deficiency in the knowledge on their biology and morphological characteristics in general. The available publications contain mainly general biological and ecological data (Vuković 1963, 1977, Vuković & Ivanović 1971, Bogut et al. 2006, Sofradžija 2009) and overall faunistic records (Kosorić 1978, Kosorić et al. 1983, Tutman et al. 2012a, Nedić et al. 2014). Several older papers concern systematic, genetic and chromosomal issues (Sofradžija & Berberović 1978, Sofradžija & Vuković 1979). There are also some data on age structure available from the Hutovo Blato wetland in the Neretva River catchment (Tutman et al. 2006, Dulčić et al. 2009). In fact, only Golub et al. (2016) coupled precise geographic location with morphometric measurements and meristic characteristics of the specimens caught. Generally, according to the last book of the freshwater fishes of Bosnia and Herzegovina available (Sofradžija 2009) five cobitid species of three genera (*Cobitis*, *Misgurnus* and *Sabanejewia*) and one from Nemacheilidae (*Barbatula*) are cited as extant in domestic waters, but without detailed data of their

distribution, ecology or biology. However, new results support differentiation of the Adriatic basin *Cobitis* and their independent evolution from the Central European group (Buj et al. 2008, 2014, 2015a, Perdices et al. 2008, 2016), revealing two new species (Freyhof & Stelbrink 2007, Buj et al. 2014).

Detailed data on the distribution of loach species have been reported in recent years from several surrounding countries (Mrakovčić et al. 2000, Povž & Šumer 2000, 2003, Bohlen et al. 2003, Erős et al. 2003, Marić & Pavlović 2005, Šanda et al. 2008c, Marić & Milošević 2010). However, no general summary paper is available from Bosnia and Herzegovina on the current status and geographical distribution of populations of the particular loach species. The fact is that knowledge of the distribution and present status of populations, including the precise taxonomic identity of species of these families is required for clarifying the taxonomic status and geographical distribution. These data provides basic information for conservation management and also contributes to more detailed knowledge of loach distribution at the continental scale. Such a compilation is needed, since the general distribution patterns of species can have deep impacts both on national as well as on European conservation. Since there is little information available about the recent status and distribution, this paper reviews the existing data on taxonomy and distribution of spined loaches from the family Cobitidae and stone loaches from the family Nemacheilidae providing an important update of their present population status in Bosnia and Herzegovina. The aims of this paper are to update the existing information about the status and to investigate more fully the geographical distribution of loaches (Cobitidae, Nemacheilidae) in Bosnia and Herzegovina.

## Material and Methods

Faunistic data on loach species' presence and distribution in Bosnia and Herzegovina waters were based on published literature in the form of research articles, monographs, books and technical reports. To promote the protection and conservation of these species, this paper attempts to give a comprehensive review of available published, analysed together with more recent data gathered by the authors from 2000 until present. Also, some conservation management recommendations are proposed in order to improve their natural status. Additional data, like information on the main threats are derived from different ichthyological monitoring surveys carried out in the last twenty years in Bosnia and Herzegovina by the Faculty of Science, University of Sarajevo.

Rivers in Bosnia and Herzegovina are divided into two separate drainage systems: the Danube (Black Sea) (38719 km<sup>2</sup> or 75.7 % of total Bosnia and Herzegovina surface area), and the Adriatic Sea catchment (12410 km<sup>2</sup> or 24.3 %), which are separated by the Dinaric mountain range (Fig. 1). Both catchments are characterised by a different ichthyofaunal composition: in general, the Danube

basin accommodates species with wider distribution ranges, while the Adriatic Sea catchment area is characterized by a number of endemic forms (Vuković 1977, Kottelat & Freyhof 2007, Sofradžija 2009). The main part of the Danube catchment area is drained by the River Sava, which runs towards the northern border of Bosnia and Herzegovina and Croatia, and its major tributaries (the rivers Una, Sana, Vrbas, Ukrina,

**Table 1.** List of loach species and their distribution within the river basins of Bosnia and Herzegovina based on available literature data.

Family/Species	Rivers – locality	River basin	Catchment	Source
<b>Cobitidae</b>				
<i>Cobitis elongata</i>	Una	Una	Danube	Brusina 1892, Medić 1901, Sofradžija 2009, Drešković et al. 2011
	Sana			Drešković et al. 2011
	Sava	Sava		Bogut et al. 2006, Sofradžija 2009, Drešković et al. 2011, Nedić et al. 2014, Simonović et al. 2015
	Brka, Tinja			Drešković et al. 2011
	Modrac Reservoir, Gostilja			Adrović et al. 2012
	Vrbas	Vrbas		Sofradžija 2009, Drešković et al. 2011
	Bosna	Bosna		Sofradžija 2009, Drešković et al. 2011
	Spreča			Sofradžija 2009
	Fojnica			Pavličević et al. 2014
	Kozica – Lašva river tributary			Pavličević et al. 2014
	Misoča			Bogut et al. 2006, Škrijelj et al. 2016
	Drina	Drina		Hamzić et al. 2015
<i>Cobitis elongatoides</i>	rivers and tributaries of Black Sea basin	Sava	Danube	Bogut et al. 2006
	Modrac Reservoir, Brka, Gostilja, Tinja			Adrović et al. 2012
	Sava – lower part			Sofradžija 2009, Nedić et al. 2014
	Vrbas – middle courses	Vrbas		Sofradžija 2009, Hamzić et al. 2015
	Miljacka, Zujevina	Bosna		Zaplata & Taler 1932
	Bosna – middle courses			Sofradžija 2009, Hamzić et al. 2015
	Spreča			Memić & Adrović 2010
	Tinja			Škrijelj et al. 2016
Drina – middle courses	Drina		Sofradžija 2009, Hamzić et al. 2015	
Una – middle courses	Una		Sofradžija 2009, Hamzić et al. 2015	
<i>Cobitis herzegoviniensis</i>	Mostarsko Blato	Neretva	Adriatic	Šanda et al. 2008a, Glamuzina et al. 2013, Buj et al. 2014, 2015a, b
<i>Cobitis illyrica</i>	Lake Krenica	Neretva	Adriatic	Šanda et al. 2008a, b, Glamuzina et al. 2013, Buj et al. 2014, 2015a, b, Čaleta et al. 2015
<i>Cobitis narentana</i>	lower River Neretva from Čapljina city	Neretva	Adriatic	Vuković 1963, 1977, Šanda et al. 2008a, b, Sofradžija 2009, Glamuzina et al. 2013, Buj et al. 2014, 2015a, b, Čaleta et al. 2015
	Hutovo Blato wetland			Tutman et al. 2006, 2012a, Šanda et al. 2008a, b, Glamuzina et al. 2013, Buj et al. 2014, 2015a, b, Čaleta et al. 2015
	rivers Bregava and Trebišnjica			Šanda et al. 2008a, b, Buj et al. 2014, 2015a, b

**Table 1.** Continued.

Family/Species	Rivers – locality	River basin	Catchment	Source
<i>Misgurnus fossilis</i>	Sava	Sava	Danube	Brusina 1892, Sofradžija 2009, Hamzić et al. 2015
	waters of Black Sea basin			Vuković 1963, 1977, Bogut et al. 2006
	River Sava and lower reaches of their tributaries			Sofradžija 2009
	Una – near Dubica	Una		Brusina 1892
	Bosna	Bosna		Sofradžija 2009, Drešković et al. 2011, Hamzić et al. 2015
	Krivaja			Muhamedagić et al. 2013
<i>Sabanejewia balcanica</i>	Drina	Drina		Sofradžija 2009
	waters of Black Sea basin	Sava	Danube	Vuković 1963, 1977, Bogut et al. 2006
	Sava			Hamzić et al. 2015, Simonović et al. 2015
	Gostilja			Adrović et al. 2012
	River Vrbas and tributaries Suturlija and Široka	Vrbas		Karaman 1963, Vuković 1977, Sofradžija 2009, Marešová et al. 2011, Hamzić et al. 2015
	Bosna	Bosna		Sofradžija 2009, Drešković et al. 2011
	Krivaja			Muhamedagić et al. 2013
	Usora			Marešová et al. 2011
	Drina – middle reaches	Drina		Sofradžija 2009
	Una – Kostajnica	Una		Drešković et al. 2011
Una – Bihać			Marešová et al. 2011	
Nemacheilidae				
<i>Barbatula barbatula</i>	River Bosna	Bosna	Danube	Zaplata & Taler 1932, Sofradžija 2009
	rivers of Danubian catchment	Sava		Vuković 1963, 1977, Bogut et al. 2006
	lower Sava River			Nedić et al. 2014, Hamzić et al. 2015
	upper and middle reaches of Sana	Una		Sofradžija 2009
	upper and middle reaches of Ukrina	Ukrina		Sofradžija 2009
	upper and middle reaches of Vrbas	Vrbas		Sofradžija 2009
	upper and middle reaches of Drina	Drina		Sofradžija 2009
	upper Neretva River	Neretva	Adriatic	Vegara et al. 2009

Bosna, Brka, Tinja and River Drina) that drain deeply into central and northern Bosnia and Herzegovina. They receive several mountain tributaries each. The River Sava is the main waterway connecting Croatia, Bosnia and Herzegovina and Serbia. Together, these tributaries constitute a major drainage basin of the southeastern Europe, covering the total area of approximately 97713.20 km<sup>2</sup> representing one of the most significant sub-basins of the Danube River basin, with a share of 12 % (Jukić 2008).

In contrast, the Adriatic Sea catchment area of Bosnia and Herzegovina comprises of two sub-basins; the Neretva and Cetina rivers. The only river that flows into the Adriatic Sea is the River Neretva. The

Neretva River basin, including the part of Trebišnjica River basin, has an area of 10100 km<sup>2</sup> or 81.4 % of the total Adriatic Sea catchment area of Bosnia and Herzegovina, while the Cetina river sub-basin extends over 2310 km<sup>2</sup> (18.6 % of the total Adriatic Sea catchment area of Bosnia and Herzegovina). In Herzegovina there is an extensive karst area (more than 4000 km<sup>2</sup>) with important groundwater sources, that occur as small springs, streams, rivers and lakes.

## Results

### *Diversity of loaches in Bosnia and Herzegovina*

Review of the published literature reveals that data on loach species' composition and geographical



**Fig. 1.** Map of Bosnia and Herzegovina showing main catchments, Danube (or Black Sea) catchment (unshaded) and Adriatic Sea catchment (shaded).

distribution are very scarce, mostly based on species lists only lacking detailed descriptions of species and localities. According to the bibliographic data, a total of nine species of loach (eight Cobitidae and one Nemacheilidae) were reported from freshwaters within their geographical range (Vuković 1963, 1977, Bogut et al. 2006, Šanda et al. 2008a, b, Sofradžija 2009, Buj et al. 2014, 2015a, b, Golub et al. 2016). However, some of the data are in obvious conflict with the current taxonomy. Based on all available data it is clear that reports on the presence of *Cobitis taenia* Linnaeus, 1758 in Bosnia and Herzegovina are a result of taxonomic imprecision and it cannot be expected that this species inhabits waters of Bosnia and Herzegovina, because its distribution range is restricted to northern Europe (Nalbant et al. 2001). Bosnia and Herzegovina was thought to be a part of this species' range since *C. taenia* was considered to have extremely wide distribution through Palearctic region (Nalbant et al. 2001). However, already Bacescu (1962) reported *C. taenia* to be an assemblage of morphologically similar, but phylogenetically distinct species.

The ichthyofauna of Bosnia and Herzegovina comprises eight loach species, distributed in both catchments. In the northern part (Sava River basin), five species were recorded: Balkan spined loach *C. elongata* Heckel & Kner, 1858, Danube spined loach *C. elongatoides* Bacescu & Mayer, 1969, weather loach *Misgurnus fossilis* (Linnaeus, 1758), Balkan golden loach *Sabanejewia balcanica* (Karaman, 1922) and one member of nemacheilid family: stone loach *Barbatula barbatula* (Linnaeus, 1758). For the waters of the Adriatic Sea catchment three species of Cobitidae were reported; two recently described: Illyric spined loach,

*C. illyrica* Freyhof & Stelbrink 2007; Herzegovinian spined loach, *C. herzegoviniensis* Buj & Šanda, 2014 and the Neretva spined loach, *C. narentana* Karaman 1928. From the Nemacheilidae family *B. barbatula* was recently reported in the upper part of the River Neretva (Vegara et al. 2009). Significantly more and newer data are available on the loaches in the Adriatic (Schneider et al. 2000a, b, Buj et al. 2008, 2014, 2015a, b) than for the Danube catchment. Reported species are not equally distributed across Bosnia and Herzegovina, but show a higher diversity in the north than in the south of the country. The only species shared among the both catchments is *B. barbatula* (Table 1).

#### Overview of the available literature

The earliest information comes from Brusina (1892) and Medić (1901), who indicated the presence of *B. barbatula* (as *Nemachilus barbatulus*) and *M. fossilis* in the River Sava, mentioning *C. taenia* and *B. barbatula* (as *Cobitis barbatula*), but only on the Croatian side. Later, Zaplata & Taler (1932) reported loach species in the River Bosna and its tributaries near Sarajevo recording the presence of *C. taenia* and *B. barbatula* (as *C. barbatula*). Karaman (1963) described a new subspecies *C. aurata bosniaca* from the River Vrbas which flows into the River Sava. In the first check-list, Vuković (1963) listed four species; *C. taenia*, *C. aurata*, *B. barbatula* (as *N. barbatulus*) and *M. fossilis* (as *M. fosillis*). *Cobitis taenia* was placed in the waters of both the Danube and Adriatic drainage systems with two subspecies; *C. taenia taenia* Linnaeus, 1758 in the Danube, and *C. taenia narentana* Karaman 1928 in the Adriatic basin. Vuković (1977) in a subsequent check-list just followed the previous one and mentioned *C. taenia* with the two same subspecies, *S. balcanica* (as *C. aurata*) with two subspecies; *C. aurata balcanica* Karaman, 1922 and *C. aurata bosniaca* Karaman 1963 (in the River Vrbas and their tributaries Suturlija and Široka), *B. barbatula* (as *N. barbatulus*) and *M. fossilis*, all in the waters of Danube catchment. Furthermore, Kosorić & Mikavica (1981) listed *C. taenia* in the River Krivaja (Danube basin), whilst Kosorić et al. (1983) for the River Bosna and Korjenić (2004) for the River Fojnica in middle Bosnia and Herzegovina. Sofradžija & Berberović (1978) investigated sexual dimorphism in *C. taenia*, while Sofradžija & Vuković (1979) presented the chromosomal set of *B. barbatula* (as *N. barbatulus*).

*Cobitis narentana* was the only *Cobitis* species that has been mentioned from the Neretva drainage (Karaman

1928, Taler 1953, Vuković 1977, Mrakovčić et al. 1995, 2000). Later phylogenetic research revealed their differentiation indicating independent evolution from the Central European group (Buj et al. 2008, 2014, 2015a, Perdices et al. 2008, 2016). In addition, two new species were described (Freyhof & Stelbrink 2007, Buj et al. 2014). Tutman et al. (2006) presented some biological parameters of *C. narentana* while Dulčić et al. (2009) reported a length-weight relationship of a *C. narentana* population from Hutovo Blato wetland (Neretva River basin). Furthermore, Tutman et al. (2012a) mentioned *C. narentana* in Hutovo Blato as a rare member of its ichthyofauna.

According to Bogut et al. (2006) the freshwaters of Bosnia and Herzegovina are inhabited by five loach species (*C. elongata*, *C. elongatoides*, *M. fossilis*, *S. balcanica* and *B. barbatula*), while Kottelat & Freyhof (2007) listed six, adding *C. narentana* to the list. A review of more recent studies shows that Sofradžija (2009), following the older systematics, indicated five species; *M. fossilis*, *B. barbatula* (as *N. barbatulus*), *C. elongatoides*, *C. elongata* for Danube and *C. narentana* for the Adriatic catchment and two subspecies (*C. aurata balcanica* Karaman, 1922 and *C. a. bosniaca* Karaman 1963), both in the waters of Danube basin. However, these data were presented without precise location of their distribution. Actually, Šanda et al. (2008a, b) were the first ones to analyze in detail the occurrence and the distribution of *Cobitis* species in the waters of Adriatic Sea catchment in Bosnia and Herzegovina. Further intensive research into the Cobitidae family in the last 15-20 years in the Neretva river catchment brought many crucial findings, which radically changed original insights on species structure (Schneider et al. 2000a, b, Buj et al. 2008, 2014, 2015a). Recently, data from different ichthyological monitoring surveys have provided more precise information on distribution ranges. Škrijelj et al. (2008a, b, 2016), and Pavličević et al. (2014) mentioned *C. elongata* for the upper streams of the River Fojnica and Misoča, tributaries of the River Bosna (Danube catchment), while *C. elongatoides* was recorded for the River Tinja, a tributary of the River Sava (Škrijelj et al. 2008b). However, it needs to be emphasized that with the exception of Golub et al. (2016), these investigations were not exclusively loach-oriented, nor supported by the analysis of specimens caught.

From the presented overview, it is evident that, although loaches have been known from Bosnia and Herzegovina for a long time, their species identity and geographical distribution is still incompletely

resolved. However, as the taxonomy of fishes in Bosnia and Herzegovina has not been carefully studied yet, the distribution ranges of many taxa are not defined and evidence of the presence or absence of some of them needs to be clarified through further detailed study.

#### *Status and distribution of loach species in Bosnia and Herzegovina*

The distribution and the conservation status of each species were evaluated (Table 1) and are briefly discussed below.

##### Family Cobitidae

##### *Cobitis elongata* Heckel & Kner, 1858

Common name: Balkan spined loach; local name: veliki vijun, vijunica

*Cobitis elongata* is an autochthonous species occurring in many freshwaters throughout the Danube basin of South-Eastern Europe (including Slovenia, Croatia, Serbia, Montenegro, Macedonia, Bulgaria and Romania; Bănărescu 1964, 1990, Povž & Šumer 2003, Marić & Pavlović 2005, Mrakovčić et al. 2006, Kottelat & Freyhof 2007). However, data concerning both the status and the distribution of this species from Bosnia and Herzegovina are still scarce, mostly due its mistaken identification as *C. taenia* (Vuković 1963, 1977).

The first data about this species in Bosnia and Herzegovina came from Brusina (1892) and Medić (1901) who mention the River Una and their tributaries as the extent of the distribution area. Later, Kottelat & Freyhof (2007) and Sofradžija (2009), simply repeated the same data. Additional data may be found in Bănărescu & Nalbant (1957, 1998), Deliće et al. (2003a), Buj et al. (2008), Mrakovčić et al. (2000, 2006, 2008), Dumbović et al. (2009) but only for the Croatian section of the River Sava and Una. Škrijelj et al. (2008a, 2016), and Pavličević et al. (2014) mention *C. elongata* for the upper streams of the River Misoča and Fojnica, tributaries of the River Bosna (Danube catchment, Fig. 2). This species occurs on moderate to fast-flowing stretches of shallow rivers and is more frequent in the upper parts of the rivers and streams. It also occurs on sandy banks and shores, sometimes on rocky substrates with submerged vegetation, however it is not known to enter small streams, large rivers or river stretches without currents (Mrakovčić et al. 2008). Recent phylogenetic investigations proved this species to be more closely related to the species from the Adriatic basin, than to the remaining species distributed in Central Europe (Perdices & Doadrio 2001, Perdices et al. 2008, Buj et al. 2008, 2014),

even though *C. elongata* does not possess Canestrini scale, unlike majority of the Adriatic species. Its phylogeography and evolutionary history still have to be uncovered. However, based on previous investigations it separated from the remaining species in Miocene (Buj et al. 2015a).

Habitat destruction, extraction of water for irrigation as well as industrial, agricultural and untreated sewage pollution are recognised as the main threats (Mrakovčić et al. 2008). It is categorized as Least Concern (LC) species in the IUCN Red List (Freyhof & Kottelat 2008a) and the Red List of fauna of the Federation of Bosnia and Herzegovina (Škrijelj et al. 2013).



Fig. 2. Distribution of *Cobitis elongata* in Bosnia and Herzegovina based on literature data. Danube (or Black Sea) catchment (unshaded), Adriatic Sea catchment (shaded).

### *Cobitis elongatoides* Bacescu & Mayer, 1969

Common name: Danube spined loach; local name: vijun, badelj

Despite the wide distribution of this species (Kottelat & Freyhof 2007), limited historical data are available for Bosnia and Herzegovina. Like previous species, *C. elongatoides* was in the older literature most probably mentioned under the name of *C. taenia* (Vuković 1963, 1977, Vuković & Ivanović 1971, Sofradžija & Berberović 1978, Kosorić 1981, Kosorić et al. 1983, Korjenić 2004) for the River Sava and its tributaries. For the Croatian part of the River Sava it was mentioned in Mrakovčić et al. (2000) and Schneider et al. (2000a) under the name of *C. taenia danubialis*, while in Delić et al. (2003a) and Mrakovčić et al. (2008) it was identified as *C. elongatoides*. As *C. elongatoides* it was reported in Dumbović et al. (2009) for the Croatian section of the River Una.

There are some new data about ecological and biosystematic characteristics in Adrović et al. (2008).

Memić & Adrović (2010) reported this species in the upper part of the River Spreča (Sava catchment), while Škrijelj et al. (2008b) mentioned it for the River Tinja, a tributary of the River Sava (Fig. 3). Golub et al. (2016) presented information about morphological, morphometric and meristic characteristics of this species from the River Suturlija, which colonises different habitat types, from minor stagnant waters to major rivers (Mrakovčić et al. 2008).

Phylogeography of this species has been investigated in Croatia and Central Europe, although Bosnian samples were not included. Already the first comprehensive phylogeographic investigation of the genus *Cobitis* in Europe (Perdices & Doadrio 2001) had included *C. elongatoides* inside the so called *Cobitis s. str.* lineage. Closer relatedness between *C. elongatoides* and some central and eastern European species (*C. taenia*, *Cobitis tanaitica* Bacescu & Mayer, 1969, *Cobitis stephanidisi* Economidis, 1992, etc.) was corroborated by investigation of Bohlen et al. (2006). Existence of both “pure” *C. elongatoides* populations, as well as populations comprising diploid *C. elongatoides* males and females, together with triploid hybridogenetic females has been determined in the Danube basin of Croatia (Buj et al. 2008). Thereafter, presence of hybrid individuals in Bosnia and Herzegovina is also likely. Moreover, Sofradžija & Berberović (1978) reported triploid females (with 75 chromosomes in somatic cells) from middle Danube system in Bosnia and Herzegovina.

The main threats are habitat alteration, untreated sewage, pollution, the extraction of gravel from the river, water flow and level regulations, and the introduction of non-native species. It is included as the LC species in the IUCN Red List (Freyhof




Fig. 3. Distribution of *Cobitis elongatoides* in Bosnia and Herzegovina based on literature data. Danube (or Black Sea) catchment (unshaded), Adriatic Sea catchment (shaded).



& Kottelat 2008b) and the Red List of fauna of the Federation of Bosnia and Herzegovina (Škrijelj et al. 2013), respectively.

*Cobitis herzegoviniensis* Buj & Šanda, 2014

Common name: Herzegovinian spined loach, local name: hercegovački vijun

The recently described species *C. herzegoviniensis* was first recorded by Šanda et al. (2008a) in the tributary of the River Neretva (Lištica) and surrounding karstic field (Mostarsko Blato) from where spined loaches have never previously been reported (Kosorić & Vuković 1966, Aganović 1969, Kosorić 1974, 1977, 1978, Kosorić et al. 1989). Šanda et al. (2008a, b) already stated that *Cobitis* diversity in the Neretva basin is probably underestimated and further detailed taxonomic research is needed to resolve this issue, suggesting the existence of possible new species. Furthermore, a recent taxonomic analysis has revealed that the population distributed in the Mostarsko Blato karstic field form a new species (Buj et al. 2014). This species lives only in the Mostarsko Blato karstic field (Buj et al. 2014) (Fig. 4). Phylogenetically, it represents a sister species to *C. illyrica*, and belongs to the so called Adriatic phylogenetic group (Buj et al. 2014).

Life history traits are not sufficiently known so an assessment of conservation status and extinction risk has not yet been conducted (Buj et al. 2015a). Nevertheless, low genetic diversity and effective population size of this species are alarming in conservational sense (Buj et al. 2015a).

*Cobitis illyrica* Freyhof & Stelbrink, 2007

Common name: Illyric spined loach, local name: ilirski vijun

This is another recently described species, with a fragmentary distribution. *Cobitis illyrica* is an endemic species of Croatia and Bosnia and Herzegovina. It was first recorded from Lake Prološko Blato, in Imotsko polje in neighbouring Croatia (Freyhof & Stelbrink 2007). Šanda et al. (2008b) noted the presence of a possibly undescribed species in Lake Krenica, Bekijsko polje. Buj et al. (2014) confirmed their findings, which is the first record of *C. illyrica* in Bosnia and Herzegovina. This species inhabits Lake Prološko Blato, Baćinska lakes and the River Matica in neighbouring Croatia, as well as Krenica in Bekijsko polje in Bosnia and Herzegovina (Fig. 4). Based on the results of Buj et al. (2015a), three populations of *C. illyrica* are genetically distinct and gene flow among them is very restricted. Out of all

*C. illyrica* populations, the one from Lake Krenica contains the highest genetic diversity, but all three should be considered evolutionary significant units for this species (Buj et al. 2015a).

Assesment of the extinction risk for the Bosnia and Herzegovina has not yet been made (Škrijelj et al. 2013), but the global estimation for *C. illyrica* is Critically Endangered (CR), (Freyhof & Kottelat 2008c).



Fig. 4. Distribution of *Cobitis illyrica*, *C. narentana* and *C. herzegoviniensis* in Bosnia and Herzegovina based on literature data. Black rectangle indicate places where *C. illyrica* was recorded, black stars indicate *C. narentana* localities, while black dot shows record of *C. herzegoviniensis*. Danube (or Black Sea) catchment (unshaded), Adriatic Sea catchment (shaded).

*Cobitis narentana* Karaman 1928

Common name: Neretva spined loach; local name: neretvanski vijun

This is an endemic species distributed within the lower parts of the River Neretva, its tributaries and channels and Lake Modro Oko in Croatia (Schneider et al. 2000a, Zanella et al. 2003, Buj et al. 2014, 2015a), as well as in the River Trebišnjica and Hutovo Blato wetland in Bosnia and Herzegovina (Tutman et al. 2006, 2012a, Buj et al. 2014, 2015a, b). It was first described as a subspecies (*C. taenia narentana* Karaman 1928). Later data include only general notes from the Neretva River drainage system (Vuković 1963, 1977). The systematic status has been changed, since Schneider et al. (2000a) redescribed it as a separate species *C. narentana*. Sofradžija (2009) considered the lower Neretva River from town Čapljina as their distribution area in Bosnia and Herzegovina, while Tutman et al. (2006, 2012a), Šanda et al. (2008a, b), and Buj et al. (2015a, b) extended it to the Hutovo Blato wetland area, Šanda et al. (2008a, b) in the River Bregava and Šanda et al. (2008a, b) and Buj et al. (2015a, b) in

the River Trebišnjica, all in Bosnia and Herzegovina (Fig. 4). It prefers sandy or silty substrates or dense vegetation in slow-flowing water. Phylogenetically, *C. narentana* is a sister species to *C. dalmatina* (distributed in the River Cetina in Croatia), and also belongs to the Adriatic lineage (Perdices et al. 2008, Buj et al. 2014). Its genetic diversity is very high, due to long, unconstrained evolutionary history as well as distribution on several localities (Buj et al. 2015a). Contrary to situation with *C. illyrica*, populations of *C. narentana* are genetically similar (Buj et al. 2015a). This species is susceptible to anorganic pollution, watercourse regulation and sand and gravel extraction, as well as introduction of alien species such as pike-perch and rainbow trout (Tutman et al. 2006, 2012a, b). A proposed dam construction will decrease water flow, increase salinification and the threat of drought (Zanella et al. 2009). Based on the available data (IUCN 2016) it is categorized as a Vulnerable species (VU), which is also its status in the Red List of fauna of the Federation of Bosnia and Herzegovina (Škrijelj et al. 2013).

#### *Misgurnus fossilis* (Linnaeus, 1758)

Common name: weather loach; local name: čikov, piškor

This species is distributed throughout the whole of Europe from northern France to western Russia (including the Danube and Volga River basins) except for Scandinavia, the Mediterranean and the British Isles (Kottelat & Freyhof 2007). However, historical data on the distribution of this species in Bosnia and Herzegovina are very scarce (Vuković 1963, 1977, Bogut et al. 2006), so the actual distribution is not certain. According to the available data, *M. fossilis* occurs in the Sava River basin, namely the River Sava with its tributaries, river branches and oxbows in the lowland floodplain regions of northern Bosnia and Herzegovina (Fig. 5), where it prefers calm and stagnant waters (Sofradžija 2009). The factors threatening this species include loss of habitats due to change in hydrological regime, river regulation and pollution of backwaters, which are likely to have the most negative effect on this species in the region (Mrakovčić et al. 2000, 2008). It is categorised as the LC species in the IUCN Red List (Freyhof 2011a) and the Red List of fauna of the Federation of Bosnia and Herzegovina (Škrijelj et al. 2013), respectively. Bohlen et al. (2007) investigated mitochondrial DNA phylogeography of *M. fossilis* in Europe and found very low mitochondrial divergence with just eight closely related haplotypes. The most

southern haplotype, that is closest to the Bosnia and Herzegovina, was located in the Poganovo polje (Croatia, Park of Nature Lonjsko polje). This sample revealed specific population different from all other samples and localities. Authors also considered that southern-eastern part of the range was a refuge area for the species during the glaciation events in the Pleistocene. Another ecologically similar lowland species *Umbra krameri* Walbaum, 1792, was shown to be also genetically different in the Sava River basin (Marić et al. 2017). Phylogeography of this species has been investigated in Danube and Dniester drainages, and it has been found that population from Sava River basin belongs to specific phyletic lineage (haplogroup) and could be considered as potential Evolutionary Significant Units (ESU). Therefore, it is possible that southern part (Bosnia and Herzegovina) of the *M. fossilis* population in Sava River basin could be slightly genetically different from all other populations, and could represent an evolutionary significant unit. This need further investigation.



Fig. 5. Distribution of *Misgurnus fossilis* in Bosnia and Herzegovina based on literature data. Danube (or Black Sea) catchment (unshaded), Adriatic Sea catchment (shaded).

#### *Sabanejewia balcanica* (Karaman, 1922)

Common name: Balkan golden loach; local name: zlatni vijun

The Balkan golden loach is widely distributed in the Danube basin (Romania, Bulgaria, Croatia, Bosnia and Herzegovina, Serbia, Slovenia, Slovakia, Czech Republic, Ukraine, Russia) and it has been reported from the Aegean Sea catchment in Maritza and from Gallikos to Pinios drainages (Freyhof & Kottelat 2008d). In earlier studies, *S. balcanica* in Bosnia and Herzegovina was considered as a subspecies of *C. aurata*. Karaman (1963) conducted the first assessment of *S. balcanica* in Bosnia and Herzegovina

describing a new subspecies as *C. aurata bosniaca* in the River Vrbas and its tributaries Suturlija and Široka (Sava River catchment). Vuković (1963, 1977) and Sofradžija (2009) identified two subspecies; *C. aurata balcanica* Karaman, 1922 in waters of the Danube catchment and *C. aurata bosniaca* Karaman 1963 only in the River Vrbas and its tributaries Suturlija and Široka (Sava River catchment). Later authors (Mrakovčić et al. 2000, 2006, 2008, Bogut et al. 2006, Sofradžija 2009, Drešković et al. 2011) simply repeated previous reports without new data on the precise distribution.

However, it is now recognized that *S. aurata* is an Asian species and does not occur in European waters (Kottelat 1997). According to Kottelat & Freyhof (2007) single *Sabanejewia* species (*S. balcanica*) is present in Bosnia and Herzegovina waters. For the Croatian section of the River Una it was reported by Dumbović et al. (2009). Muhamedagić et al. (2013) reported *S. balcanica* in the upper parts of the River Krivaja in central Bosnia and Herzegovina. Recently, Golub et al. (2016) presented morphological, morphometric and meristic characteristics of this species from the River Suturlija. According to the available data, its known distribution was established in the middle and upper parts of several rivers and streams in eastern, northern and central Bosnia and Herzegovina, in the tributaries of the Sava River basin where it overlapped with the *C. elongatoides* distributed in lower tributaries (Drina, Bosna and Vrbas) (Fig. 6). It inhabits streams with a moderate current and sandy or gravely substrates. Samples from several rivers in Bosnia and Herzegovina (Una, Vrbas and Bosna) were included in the investigation of mitochondrial DNA phylogeography (Marešová

et al. 2011). They turned out to belong to so called Danubian-Balkanian complex, which is genetically heterogenous (Perdices et al. 2003) and taxonomically still not resolved.

*Sabanejewia balcanica* is categorized as LC in the IUCN Red List (Freyhof & Kottelat 2008b). Due to the lack of knowledge on its distribution range, frequency and deleterious factors, the species is categorized as Data Deficient (DD) in the Red List of fauna of the Federation of Bosnia and Herzegovina (Škrijelj et al. 2013).

#### Family Nemacheilidae

#### *Barbatula barbatula* (Linnaeus, 1758)

Common name: stone loach; local name: brkica, tivuška  
The stone loach has a wide natural distribution in Eurasia, from the Pyrenees in the west to the Ural and Emba basins in the east, but it does not occur on the Italian and Iberian Peninsulas (Kottelat & Freyhof 2007). The species was first mentioned under the name *C. barbatula* by Zaplata & Taler (1932) in the main course of the River Bosna. Furthermore, Vuković (1963, 1977) and Sofradžija (2009) indicated the species as common, but not in greater numbers in the small rivers of the Sava River basin under the *Nemachilus barbatulus* and *Nemacheilus barbatulus*, respectively. Finally, under the name of *B. barbatula* Bogut et al. (2006) mentioned it for the waters of the Danube system without precise locations. There is a very interesting record of *B. barbatula* in the upper Neretva River, Adriatic catchment (Vegara et al. 2009) (Fig. 7). No other scientific publication has previously reported *B. barbatula* in the Neretva River catchment, nor in any other river of the Adriatic drainage system in Bosnia and Herzegovina (Vuković 1977, Sofradžija 2009). This paper describes the first record of *B. barbatula* in the River Neretva and consequently also the first record for the Adriatic drainage system in Bosnia and Herzegovina as a whole. The exact taxonomic status of this population cannot be resolved at the moment, since presence of *Barbatula zetensis* (Šorić, 2000) or *Barbatula sturanyi* (Steindachner, 1892), that inhabit the nearby Ohrid-Drin-Skadar basin cannot be ruled out. It is also not clear whether this population is an introduced one (source unknown) or if it is native, but overlooked in previous investigations. Genetic structure of *B. barbatula* is very complex. It comprises several highly distinct clades and its monophyly was not confirmed in previous investigations (Šedivá et al. 2008). Although samples from Bosnia and Herzegovina were not included in molecular genetic investigation




Fig. 6. Distribution of *Sabanejewia balcanica* in Bosnia and Herzegovina based on literature data. Danube (or Black Sea) catchment (unshaded), Adriatic Sea catchment (shaded).

of this species (Šedivá et al. 2008), it is possible that different lineages, or even species, are present in different basins.

As no major threats are recognised, it is included as LC species in the IUCN Red List (Freyhof 2011b) and the Red List of fauna of the Federation of Bosnia and Herzegovina (Škrijelj et al. 2013), respectively.



**Fig. 7.** Distribution of *Barbatula barbatula* in Bosnia and Herzegovina based on literature data. Danube (or Black Sea) catchment (unshaded), Adriatic Sea catchment (shaded).

### Vulnerability

Loaches are considered threatened throughout Europe, owing to human induced alteration of freshwater ecosystems (Lelek 1987, Kotusz 1996). Even though taxonomic status and phylogenetic relationships of the Adriatic spined loaches have been investigated (Buj et al. 2014, 2015a), data on their, and other loach species vulnerability in Bosnia and Herzegovina are still lacking. The main threats affecting loach species in Europe are related to water pollution and habitat loss by inappropriate modifications of riverbeds (IUCN 2016). Their protracted residence in silt habitats and vegetation required by juveniles, makes them highly susceptible to habitat modification (IUCN 2016). Although there are no specialised investigations dedicated to this topic available for Bosnia and Herzegovina, it seems that the biggest threats are the regulation of the riverbeds, like in the River Trebišnjica (Buj et al. 2015a), and river remodeling and accumulation, thus reducing the flow of organic particles downstream. Loaches are known to be sensitive to pollution, as well as sediment and gravel extraction (Nedić et al. 2014). In Bosnia and Herzegovina none of them has any economic value and they are not subjected to economic exploitation. However, in some areas they can be used as live bait for salmon fishing.

### Discussion

The region of the Balkans is often considered as an ichthyologic “hot spot”, with a great number of species and a high portion of endemism for the loach species living in freshwaters in a relatively small area (Buj et al. 2015a). Bosnia and Herzegovina’s zoogeographic location and climatic conditions provide a rich fish fauna in which several loach species have been identified in different types of water bodies with a soft substratum. Unfortunately, this rich ichthyofauna has been understudied and is not receiving adequate protection.

#### *Level of investigation of loaches in Bosnia and Herzegovina*

According to the available ichthyological literature, a total of nine species of loaches were reported to inhabit the freshwaters of Bosnia and Herzegovina. Although their presence has been reported for a long time (Brusina 1892, Medić 1901, Zaplata & Taler 1932), their species identity and geographical distribution is still poorly known. The incorrect identification, coupled with improvements in taxonomy, has resulted in confusing conclusions in several reports, making them unreliable. Considering data about biosystematic position of these species in the waters of Bosnia and Herzegovina, Golub et al. (2016) suggested that it could be at least partially due to an inconsistent approach to the identification of these species.

The list of eight species of Cobitidae and Nemacheilidae in Bosnia and Herzegovina established in this paper, exceeds the estimation of the most recent check-list of Sofradžija (2009). Such a pronounced difference in a number of listed species reflects shortcomings in the knowledge on the species composition and geographic distribution and confirms the need for more detailed analysis in this field. In the absence of precise scientific research, although rare, data from various ichthyological monitoring surveys present an important source of information. The obvious deficiency of data in published literature is probably associated with several problems: lack of adequately targeted loach-oriented research, as well as sampling techniques; small size and lack of commercial value of loaches, which is the reason that they can easily be overlooked (Mustafić et al. 2003, Šanda et al. 2008a, b), like other fishes with no economic value (Tutman et al. 2012b, 2013). Actually, the only targeted investigation into the morphological, morphometric and meristic characteristics of loach species were those presented by Golub et al. (2016).

### *Presence and distribution of loaches in Bosnia and Herzegovina*

Scarce and sometimes unreliable data in the literature have led to problematic assessments of the presence of different loach species in the freshwaters of Bosnia and Herzegovina. Particularly problematic is the status of *C. taenia*. Despite its listing as a member of the Bosnia and Herzegovina ichthyofauna since Zaplata & Taler (1932) and Vuković (1977), it is now evident that its distribution range extends far beyond this area (Šlechtová et al. 2000, Bohlen et al. 2002) and it should not be considered as a part of the ichthyofauna of Bosnia and Herzegovina. However, it is not clear which species reports of *C. taenia* (Vuković 1963, 1977, Kosorić & Mikavica 1981 and others) should be assigned to.

Assessing distribution ranges from older reports is also problematic, since only general notes are sometimes given instead of the exact localities. Two catchments in Bosnia and Herzegovina accommodate different loach species. The Black Sea catchment is inhabited by five species with wider distribution ranges. The distribution area of *C. elongata* in Bosnia and Herzegovina includes the lower reaches of the rivers Una, Vrbas, Ukrina and Bosna, tributaries of the River Sava (Danube drainage), distribution range of *C. elongatoides* includes middle and lower reaches of tributaries of the River Sava (Una, Sana, Vrbas, Ukrina, Bosna and Drina), and River Sava itself. *Misgurnus fossilis* is distributed in the River Sava, *B. barbatula* in upper reaches of the rivers Sana, Vrbas, Bosna, Drina, while the area of distribution of *S. balcanica* includes the middle reaches of the same rivers in central Bosnia and Herzegovina.

On the other hand, the Adriatic catchment in Bosnia and Herzegovina is inhabited by three loach species, all of which are endemic: *C. narentana* is endemic to the Neretva River basin, *C. illyrica* is distributed fragmentarily in Croatia and Bosnia and Herzegovina and *C. herzegoviniensis* is found only in the Mostarsko Blato karstic field (Buj et al. 2014, 2015a). There is a single report of the occurrence of *B. barbatula* in the upper River Neretva (Vegara et al. 2009), which would mean that *B. barbatula* is the only species found in both catchments. However, from the report it is not possible to make definitive conclusions about its status and distribution. Therefore, further detailed phylogenetic and taxonomic research is needed to resolve this issue.

The most common and widespread species of loaches in Bosnia and Herzegovina, as well as throughout central and northern Europe (Bohlen & Ráb 2001) is

*C. elongatoides*. It was found on all major drainage systems inside the Danube catchment, in relatively numerous populations. The population data of this species in several rivers of the continental part of Croatia display the same pattern (Delić et al. 2003a, b, 2009). On the other hand, *B. barbatula* is the rarest with regards to its occurrence in both the Danube and Adriatic catchment.

The reported occurrence and distribution of loach species in Bosnia and Herzegovina follows the pattern observed throughout Europe, with only rare syntopic occurrences of different species (Bohlen & Ráb 2001). It seems that smaller water systems usually contain only a single species, while in the larger ones sympatric species are often separated in different river sections. For example, in the river systems of the Vrbas and Bosna, *S. balcanica* inhabits the upper part of the basins, *C. elongatoides* is found in the middle, while *C. elongata* in the lower stretches (Sofradžija 2009). The same pattern is observed in the river systems of the Elbe and Odra (Boroń & Kotusz 1999, Ráb et al. 2000, Bohlen et al. 2002). The opposite pattern is observed in the Adriatic catchment, with only a single species of loach found in a particular locality or fish community (Šanda et al. 2008a, b, Buj et al. 2008, 2014, 2015a, b).

### *Extinction risk and conservation of loaches in Bosnia and Herzegovina*

Unfortunately, regional assessment of the extinction risk of loaches in Bosnia and Herzegovina has not yet been conducted. Up to the present day, the Republic of Bosnia and Herzegovina has no Red Book of Freshwater Fishes nor a formal strategy for their conservation. There is only the Red List of fauna of the Federation of Bosnia and Herzegovina as part of the Republic of Bosnia and Herzegovina (Škrijelj et al. 2013). In this list six loach species were included; *C. narentana* as VU, *S. balcanica* as DD, and the *C. elongata*, *C. elongatoides*, *M. fossilis* and *B. barbatula* as LC species. Global assessment following IUCN criteria, however, has been conducted for the majority of loaches. Two endemic loach species present in Bosnia and Herzegovina are considered endangered according to the IUCN categories and significant anthropologic impact is posing a threat to their survival. On the other hand, both of them are endemic, distributed in a single river, river basin or only on several localities in a very small area (Buj et al. 2014, 2015a, b), implying the necessity of immediate effective protection. *C. narentana* is estimated as VU, and *C. illyrica* is considered CR. It is important to mention that the *C. illyrica* population

in Lake Krenica is genetically different from the Croatian populations (Buj et al. 2015a) and should be considered as a distinct conservation unit. For *C. herzegoviniensis* extinction risk estimation has not yet been conducted (IUCN 2016). The remaining species are included as the LC species in the IUCN Red List. Considering the extinction risk estimations of these species and the relatively poor understanding of their taxonomical position in Bosnia and Herzegovina, basic biological features, and balances in place with regard to habitat ecology, there is an imperative to employ ichthyological surveys of these species, as a basis for taking species protection security measures.

The need for detailed ichthyologic research in order to provide precise data about Bosnia and Herzegovina loach fish fauna is still required. A main limitation in active protection of a species is incomplete knowledge about its geographical distribution combined with uncertain taxonomic status. This highlights the importance of establishing conservation measures, considering the high faunistic and scientific value of this family.

## Acknowledgements

We are thankful to Jason Kirby from Liverpool John Moores University, U.K., for improving the English in the manuscript.

## Literature

- Adrović A., Skenderović I., Salihović J. & Stjepić Š. 2012: Biodiversity of fish fauna of the Gostelja River and reservoir Modrac. Structure and dynamics of ecosystems Dinarides – status, possibilities and prospects. *Academy of Sciences and Arts of Bosnia and Herzegovina, Department of Natural Sciences and Mathematics, Proceedings, Special Editions CXLIX 23*: 277–290.
- Adrović A., Škrijelj R. & Đug S. 2008: Ecological and biosystemic characteristics of *Cobitis elongatoides* Bacescu & Maier, 1969 from the rivers Brka, Gostelja and Tinja in the basin Sava River. *XIV Proceedings Biotechnological consultancy, Čačak*: 359–366. (in Bosnian)
- Aganović M. 1969: Composition of fish populations of the River Jaruga. *Ichthyologia 1*: 3–10. (in Serbo-Croatian/Croato-Serbian with English summary)
- Bacescu M. 1962: Contribution a la systematique du genre *Cobitis*. Description d'une espèce nouvelle, *Cobitis calderoni*, provenant de l'Espagne. *Rev. Biol. (Bucarest) 6*: 435–438.
- Bănărescu P. 1964: Fauna Republicii Populare Române: Pisces-Osteichthyes, vol. 13. *Academia Republicii Populare Române*.
- Bănărescu P. 1990: Zoogeography of fresh waters, vol. 1. General distribution and dispersal of freshwater animals. *Aula-Verlag, Wiesbaden, Germany*.
- Bănărescu P. & Nalbant T.T. 1957: Eidionomische und taxionomische Untersuchungen an *Cobitis elongata* (Pisces, Cobitidae). *Senckenb. Biol. 38*: 283–294.
- Bănărescu P. & Nalbant T.T. 1998: *Cobitis elongata* Heckel and Kner, 1858 (Pisces, Ostariophysi, Cobitidae), distribution, relationships, geographical variation and protection status. *Rev. Roum. Biol. Anim. 43*: 57–65.
- Bogut I., Novoselić D. & Pavličević I. 2006: Biology of fishes. *University of J. J. Strossmayer, University of Mostar, Faculty of Agriculture, Osijek*. (in Croatian)
- Bohlen J., Perdices A., Doadrio I. & Economidis P.S. 2006: Vicariance, colonisation, and fast local speciation in Asia Minor and the Balkans as revealed from the phylogeny of spined loaches (Osteichthyes; Cobitidae). *Mol. Phylogenet. Evol. 39*: 552–561.
- Bohlen J. & Ráb P. 2001: Species and hybrid richness in spined loaches of the genus *Cobitis* (Teleostei: Cobitidae), with a checklist of European forms and suggestions for conservation. *J. Fish Biol. 59 (Suppl. A)*: 75–89.
- Bohlen J., Ráb P., Šlechtová V. et al. 2002: Hybridogeneous biotypes in spined loaches (genus *Cobitis*) in Germany with implications for the conservation of such fish complexes. In: Collares-Pereira M.J., Cowx I. & Coelho M. (eds.), *Conservation of freshwater fishes: options for the future* (chapter 28). *Fishing News Books, Blackwell Science, Oxford*: 311–321.
- Bohlen J., Šlechtová V., Doadrio I. & Ráb P. 2007: Low mitochondrial divergence indicates a rapid expansion across Europe in the weather loach, *Misgurnus fossilis* (L.). *J. Fish Biol. 71*: 186–194.
- Bohlen J., Šlechtová V., Šanda R. et al. 2003: *Cobitis ohridana* and *Barbatula zetensis* in the River Morača basin, Montenegro: distribution, habitat, population structure and conservation needs. *Folia Biol. (Kraków) 51 (Suppl.)*: 147–153.
- Boroń A. & Kotusz J. 1999: A preliminary report and karyotype of a new *Cobitis* species in the ichthyofauna of Poland. *Cytobios 98*: 59–64.
- Brusina S. 1892: Gleanings for the Croatian ichthyology and fisheries. *Glas. Hrv. Narav. Društva 1–6*: 221–288. (in Croatian)
- Buj I., Čaleta M., Marčić Z. et al. 2015a: Different histories, different destinies – impact of evolutionary history and population genetic structure on extinction risk of the Adriatic spined loaches (genus *Cobitis*; Cypriniformes, Actinopterygii). *PLOS ONE 10*: e0131580.
- Buj I., Podnar M., Mrakovčić M. et al. 2008: Genetic diversity and phylogenetic relationships of spined loaches (genus *Cobitis*) in Croatia based on mtDNA and allozyme analyses. *Folia Zool. 57*: 71–82.
- Buj I., Šanda R., Marčić Z. et al. 2014: Combining morphology and genetics in resolving taxonomy – a systematic revision of spined loaches (genus *Cobitis*; Cypriniformes, Actinopterygii) in the Adriatic watershed. *PLOS ONE 9*: e99833.
- Buj I., Šanda R., Marčić Z. et al. 2015b: Sexual dimorphism of five *Cobitis* species (Cypriniformes, Actinopterygii) in the Adriatic watershed. *Folia Zool. 64*: 97–103.
- Čaleta M., Buj I., Mrakovčić M. et al. 2015: Croatian endemic fishes. *Environmental Protection Agency, Zagreb, Croatia*. (in Croatian)

- Delić A., Bučar M., Jugović D. et al. 2009: New data on the distribution *Cobitis elongatoides* Bacescu & Maier, 1969 (Cobitidae) in Central Croatia. *Nat. Croat.* 18: 255–262.
- Delić A., Bučar M., Kučinić M. & Mrakovčić M. 2003a: New data about the distribution of *Sabanejewia balcanica* (Karaman, 1922) (Cobitidae) in Croatia. *Folia Biol. (Kraków)* 51 (Suppl.): 39–42.
- Delić A., Kučinić M., Bučar A. et al. 2003b: Morphometric and meristic characteristics of the goldside loach *Sabanejewia balcanica* Kar. in central Croatia. *Folia Biol. (Kraków)* 51 (Suppl.): 33–38.
- Drešković N., Đug S., Stupar V. et al. 2011: NATURA 2000 in Bosnia and Herzegovina. *Center for Environment and Sustainable Development, Sarajevo. (in Bosnian)*
- Dulčić J., Tutman P., Prusina I. et al. 2009: Length-weight relationships for six endemic freshwater fishes from Hutovo blato wetland (Bosnia and Herzegovina). *J. Appl. Ichthyol.* 25: 499–500.
- Dumbović V., Posavec Vukelić V., Duplić A. et al. 2009: Action plan for protection of biodiversity of the river Una and surrounding area. *Sisačko-Moslavačka region, Sisak. (in Croatian)*
- Erős T., Sallai Z. & Korusz J. 2003: Distribution and conservation status of loaches in Hungary. *Folia Biol. (Kraków)* 51 (Suppl.): 17–19.
- Freyhof J. 2011a: *Misgurnus fossilis*. The IUCN Red List of Threatened Species 2011. Downloaded on 02 June 2017. <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T40698A10351495.en>
- Freyhof J. 2011b: *Barbatula barbatula*. The IUCN Red List of Threatened Species 2011. Downloaded on 02 June 2017. <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T14494A4439010.en>
- Freyhof J. & Kottelat M. 2008a: *Cobitis elongata*. The IUCN Red List of Threatened Species 2008. Downloaded on 02 June 2017. <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T5031A11108572.en>
- Freyhof J. & Kottelat M. 2008b: *Cobitis elongatoides*. The IUCN Red List of Threatened Species 2008. Downloaded on 02 June 2017. <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T135597A4156685.en>
- Freyhof J. & Kottelat M. 2008c: *Cobitis illyrica*. The IUCN Red List of Threatened Species 2008. Downloaded on 02 June 2017. <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T135712A4188832.en>
- Freyhof J. & Kottelat M. 2008d: *Sabanejewia balcanica*. The IUCN Red List of Threatened Species 2008. Downloaded on 02 June 2017. <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T135661A4173614.en>
- Freyhof J. & Stelbrink B. 2007: *Cobitis illyrica*, a new species of loach from Croatia (Teleostei: Cobitidae). *Ichthyol. Explor. Fresh.* 18: 269–275.
- Geiger M.F., Herder F., Monaghan M.T. et al. 2014: Spatial heterogeneity in the Mediterranean Biodiversity Hotspot affects barcoding accuracy of its freshwater fishes. *Mol. Ecol. Resour.* 14: 1210–1221.
- Glamuzina B., Pavličević J., Tutman P. et al. 2013: Fishes of Neretva. *Association CEAV and Modrozeleno – Team Protector, Mostar/Metković, Croatia. (in Croatian)*
- Golub D., Marić Ž., Šukalo G. et al. 2016: Morphological characteristics of *Cobitis elongatoides* and *Sabanejewia balcanica* (Cobitidae) from river Suturlija. *Third Symposium of biologists and ecologists of Republic of Srpska, 12-14 November 2015, Banja Luka: 139–148. (in Serbian with English abstract)*
- Hamzić A., Ugarčina N., Péteri A. et al. 2015: Analysis of the fishery and aquaculture sector in Bosnia and Herzegovina. *Food and Agriculture Organization of the United Nations, Regional Office for Europe and Central Asia.*
- IUCN 2016: The IUCN Red List of Threatened Species, version 2016-3. Downloaded on 07 December 2016. <http://www.iucnredlist.org>
- Janko K., Kotlík P., Culling M.A. & Ráb P. 2005a: Ice age cloning – comparison of quaternary evolutionary histories of sexual and clonal forms of European loaches (*Cobitis*; Teleostei) using the analysis of mitochondrial DNA variation. *Mol. Ecol.* 14: 2991–3004.
- Janko K., Vasiļev V.P., Ráb P. et al. 2005b: Genetic and morphological analyses of 50-chromosome spined loaches (*Cobitis*, Cobitidae, Pisces) from the Black Sea basin that are morphologically similar to *C. taenia*, with the description of a new species. *Folia Zool.* 54: 405–420.
- Jukić M. 2008: The River Sava as a resource in spatial planning. *Geoadria* 13: 81–96.
- Karaman M. 1963: Eine neue Unterart des Steinpetzgers, *Cobitis aurata bosniaca* n. ssp. (Cobitidae, Pisces) aus Jugoslawien. *Int. Rev. Ges. Hydrobiol.* 48: 629–632.
- Karaman S. 1928: Contribution to the Yugoslavian ichthyology. *Glasnik Skopskog naučnog društva, Skoplje* 6: 147–176. (in Macedonian and German)
- Korjenić E. 2004: Variation of the basic body measure of some fish species from the Fojnica River. *Veterinaria* 53: 25–35. (in Bosnian)
- Kosorić Đ. 1974: Fishes of the Rama River. *Ichthyologia* 6: 69–78. (in Serbo-Croatian/Croato-Serbian with English summary)
- Kosorić Đ. 1977: Fish populations of the middle part of the Neretva River, with projection of spreading possibilities after water reservoir construction. *Ichthyologia* 9: 121–129. (in Serbo-Croatian/Croato-Serbian with English summary)
- Kosorić Đ. 1978: The composition of the Hutovo blato fish population. *Godišnjak BIUS-a* 31: 69–81. (in Serbo-Croatian/Croato-Serbian with English summary)
- Kosorić Đ. 1981: Fish population of Krivaja River. *Godišnjak BIUS-a* 34: 57–73. (in Serbo-Croatian/Croato-Serbian)
- Kosorić Đ. & Mikavica D. 1981: Fish populations of Krivaja River. *Annual BIUS University in Sarajevo* 34: 57–71. (in Serbo-Croatian/Croato-Serbian with English summary)
- Kosorić D. & Vuković T. 1966: Fishes of the River Buna. *Glasnik Zemaljskog muzeja Bosne i Hercegovine u Sarajevu, Prirodne Nauke* 5: 179–190. (in Serbo-Croatian/Croato-Serbian with German summary)
- Kosorić D., Vuković T., Guzina N. et al. 1989: Ichthyofauna of River Neretva and its changes after the building of hydroelectricity power stations. *Zbornik radova – Savjetovanje o ribarstvu na hidroakumulacijama, Mostar: 133–138. (in Serbo-Croatian/Croato-Serbian)*

- Kosorić D., Vuković T., Kapetanović N. et al. 1983: Fish composition of the Neretva River in Bosnia and Herzegovina. *Godišnjak BIUS-a* 36: 117–128. (in Serbo-Croatian/Croato-Serbian)
- Kottelat M. 1997: European freshwater fishes. *Biologia (Bratislava)* 52 (Suppl. 5): 1–271.
- Kottelat M. & Freyhof J. 2007: Handbook of European freshwater fishes. Kottelat, Cornol, Switzerland and Freyhof, Berlin, Germany.
- Kotusz J. 1996: Species protection of loaches (Cobitoidea, Cypriniformes) in Poland in relation to their distribution and status in other European countries. *Zool. Polon.* 41 (Suppl.): 147–155. (in Polish with English summary)
- Lelek A. 1987: Threatened fishes of Europe. The freshwater fishes of Europe, vol. 9. *Aula-Verlag, Wiesbaden, Germany.*
- Marešová E., Delić A., Kostov V. et al. 2011: Genetic diversity of *Sabanejewia balcanica* (Actinopterygii: Cobitidae) in the western Balkans and comparison with other regions. *Folia Zool.* 60: 335–342.
- Marić D. & Milošević D. 2010: First record and morphological characteristics of the Balkan golden loach *Sabanejewia balcanica* (Cobitidae) in Montenegro. *Period. Biol.* 112: 149–152.
- Marić D. & Pavlović V. 2005: First records and description of *Cobitis elongata* Heckel & Kner 1858 (Cobitidae) in Montenegro. *Nat. Montenegr.* 5: 109–115.
- Marić S., Stanković D., Wanzenböck J. et al. 2017: Phylogeography and population genetics of the European mudminnow (*Umbra krameri*) with a time-calibrated phylogeny for the family Umbridae. *Hydrobiologia* 792: 151–168.
- Medić M. 1901: Second round of ichthyological notes. *Rad Jugosl. Akad. Znan. i Umj.* 30: 139–191. (in Croatian)
- Memić K. & Adrović A. 2010: Biodiversity of ichthyofauna of the Spreča river as a part of natural heritage of Tuzla area. Condition of the cultural and natural heritage in the Balkan region. *Central Institute for Conservation in Belgrade, Institute Goša d.o.o., Ministry of Culture of Republic of Serbia:* 215–226.
- Mrakovčić M., Brigić A., Buj I. et al. 2006: Red Book of Croatian freshwater fishes. *Ministry of Culture, DZZP, Zagreb, Republic of Croatia.* (in Croatian with English summary)
- Mrakovčić M., Duplić A., Mustafić P. & Marčić Z. 2008: Conservation status of the genus *Cobitis* and related genera in Croatia. *Folia Zool.* 57: 35–41.
- Mrakovčić M., Mišetić S. & Povž M. 1995: Status of freshwater fish in Croatian Adriatic river systems. *Biol. Conserv.* 72: 179–185.
- Mrakovčić M., Schneider D., Mustafić P. & Kerovec M. 2000: Status of genus *Cobitis* and related species in Croatia. *Folia Zool.* 49 (Suppl. 1): 116–119.
- Muhamedagić S., Korjenić E., Hamzić A. et al. 2013: Ichthyofauna of Krivaja River catchment area. In: Poleksić V. (ed.), *Water & Fish. Conference proceedings from VI International Conference “Water & fish”, University of Belgrade, Faculty of Agriculture, Serbia:* 310–321.
- Mustafić P., Mrakovčić M., Čaleta M. et al. 2003: Loaches in long term study of the Drava River in Croatia. *Folia Biol. (Kraków)* 51 (Suppl.): 143–146.
- Nalbant T.T., Ráb P., Bohlen J. & Saitoh K. 2001: Evolutionary success of the loaches of the genus *Cobitis* (Pisces: Ostariophysi: Cobitidae). *Trav. Mus. Natl. Hist. Nat. „Grigore Antipa“* 43: 277–289.
- Nedić Z., Begović S., Dogan M. et al. 2014: Decreasing of biodiversity of fish populations from the Sava River in Bosnia and Herzegovina. *BJASR* 3: 35–40.
- Pavličević J., Petrović D., Bogut I. et al. 2014: Revision of the fishing basis for the fishing zone of Central Bosnia Canton. *Faculty of Agricultural and Food Technology, University of Mostar, Department of Fisheries, Zoology and Water Protection, Bosnia and Herzegovina, Mostar.* (in Croatian)
- Perdices A., Bohlen J. & Doadrio I. 2008: The molecular diversity of Adriatic spined loaches (Teleostei, Cobitidae). *Mol. Phylogenet. Evol.* 46: 382–390.
- Perdices A., Bohlen J., Šlechtová V. & Doadrio I. 2016: Molecular evidence for multiple origins of the European spined loaches (Teleostei, Cobitidae). *PLOS ONE* 11: e0144628.
- Perdices A. & Doadrio I. 2001: The molecular systematics and biogeography of the European cobitids based on mitochondrial DNA sequences. *Mol. Phylogenet. Evol.* 19: 468–478.
- Perdices A., Doadrio I., Economidis P.S. et al. 2003: Pleistocene effect on the European freshwater fish fauna: double origin of the cobitid genus *Sabanejewia* in the Danube basin (Osteichthyes: Cobitidae). *Mol. Phylogenet. Evol.* 26: 289–299.
- Povž M. & Šumer S. 2000: Present status and distribution of the species of the genera *Misgurnus*, *Cobitis* and *Sabanejewia* in Slovenia. *Folia Zool.* 49 (Suppl. 1): 107–112.
- Povž M. & Šumer S. 2003: Status, distribution and morphometric/meristic characteristics of *Cobitis elongata* Heckel & Kner 1858 from Slovenia. *Folia Biol. (Kraków)* 51 (Suppl. 1): 21–25.
- Ráb P., Rábová M., Bohlen J. & Lusk S. 2000: Genetic differentiation of the two hybrid diploid-polyploid complexes of loaches, genus *Cobitis* (Cobitidae), involving *C. taenia*, *C. elongatoides* and *C. spp.* in the Czech Republic. Karyotypes and cytogenetic diversity. *Folia Zool.* 49 (Suppl. 1): 55–66.
- Schneider D., Mrakovčić M., Mustafić P. & Kerovec M. 2000b: Morphological difference in some *Cobitis* populations from Croatia. *Folia Zool.* 49 (Suppl. 1): 227–234.
- Schneider D., Mustafić P., Mrakovčić M. & Mihaljević Z. 2000a: Some aspects of the biology of the Neretvan spined loach. *Folia Zool.* 49 (Suppl. 1): 159–165.
- Simonović P., Povž M., Piria M. et al. 2015: Ichthyofauna of the River Sava system. In: Milačić R., Ščančar J. & Paunović M. (eds.), *The Sava River. The Handbook of Environmental Chemistry* 31, Springer; Berlin-Heidelberg: 361–400.
- Sofradžija A. 2009: Freshwater fishes of the Bosnia and Herzegovina. *Council of the Congress of Bosnian Intellectuals, Sarajevo.* (in Bosnian)
- Sofradžija A. & Berberović Lj. 1978: Diploid-triploid sexual dimorphism in the loach (*Cobitis taenia taenia* L., Cobitidae, Pisces). *Genetika* 10: 389–397. (in Serbo-Croatian/Croato-Serbian)



- Sofradžija A. & Vuković T. 1979: Chromosomes of *Nemachilus barbatus*. *Ichthyologia* 11: 5–13. (in Serbo-Croatian/Croato-Serbian)
- Šanda R., Bogut I., Doadrio I. et al. 2008a: Distribution and taxonomic relationships of spined loaches (Cobitidae, *Cobitis*) in the River Neretva basin, Bosnia and Herzegovina. *Folia Zool.* 57: 20–25.
- Šanda R., Bogut I. & Vukić J. 2008b: New data of the ichthyofauna of lower Neretva River catchment and surrounding karstic fields. In: Bogut I. (ed.), Freshwater fish farming, status and perspectives. *Croatian Chamber of Economy, Zagreb: 119–125.* (in Croatian)
- Šanda R., Vukić J., Choleva L. et al. 2008c: Distribution of loach fishes (Cobitidae, Nemacheilidae) in Albania, with genetic analysis of populations of *Cobitis ohridana*. *Folia Zool.* 57: 42–50.
- Šedivá A., Janko K., Šlechtová V. et al. 2008: Around or across the Carpathians: colonization model of the Danube basin inferred from genetic diversification of stoneloach (*Barbatula barbatula*) populations. *Mol. Ecol.* 17: 1277–1292.
- Škrijelj R., Korjenić E., Đug S. et al. 2016: Fishing bases of the canton Sarajevo for fishing area VI. *PMF University in Sarajevo, Sarajevo, Bosna and Hercegovina.* (in Bosnian)
- Škrijelj R., Lelo S., Drešković N. et al. 2013: Red List of endangered plants, animals and mushrooms in the Federation of Bosnia and Herzegovina. Red List of Fauna of the Federation of Bosnia and Herzegovina, book 3. *The Federal Ministry of Environment and Tourism, EU „Greenway“ Sarajevo and PMF University in Sarajevo, Sarajevo, Bosna and Hercegovina.* (in Bosnian)
- Škrijelj R., Muhamedagić S., Hamzić A. et al. 2008a: Fishing-economic basis for the fishing zone of the Central Bosnia Canton. *Faculty of Agriculture and Food Science, University in Sarajevo, Sarajevo, Bosna and Hercegovina.* (in Bosnian)
- Škrijelj R., Muhamedagić S., Hamzić A. et al. 2008b: Economic basis for the fishing area of Brčko District of B & H. *Faculty of Agriculture and Food Science, University in Sarajevo, Sarajevo, Bosna and Hercegovina.* (in Bosnian)
- Šlechtová V., Lusková V., Šlechta V. et al. 2000: Genetic differentiation of two diploid-polyploid complexes of spined loach, genus *Cobitis* (Cobitidae), in the Czech Republic, involving *C. taenia*, *C. elongatoides*, and *C. spp.*: Allozyme interpopulation and interspecific difference. *Folia Zool.* 49 (Suppl. 1): 67–78.
- Taler Z. 1953: Distribution and list of freshwater fish of Yugoslavia. *Glas. Prirod. Muzeja Srps. Zem. (serija B), knjiga 5–6, Beograd.* (in Serbo-Croatian/Croato-Serbian)
- Tutman P., Burić M. & Skaramuca B. 2012b: First substantiated record of the blackstriped pipefish, *Syngnathus abaster* (Actinopterygii, Syngnathiformes, Syngnathida) in the freshwaters of Bosnia and Herzegovina. *Acta Ichthyol. Pisc.* 42: 259–262.
- Tutman P., Glamuzina B., Dulčić J. & Zovko N. 2012a: Ichthyofauna of the Hutovo blato wetland (lower Neretva River, Bosnia and Herzegovina): status and vulnerability. *Croat. J. Fish.* 70: 169–185.
- Tutman P., Matić-Skoko S., Glamuzina B. et al. 2006: Some biological parameters of Neretvan spined loach, *Cobitis narentana* (Karaman, 1928) from Hutovo blato wetland, Bosnia and Herzegovina. In: Mrakovčić M., Buj I. & Zanella L. (eds.), Third International Conference: loaches of the genus *Cobitis* and related genera: biology, systematics, genetics, distribution, ecology and conservation. *Croatian Information and Documentation Society, Zagreb.*
- Tutman P., Šanda R., Glamuzina B. & Dulčić J. 2013: First confirmed record of *Pomatoschistus canestrinii* (Ninni, 1883) (Gobiidae) from Bosnia and Herzegovina. *J. Appl. Ichthyol.* 29: 937–939.
- Vegara M., Muhamedagić S., Hamzić A. et al. 2009: Mid-term improvement program of fisheries for the fishing area of Kalinovik municipality. *Norwegian University of Life Sciences (UMB), NORAGRIC, AAS, Norway. Norwegian SEE programme: Institutional collaboration between academic institutions in agriculture, forestry and veterinary medicine in Norway and Bosnia and Herzegovina, Croatia and Serbia and Montenegro, 2006-2009.* (in Bosnian)
- Vuković T. 1963: Fishes of Bosnia and Herzegovina (determination key). *The Institute for publishing textbooks, Sarajevo.* (in Serbo-Croatian/Croato-Serbian)
- Vuković T. 1977: Fishes of Bosnia and Herzegovina. *IGKRO „Svjetlost“, Sarajevo.* (in Serbo-Croatian/Croato-Serbian)
- Vuković T. & Ivanović B. 1971: Freshwater fishes of Yugoslavia. *Zemaljski muzej Sarajevo, Sarajevo.* (in Serbo-Croatian/Croato-Serbian)
- Zanella D., Mrakovčić M., Mustafić P. et al. 2009: Threatened fishes of the world, *Cobitis narentana* Karaman 1928 (Cobitidae). *Environ. Biol. Fishes* 86: 523–524.
- Zanella D., Mrakovčić M., Schneider D. et al. 2003: Growth of *Cobitis narentana* Karaman, 1928 in the Neretva River, Croatia. *Folia Biol. (Kraków)* 51 (Suppl.): 155–157.
- Zaplata R. & Taler Z. 1932: Fishes of Sarajevo and wider area. *Glasnik Zemaljskog Muzeja* 44: 1–32.