

## **Ecology and conservation of the Japanese flying squirrel *Pteromys momonga***

Author: Suzuki, Kei K.

Source: Journal of Vertebrate Biology, 72(23054)

Published By: Institute of Vertebrate Biology, Czech Academy of Sciences

URL: <https://doi.org/10.25225/jvb.23054>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Ecology and conservation of the Japanese flying squirrel *Pteromys momonga*

Kei K. SUZUKI

Kyushu Research Centre, Forestry and Forest Products Research Institute, Kumamoto, Japan;  
e-mail: pteromysuzuki@affrc.go.jp

► Received 24 May 2023; Accepted 16 June 2023; Published online 7 September 2023

**Abstract.** Flying squirrels have important roles in ecosystems, dispersing seeds and spores. However, flying squirrel species, which strongly depend on old mature forests, have declined in abundance due to the logging of mature forests. Guidelines for the conservation of flying squirrels have been developed, but it is vital to have an accurate understanding of their ecology to revise the guidelines to be more effective. The Japanese flying squirrel (*Pteromys momonga*, JFS), endangered in wide areas of Japan, is one of the flying squirrel species for which the least ecological information is available. This paper reviews individual reports and attempts to address knowledge gaps in JFS ecology to facilitate conservation strategies. Of particular importance is the fact that JFS uses planted Japanese cedar (*Cryptomeria japonica*) for nesting, gliding and winter forage, and JFS habitat may be strongly influenced by human activities such as forestry. Plantations are regularly logged, so management of plantations needs to be implemented with attention to the JFS habitat. It was also found that some ecological information necessary to develop guidelines for conservation, such as diet, gliding ability, and home-range size, is missing. Clarification of these issues is an important task.


**Key words:** food content, habitat use, nest site, plantation, reproduction

## Introduction

Gliding mammals can be divided into three main groups: flying squirrels, gliding marsupials, and colugos. Flying squirrels are the most diverse and widely distributed of these, inhabiting Asia, Europe, and North America (Jackson 2012) and playing essential roles in ecosystems as seed and spore dispersers (Maser et al. 1985, Nandini & Parthasarathy 2008) and as essential prey for predators vulnerable to environmental changes (Carey et al. 1992, Fryxell et al. 1998, Byholm et al. 2012). As such, they are of worldwide conservation interest (e.g. Smith et al. 2005, Koli 2016, Selonen & Mäkeläinen 2017). Flying squirrels are strongly reliant on old-growth forests and are sensitive to the isolation of forests (Smith

2012). They have experienced declines in abundance due to the logging of mature forests (Hokkanen et al. 1982, Holloway & Smith 2011). Although guidelines for their conservation have been developed, it is important to gain an accurate and comprehensive understanding of their ecology to revise the guidelines and make them more effective (Santangeli et al. 2013).

In Japan, three species of flying squirrels have been described: Japanese giant flying squirrel (GFS, *Petaurista leucogenys*), Japanese flying squirrel (JFS, *Pteromys momonga*), and Siberian flying squirrel (SFS, *Pteromys volans*) (Ohdachi et al. 2009). Japan is comprised of four main islands: Hokkaido, Honshu, Shikoku, and Kyushu, and the fauna inhabiting these islands vary along the Blakiston's Line



**Fig. 1.** Prefectural Red List rank of Japanese flying squirrel (JFS) in Japan. Since some prefectures do not distinguish between Critically Endangered and Endangered, the colour of both ranks is unified here. There is no prefecture that ranked JFS as Least Concern.

between Hokkaido and Honshu (Fig. 1). GFS and JFS are endemic species in Japan, found south of the Blakiston's Line in Honshu, Shikoku, and Kyushu Islands. In contrast, SFS is found in Hokkaido Island north of the Blakiston's Line but is also distributed in a wide area of the Eurasian Continent. The Hokkaido population of SFS is thought to have separated from the Eurasian population during the Holsteinian interglacial (Oshida et al. 2005) and is treated as an endemic subspecies, *P. volans orii*. Ecological information pertaining to these species was reviewed in "The wild mammals of Japan" (Ohdachi et al. 2009) and was subsequently revised in 2015 (Ohdachi et al. 2015). The GFS is more readily observable than the other two flying squirrels due to its larger size and vocalizations. In addition, the

fact that the GFS is found in shrine forests on flatland is probably a significant factor. Thus, the ecology of GFS has been actively studied since the 1980s (Ando & Imaizumi 1982, Ando et al. 1983) and has been reviewed in great detail (Kawamichi 2015). Research on SFS has seen a marked increase in the rate of investigation since the 1990s (Yamaguchi & Yanagawa 1995, Yanagawa 1999), with discoveries still being made frequently (Suzuki et al. 2013, Suzuki & Yanagawa 2019, Murakami et al. 2021). In particular, the ecology and conservation of SFS as a flagship species in the Eurasian Continent have been studied in depth (Selonen & Mäkeläinen 2017). The ubiquity of SFS in urban and sub-urban forests on flatlands makes it a species that is easy to observe, which has likely been a contributing factor to the




Fig. 2. Flowchart of the literature search.

advancement of the research. In contrast, although there were a few records of behavioural observations in captivity (Tezuka 1959, Ando et al. 1985), little was known about the ecology of JFS.

JFS has been categorized as of Least Concern on the IUCN Red List. The Ministry of the Environment of Japan has not placed JFS on the red list. Although JFS has been found in some prefectures, 88% of the prefectural governments are concerned about its potential extinction (Fig. 1). This is partly due to the difficulty in its detection, as its actual range is likely to be small. Direct observation of JFS is challenging due to their habitat preferences; JFS inhabit steep mountainous regions. However, the development of survey methods, such as nest box (Ando 2005) and camera trapping (Suzuki & Ando 2019), have enabled researchers to collect field data and have yielded new ecological insights. However, the findings on the ecology of JFS have often been reported simply as case studies and have never been discussed

comprehensively. This paper reviews individual reports and attempts to address knowledge gaps in JFS ecology to facilitate conservation strategies.

### Material and Methods

A flowchart of the literature search is shown in Fig. 2. I conducted an exhaustive search of the literature related to JFS using four search engines, such as Google Scholar, PubMed, Citation Information by the National Institute of Informatics, and NDL DIGITAL COLLECTIONS, because it is suggested that more than two databases should be used for a comprehensive search of the literature (Atkinson & Cipriani 2018). Only species names were used as search terms to ensure a comprehensive search of the literature on JFS: Japanese flying squirrel, *P. momonga*, and four Japanese names, such as “Nihon momonga”, “Honshu momonga”, “Hondo momonga”, and “Momonga”. However, not only JFS but also SFS is called Momonga in Japan. Thus, in

**Table 1.** Measurements of Japanese flying squirrels.

Prefecture	Sex	Head and body length (mm)	Body weight (g)	Cecum length (mm)	Condition
Fukushima	Female	147.5	99.5	132.0	Death
Kanagawa	Female	202.5	134.0	-	Alive
Kanagawa	Female	176.0	146.0	-	Alive
Kanagawa	Female	172.0	122.0	-	Alive
Kanagawa	Female	170.5	162.1	-	Alive
Kanagawa	Female	162.0	118.7	-	Alive
Kanagawa	Female	170.0	116.6	-	Alive
Kanagawa	Female	168.0	110.7	-	Alive
Kanagawa	Female	172.5	108.0	-	Alive and pregnant
Kanagawa	Female	157.0	157.8	-	Alive and pregnant
Kanagawa	Female	174.0	105.6	100.0	Death and pregnant
Kanagawa	Female	156.0	80.0	-	Death
Kanagawa	Male	158.0	94.7	110.0	Death
Yamanashi	Female	167.0	103.7	160.0	Death
Saitama	Female	165.0	91.8	97.0	Death

the literature, those related to JFS were collected, and those related to SFS excluded.

In addition, the inclusion of grey literature in the review article will improve research accuracy (Atkinson & Cipriani 2018). Thus, grey literature, such as essays and theses describing JFS in my possession, were used, even if they could not be found through the search engines listed above. However, since these references were not peer-reviewed, I did not gather information on the results of statistical analyses but only objective facts such as weight, litter size, and nesting material. From the collected literature, I gathered the following information: measurements of JFS, activity, food contents, nest site, habitat use, and reproduction.

## Results and Discussion

### Morphological characteristics

The head and body (HB) and tail (TL) of JFS range from 139 to 200 mm and 95 to 140 mm, respectively (Oshida 2009a). Hind-foot length (HF) ranges from 32 to 39 mm, excluding claws. Although JFS is slightly larger than SFS (Oshida 2009b), both species are difficult to distinguish visually due to their similar size. However, these two species can be identified by the difference in mammary number: ten for JFS (Oshida 2009a) and eight for SFS (Oshida 2009b).

Many illustrated books report the body weight (BW) of JFS as ranging from 150 to 220 g (Oshida 2009a, Jackson 2012). However, these values are overestimates. HB and BW of 15 JFS captured in Suzuki (2006) are presented in Table 1. While the HB measurements agreed with previous information, the BW of almost all individuals were less than 150 g. Two female individuals had offspring, and those weights were 134 g and 146 g, respectively. Also, the minimum BW of pregnant females was 108 g (Table 1). In addition, another study also reported a 146 g female giving birth (Kobayashi 2012a). The average weight of three 2-year-old females was reported to be 135.9 g (Okubo et al. 2014). In other words, even individuals weighing less than 150 g are mature. In addition, live males weighed 100 to 165 g, except for young living with their mothers (Kobayashi 2012a, Yano 2009, Okubo et al. 2015). Thus, based on previous reports (Table 1; Kobayashi 2012a, Okubo et al. 2014), I suggest revising the BW of adult JFS to be 108 to 173 g in females and 100 to 165 g in males. These values are approximately 20-30% lighter than previously described. This discrepancy is of particular significance, as it risks misidentifying a mature adult as a subadult.

### Movement

The nocturnal activity of the JFS commences shortly after sunset (Suzuki & Ando 2017). Upon leaving the nest, JFS typically defecates in the tree hosting the

Table 2. Records of Japanese flying squirrel.

Prefecture	Year	Elevation (m)	Method	Tree species	References
Aomori	1976 to 2016	270 to 300	Direct observation	Tdo, broad-leaved trees	Sasamori & Mineshita (2019)
Aomori	1995	-	Direct observation	Crj, broad-leaved trees	Sasamori & Mineshita (2019)
Aomori	2002	-	Faeces	Tdo	Sasamori & Mineshita (2019)
Aomori	2005	-	Nest box	Crj	Sasamori & Mineshita (2019)
Aomori	2013 to 2017	190	Direct observation	Tdo, broad-leaved trees	Sasamori & Mineshita (2019)
Fukushima	2002 to 2004	350 to 700	Faeces	Crj, Pd, broad-leaved trees	Iwasaki & Takahashi (2005)
Fukushima	2004	1,850	Camera trapping	Am	Iwasaki & Takahashi (2005)
Fukushima	2004	740	Direct observation	-	Yasuda & Yagihashi (2004)
Fukushima	2005	600 to 1,000	Camera trapping	Crj, Fc, Qc, At	Iwasaki (2012)
Fukushima	2005	820 to 1,200	Direct observation and faeces	Crj, Fc, Qc	Iwasaki & Takahashi (2005)
Fukushima	2005	680 to 750	Direct observation and faeces	broad-leaved trees	Iwasaki & Takahashi (2005)
Tokyo	-	650	Direct observation	Crj, Co, Mo, Fp, Cej, Acer sp.	Okazaki (2012)
Tokyo	1996	310	Direct observation	Crj, Co, Zs, Qm, Caj, Tn	Okazaki (2012)
Tokyo	2000 to 2001	929	Direct observation	Crj, Qc, Acer sp.	Asari (2012)
Tokyo	2002	440	Direct observation	Crj	Iwasaki (2012)
Tokyo	2010	240	Direct observation	Crj, Co, Zs, Cca, Cerasus sp., Caj, Ap	Okazaki (2012)
Kanagawa	1995	-	Direct observation and Nest box	Af, Crj, Co, Broad leaved trees	Yamaguchi (1997)
Kanagawa	2001	500	Nest box	-	Yamaguchi et al. (2004)
Kanagawa	2004	650 to 750	Nest box	Crj, Co, Qse, Qa	Ookubo & Ando (2005)
Kanagawa	2005 to 2008	560 to 850	Camera trapping	Crj, Co, Fc, Qse, Af	Suzuki & Ando (2019)
Tochigi	1991 to 1993	650 to 950	Nest box	Fc, Qc	Sato (1997)
Tochigi	1991 to 1993	990 to 1,080	Nest box	broad-leaved trees	Sato (1997)
Tochigi	1991 to 1993	1,160 to 1,450	Nest box	Fc, Td	Sato (1997)
Yamanashi	1981	800	Nest box	Qc, Qse	Ando (2005)
Yamanashi	2007 to 2008	1,350 to 1,600	Camera trapping	Fc, Qc, Bp, Ah	Suzuki & Ando (2019)
Yamanashi	2008	1,074 to 1,399	Camera trapping	Broad-leaved and coniferous trees	Matsubayashi et al. (2009)

Table 2. continued

Prefecture	Year	Elevation (m)	Method	Tree species	References
Shizuoka	2004 to 2005	830 to 863	Nest box	Crj, Co	Takanaka et al. (2008)
Fukui	1993	930	Direct observation	Qc, Cs, Cb	Matsumura (1995)
Fukui	1995	140	Direct observation	Crj, Co, Qm, Ns	Yanagawa et al. (1996)
Toyama	1997	1,180	Direct observation	Fc	Murayama & Nambu (1998)
Nagano	2016	1,480 to 2,470	Nest box and camera trapping	Qc, Be, Av, Am	Kikuchi & Izumiya (2020)
Kyoto	1984 to 1991	360	Direct observation	Fc	Nimura et al. (1997)
Shiga	1996	400 to 700	Nest box	Crj, Qc, Qs, Pr, Cco	Ando (2005)
Mie	1978 to 1995	550	Direct observation	broad-leaved trees	Shimizu (2014)
Mie	1997	400	Direct observation	Crj, Co	Shimizu (2014)
Mie	1998	340	Direct observation	Crj, Co	Shimizu (2014)
Mie	2001	400	Direct observation	-	Shimizu (2014)
Mie	2003	200	Direct observation	Crj, Co	Shimizu (2014)
Mie	2005	700	Direct observation	Crj, Co, Qse	Shimizu (2014)
Mie	2005	1,150	Direct observation	broad-leaved trees	Shimizu (2014)
Mie	2006	370	Direct observation	Crj, Co	Shimizu (2014)
Mie	2014	470	Direct observation	Crj, Co	Shimizu (2014)
Yamaguchi	1996	500	Nest box	Crj, Co, natural forest	Fukamachi (2004)
Tottori	2010 to 2011	660 to 750	Nest box	Crj, Fc, Qc, Ct, Ic, At, Pr, Af	Kobayashi (2012a)
Ehime	-	1,000	Direct observation	Crj, broad-leaved trees	Yano (2009)
Ehime	1983 to 1985	1,100 to 1,325	Direct observation	Fc	Yano (2009)
Ehime	1992 to 2005	650	Direct observation	Crj, Co, Zs, Qs, Acer sp., Pr	Yano (2009)
Ehime	1992 to 2008	500 to 800	Direct observation	Crj, broad-leaved trees	Yano (2009)
Ehime	1998	-	Direct observation	broad-leaved trees	Yano (2009)
Ehime	1999	400 to 500	Direct observation	-	Yano (2009)
Ehime	2004	200	Nest box	Crj, Co	Yano (2009)
Ehime	2004 to 2005	100 to 220	Direct observation and Nest box	broad-leaved trees	Yano (2009)
Ehime	2006	1,030	Nest box	Fc, Qac, Sm, Acer sp.	Furukawa & Miyamoto (2010)
Ehime	2009 to 2010	650	Nest box and camera trapping	Crj, Co, Zs, Qs, Acer sp., Pr	Yano (2009)



Table 2. continued

Prefecture	Year	Elevation (m)	Method	Tree species	References
Tokushima	1996	1,100	Direct observation	-	Kawamichi (2009)
Tokushima	2007	730	Direct observation	-	Kawamichi (2009)
Kochi	2007	600 to 840	Direct observation	-	Kawamichi 2009)
Kumamoto	2006 to 2008	680 to 855	Nest box	Fc, Qm, Qac, Acer sp, Af, Ts	Sakata et al. (2009)
Kumamoto	2008	755 to 860	Nest box and camera trapping	Crj, Fc, Qm, Qac, Acer sp, Af, Ts	Sakata et al. (2009)
Kumamoto	2009 to 2010	400 to 480	Camera trapping	Crj, Co, Le, Cc, Qs	Sakata et al. (2011)
Kumamoto	2010	400 to 600	Camera trapping	Le, Cc, Qs, Qac, Qg	Sakata et al. (2010)
Miyazaki	2006	1,000 to 1,607	Direct observation	Crj, Co, broad-leaved trees	Kabemura et al. (2010)
Miyazaki	2007	760	Camera trapping	Crj, Co, Qs, Caj, Af, Ts	Okubo et al. (2009)
Miyazaki	2008	900	Camera trapping	Pd, broad-leaved trees	Yasuda & Kurihara (2009)

Tree species: Crj – *Cryptomeria japonica*; Co – *Chamaecyparis obtuse*; Tdo – *Thujaopsis dolaburata*; Af – *Abies firma*; Ah – *Abies homolepis*; Av – *Abies veitchii*; Am – *Abies mariesii*; Ts – *Tsuga sieboldii*; Td – *Tsuga diversifolia*; Pd – *Pinus densiflora*; Tn – *Torreya nucifera*; Fc – *Fagus crenata*; Qs – *Quercus salicina*; Qse – *Quercus serrata*; Qa – *Quercus acutissima*; Qc – *Quercus crispula*; Qm – *Quercus myrsinaefolia*; Qac – *Quercus acuta*; Qg – *Quercus gilva*; Le – *Lithocarpus edulis*; Cc – *Castanopsis cuspidate*; Bp – *Betula platyphylla*; Be – *Betula ermanii*; Ct – *Carpinus tschonoskii*; Zs – *Zelkova serrata*; Cs – *Chengiopteris sciadophylloides*; Cca – *Cinnamomum camphora*; Ns – *Neolitsea sericea*; Mo – *Magnolia obovate*; Ic – *Ilex crenata*; Caj – *Camellia japonica*; Sm – *Stewartia monadelpha*; Cb – *Clethra barbinervis*; At – *Aesculus turbinata*; Ap – *Acer palmatum*; Cco – *Cornus controversa*; Fp – *Fraxinus platypoda*; Cej – *Cercidiphyllum japonicum*; Pr – *Pterocarya rhoifolia*.






Fig. 3. Histogram showing seasonal changes in litter size of Japanese flying squirrel.

nest or in adjacent trees (Iwasaki & Takahashi 2009). After that, they usually move by gliding, but the details of their gliding ability are not known at all. If the forest is fragmented by logging, the gap size should be based on their glide ratio for JFS habitat conservation. Thus, clarifying the glide ratio of JFS is an important issue and logging based on gliding ability is desired.

However, the movements of JFS may be similar to those of SFS, a closely related species with a similar body size. Given that SFS have an average glide ratio of 1.7-2.0 (Asari et al. 2007, Suzuki et al. 2012, Suzuki & Yanagawa 2019), it is likely that JFS possess a comparable gliding capacity. In addition, the dispersal distances of JFS are unknown, but SFS is observed to move long distances (Selonen et al. 2010). Because dispersal distance is essential information for determining conservation areas and gene flow of the species (Clobert et al. 2004), another critical issue is determining the dispersal distance of JFS.

Camera trapping has revealed that JFS are often photographed on the trunk of Japanese cedar (*Cryptomeria japonica*) with a high tree height (at least

14 m), suggesting that JFS may prefer these trees as landing sites for gliding (Suzuki & Ando 2019). Japanese cedar is in demand as timber because of its straight tree shape. Straight trees may be easier for JFS to land on. In addition, SFS selectively lands on tall trees that allow it to glide farther in the next glide (Suzuki et al. 2012, Suzuki & Yanagawa 2019). JFS would land in a tall tree for the same reason. Straight, tall trees are necessary for habitat as a migration route for JFS.

#### Food contents

JFS are considered to be folivorous (Oshida 2009a), foraging mainly on leaves, flowers (pollen), buds, seeds, cones, fruits, and bark, with the dietary composition changing seasonally. However, there is little information on the types of plants foraged by JFS, with only a few captive experiments and field observations available. Two JFS captured in the northern part of Honshu Island have been found to forage on leaves of Japanese cedar, cherry (*Cerasus* sp.), and beech (*Fagus crenata*), as well as cones of red pine (*Pinus densiflora*) (Iwasaki & Takahashi 2009). A JFS captured in the central part of Honshu Island was observed to forage on leaves of red (*Quercus acuta*)



and sawtooth (*Q. acutissima*) oaks, young leaves of northern Japanese hemlock (*Tsuga diversifolia*), and fruit and seed of loquat (*Eriobotrya japonica*) (Tezuka 1959). Captive JFS have also been reported to forage on Asian hazel (*Corylus heterophylla*) and Japanese green alder (*Alnus firma*) (Kurota 1941), but the parts selected, such as the leaf, bud, or seed, is unknown. Insectivory by JFS is controversial, as some records indicate that JFS foraged on orthoptera and beetles (Kurota 1941), while other records suggest the contrary (Tezuka 1959). In the wild, only one recorded instance of JFS foraging on Japanese zelkova (*Zelkova serrata*) (Okazaki 2012). From winter to early spring, however, faeces of JFS have been found to contain pollen of Japanese cedar (Ichikawa et al. 2004, Iwasaki & Takahashi 2009).

While detailed food contents of JFS are unknown, it is considered that JFS has a lower digestive capacity than GFS (Okubo et al. 2015). Herbivores of smaller body size require more protein and energy per unit body mass than larger species (Demment & Van Soest 1985), and arboreal mammals that have adapted to folivory tend to have a longer alimentary tract (Chivers & Hladik 1980). Mitsuzuka & Oshida (2018) revealed that the cecum length per HB of the GFS (0.85 and 0.89) with large body size (approx. 1,000 g) is longer than that of SFS (0.56), and this means that GFS better adapts to folivory than SFS. Because the caecum length per HB of JFS (Table 1; mean = 0.74, n = 5, SD = 0.176) is located between GFS and SFS, food contents of JFS may also be intermediate between those species.

Given the importance of food availability in determining animal distribution (Lurz et al. 2000) and its role in optimizing animal survival (Pyke 1984), knowledge of food contents is a critical component of habitat management (Birnie-Gauvin et al. 2017). However, there is limited knowledge of food resources for JFS, and further investigation into their diet is urgently needed. In contrast, the food contents of SFS are relatively well known, with 16 species of plants from seven families identified by direct observation (Fujimaki 1963, Yanagawa 1999, Asari et al. 2008, Nambu & Yanagawa 2010). The large disparity in the amount of knowledge regarding the food contents of JFS compared to the SFS is likely attributable to the difficulty in directly observing JFS foraging in steep mountain environments, in contrast to SFS, which inhabit urban forests. Nevertheless, recent findings based on DNA analysis of faeces (Murakami et al. 2021) suggest that the diet of JFS can be identified. Faeces of JFS accumulate at the base of trees with

cavities, where JFS nest (Iwasaki & Takahashi 2009). Moreover, due to the ease of collecting SFS faeces by placing umbrellas upside down at the base of trees with cavities (Suzuki et al. 2011a), JFS faeces could similarly be collected. Identifying the contents of JFS by faeces may be more feasible than by direct observation.

### Nest site uses

JFS uses tree cavities as their primary nesting sites for resting and breeding. Although they are unable to create their own cavities, they utilize cavities created by decaying branches and old nests of woodpeckers. JFS uses multiple nests (Kobayashi 2012a). In addition, JFS generally exhibit solitary nesting, but communal nesting is observed from autumn to winter (Suzuki et al. 2008, Kobayashi 2013, Kikuchi et al. 2022). In SFS populations in Finland, communal nesting is likely associated with mating (Selonen et al. 2014). However, in JFS populations, communal nesting may involve only females or only males (Suzuki et al. 2008, Kobayashi 2013), indicating that the function of communal nesting in JFS may not necessarily be related to mating. The purpose of communal nesting in JFS remains to be determined.

The long and short diameters of the tree cavities used by JFS average 8.2 and 6.9 cm, respectively, with a minimum of 4.6 cm (Suzuki et al. 2011b). However, since JFS will also use nest boxes with entrances larger than 3.5 cm (Sakata et al. 2009), they will likely use tree cavities with a similar entrance size. The average height of nesting cavities is 6.2 m (Suzuki et al. 2011b). In experiments with nest boxes, JFS tended to prefer higher nest boxes (Okubo & Ando 2005, Kobayashi 2014a). Studies have shown that Japanese cedar cavities are preferred in Kanagawa Prefecture (Suzuki et al. 2011b). Additionally, although JFS nests in a cavity of Japanese white bark magnolia (*Magnolia obovata*) (Iwasaki 2012), nesting in cavities of Japanese cedar have been frequently confirmed in also other areas (Asari 2012, Iwasaki 2012, Okazaki 2012). Thus, the cavities with an entrance of more than 3.5 cm with openings high on the trunks of Japanese cedar appear to be important for the conservation of JFS.

JFS predominantly utilizes Japanese cedar bark as its primary nesting material (Ando 2005, Suzuki et al. 2008, Sakata et al. 2009). However, if Japanese cedar is absent, JFS uses white birch bark and moss as the materials (Kakuta 2006). It is hypothesized that the preference for Japanese cedar bark is likely attributed to its superior waterproofing and insulating properties (Kobayashi 2012b, 2014b). The preference



of JFS for the cedar cavities may be due to the easy availability of nest materials.

### Habitat uses

JFS prefer an environment where Japanese cedar and cypress (*Chamaecyparis obtuse*) plantations and natural broad-leaved forests are adjacent to each other (Suzuki et al. 2008). However, they do not necessarily inhabit only this environment. Although not quantitatively evaluated, several reports of JFS observations have been documented (Table 2). It appears that JFS habitats encompass a variety of vegetation, with approximately 60% of their habitats containing plantations of Japanese cedar and cypress. This situation may be due to the utility of Japanese cedar as a gliding path, winter food item, nest site, and nest material. These reports were mainly based on nest box use, camera trapping, and direct observation. In natural habitats without plantations, the main vegetation seems to be Pinaceae, Fagaceae, and Betulaceae.

JFS has never been surveyed for their home range. Determining the size of their home range and what type of environment they prefer within the home range is an important issue for their habitat conservation.

### Reproduction

JFS have two breeding seasons (spring and summer) per year because their neonates, described as hairless, were found in mid-March (Kakuta 2006) and mid-August (Kobayashi 2012a). The neonates in summer weighed around 7 g. Because the mean weight of neonates upon birth is 4.6 g in captivity and the daily body weight gain ranges from 1.2 to 1.3 g (Kakuta 2006), the neonates must be born within a few days. Although mating has been observed from January to February (Kikuchi et al. 2022), summer mating has probably never been observed. Given the 40 to 42-day gestation period of the SFS (Airapetyants & Fokin 2003), which is almost the same size and belongs to the same genus, there will likely be a mating season in July. However, it is unknown whether the same individuals breed twice a year.

The litter size of JFS vary widely from one to eight (Kobayashi 2012a). Although it has rarely been mentioned, JFS litter size will likely change seasonally. Based on 16 references (Sakaguchi 1957, Shirai 1963, Ueno et al. 1996, Yanagawa et al. 1996, Suzuki 2001, Kakuta 2006, Suzuki et al. 2008, Hosoda 2009, Oshida 2009a, Sakata et al. 2009, Yano 2009, Imaizumi 2012, Kobayashi 2012a, 2019, 2022, 2023) and one personal

observation (two young in spring), the litter size in spring ranges one to five (37 young in 18 litters) and that in summer is three to eight (84 young in 16 litters) (Fig. 3). It should be noted, however, that some offspring may have died prior to the observations, and these litter sizes may be slightly underestimated.

Although seasonal changes in litter sizes are shown in southern flying squirrels (*Glaucomys volans*), average litter sizes for spring and summer were 2.4 and 3.4, respectively (Stapp & Mautz 1991), which is not as large a difference as found in the JFS. The summer litter size of JFS is remarkably large for a flying squirrel species, considering that almost all *Pteromyini species* in the family Sciuridae generally have small litter sizes of one to three (Goldingay 2000, Hayssen 2008). It is unknown why JFS litter size changes seasonally. With such an unusual litter size for an arboreal squirrel, JFS is a vital research target for understanding the evolution of litter size in Sciuridae, and they are worthy of conservation.

### Conservation and Conclusions

JFS is one of the least ecologically understood species among flying squirrels. This review highlights knowledge gaps: gliding ability, diet composition, and home range size. In addition, recent observations of the species have been recorded both directly and through the use of nest boxes and camera trapping (Table 2). This review provides some insight into the conservation of JFS; of particular importance is that JFS uses planted Japanese cedar for nesting, gliding, and winter forage. These conclusions indicate that JFS habitat may be strongly influenced by human activities such as forestry.

Despite being generally thought of as having low biodiversity, plantations actually serve as habitats for many species (Brockerhoff et al. 2008). In addition, evidence suggests that plantations have an important role in providing habitat to several endangered species (Brockerhoff et al. 2005, 2008, Barbaro et al. 2009, Berndt et al. 2009). For reasons of human economics, it is impossible to eliminate plantations altogether. Recently, therefore, management regimes of plantations that promote biodiversity have garnered great international interest in the conservation of endangered species (Castano-Villa et al. 2019, Gadoth-Goodman & Rothstein 2020, Jamhuri et al. 2020, Wang et al. 2022).

Plantations are regularly logged, so management of plantations needs to be implemented with attention

to the JFS habitat. For instance, it is preferable to avoid cutting down cavity trees and trees in their vicinity, and if it is inevitable, then a nest box should replace the nest. In addition, the creation of large habitat gaps through thinning should be avoided. Since the glide ability of the JFS is unclear, it is not possible to state here how long of a gap will make it difficult for the JFS

to move. Identifying gliding ability is a particularly important issue for developing guidelines for their conservation.

### Acknowledgements

---

*I am grateful to Dr M. Yasuda for helping collect literature.*



## Literature

- Airapetyants A.E. & Fokin I.M. 2003: Biology of European flying squirrel *Pteromys volans* L. (Rodentia: Pteromyidae) in the north-west of Russia. *Russ. J. Theriol.* 2: 105–113.
- Ando M. 2005: Improvement of nest box investigation techniques for study of arboreal rodents. *Mammal. Sci.* 45: 165–176.
- Ando M., Funakoshi K. & Shiraishi S. 1983: Use patterns of nests by the Japanese giant flying squirrel, *Petaurista leucogenys*. *Sci. Bull. Fac. Agric. Kyushu Univ.* 38: 27–43.
- Ando M. & Imaizumi Y. 1982: Habitat utilization of the white-cheeked giant flying squirrel *Petaurista leucogenys* in a small shrine grove. *J. Mamm. Soc. Japan* 9: 70–81. (in Japanese with English abstract)
- Ando M., Shiraishi S. & Uchida T.A. 1985: Feeding behaviour of three species of squirrels. *Behaviour* 95: 76–86.
- Asari Y. 2012: Sightings of *Pteromys momonga* at the study site of *Petaurista leucogenys* in Tokyo. *Sciurid Information* 28: 7–8. (in Japanese)
- Asari Y., Yamaguchi Y. & Yanagawa H. 2008: Field observations of the food items of the Siberian flying squirrel, *Pteromys volans orii*. *J. Jpn. Wild. Res. Soc.* 33: 7–11.
- Asari Y., Yanagawa H. & Oshida T. 2007: Gliding ability of the Siberian flying squirrel *Pteromys volans orii*. *Mamm. Stud.* 32: 151–154.
- Atkinson L.Z. & Cipriani A. 2018: How to carry out a literature search for a systematic review: a practical guide. *BJ Psych. Adv.* 24: 74–82.
- Barbaro L., Couzi L., Bretagnolle V. et al. 2009: Multi-scale habitat selection and foraging ecology of the Eurasian hoopoe (*Upupa epops*) in pine plantations. In: Brockerhoff E.G., Jactel H., Parrotta J.A. et al. (eds.), *Plantation forests and biodiversity: oxymoron or opportunity?* Springer, Dordrecht, Netherlands: 149–163.
- Berndt L.A., Brockerhoff E.G. & Jactel H. 2009: Relevance of exotic pine plantations as a surrogate habitat for ground beetles (Carabidae) where native forest is rare. In: Brockerhoff E.G., Jactel H., Parrotta J.A. et al. (eds.), *Plantation forests and biodiversity: oxymoron or opportunity?* Springer, Dordrecht, the Netherlands: 247–261.
- Birnie-Gauvin K., Peiman K.S., Raubenheimer D. & Cooke S.J. 2017: Nutritional physiology and ecology of wildlife in a changing world. *Conserv. Physiol.* 5: cox030.
- Brockerhoff E.G., Berndt L.A. & Jactel H. 2005: Role of exotic pine forests in the conservation of the critically endangered New Zealand ground beetle *Holcaspis brevicula* (Coleoptera: Carabidae). *N. Z. J. Ecol.* 29: 37–43.
- Brockerhoff E.G., Jactel H., Parrotta J.A. et al. 2008: Plantation forests and biodiversity: oxymoron or opportunity? *Biodivers. Conserv.* 17: 925–951.
- Byholm P., Burgas D., Virtanen T. & Valkama J. 2012: Competitive exclusion within the predator community influences the distribution of a threatened prey species. *Ecology* 93: 1802–1808.
- Carey A.B., Horton S.P. & Biswell B.L. 1992: Northern spotted owls: influence of prey base and landscape character. *Ecol. Monogr.* 62: 223–250.
- Castano-Villa G.J., Estevez J.V., Guevara G. et al. 2019: Differential effects of forestry plantations on bird diversity: a global assessment. *For. Ecol. Manag.* 440: 202–207.
- Chivers D.J. & Hladik C.M. 1980: Morphology of the gastrointestinal tract in primates: comparisons with other mammals in relation to diet. *J. Morphol.* 166: 337–386.
- Clobert J., Ims R.A. & Rousset F. 2004: Causes, mechanisms and consequences of dispersal. In: Hanski I. & Gaggiotti O.E. (eds.), *Ecology, genetics and evolution of metapopulations.* Elsevier, Amsterdam, the Netherlands: 307–335.
- Demment M.W. & Soest P.J. Van 1985: A nutritional explanation for body-size patterns of ruminant and nonruminant herbivores. *Am. Nat.* 125: 641–672.
- Fryxell J.M., Falls J.B., Falls E.A. & Brooks R.J. 1998: Long-term dynamics of small-mammal populations in Ontario. *Ecology* 79: 213–225.
- Fujimaki Y. 1963: Observations on the Japanese flying squirrel, *Pteromys volans orii* Kuroda, in captivity. *J. Mamm. Soc. Japan.* 2: 42–45. (in Japanese with English summary)
- Fukamachi O. 2004: Fun nest box. *Sciurid Information* 14: 5–7. (in Japanese)
- Furukawa M. & Miyamoto T. 2010: A record of the Japanese flying squirrel *Pteromys momonga* for Uwajima city, Ehime Prefecture. *Bull. Shikoku Inst. Nat. Hist.* 6: 18–21. (in Japanese)
- Gadoth-Goodman D. & Rothstein D.E. 2020: Alternative silvicultural approaches to managing Jack pine plantations for endangered species habitat and forest products. *For. Sci.* 66: 210–220.
- Goldingay R.L. 2000: Gliding mammals of the world: diversity and ecological requirements. In: Goldingay R. & Scheibe J.S. (eds.), *Biology of gliding mammals.* Filander Verlag, Furth, Germany: 9–44.
- Hayssen V. 2008: Reproductive effort in squirrels: ecological, phylogenetic, allometric, and latitudinal patterns. *J. Mammal.* 89: 582–606.



- Hokkanen H., Törmälä T. & Vuorinen H. 1982: Decline of the flying squirrel *Pteromys volans* L. populations in Finland. *Biol. Conserv.* 23: 273–284.
- Holloway G.L. & Smith W.P. 2011: A meta-analysis of forest age and structure effects on northern flying squirrel densities. *J. Wildl. Manag.* 75: 668–674.
- Hosoda T. 2009: Reproduction of Japanese flying squirrel. *Tokyo Zoo Net News*. [https://www.tokyo-zoo.net/topic/topics\\_detail?kind=news&link\\_num=11936](https://www.tokyo-zoo.net/topic/topics_detail?kind=news&link_num=11936) (in Japanese)
- Ichikawa T., Motoki T., Hara R. et al. 2004: Seasonal changes in habitats of Japanese flying squirrels in Nagano Prefecture. *Proceedings of Japanese Society of Environmental Entomology and Zoology, Osaka, Japan.* (in Japanese)
- Imaizumi Y. 2012: If I build a small house in the forest. In: Imaizumi T. (ed.), *My mountain hut diary. RONSOSHA, Tokyo, Japan: 72–75.* (in Japanese)
- Iwasaki Y. 2012: Sympatric distribution of *Petaurista leucogenys* and *Pteromys momonga* in Fukushima Prefecture and Tokyo. *Sciurid Information* 28: 2–6. (in Japanese)
- Iwasaki Y. & Takahashi M. 2005: Inhabitation of Japanese flying squirrel in Fukushima Prefecture. *Fukushima Seibutsu* 48: 13–16. (in Japanese)
- Iwasaki Y. & Takahashi M. 2009: Excrements and other signs of *Pteromys momonga* for detecting its habitat. *Sciurid Information* 22: 16–19. (in Japanese)
- Jackson S.M. 2012: *Gliding mammals of the world.* Csiro Publishing, Melbourne, Australia.
- Jamhuri J., Edinoor M.A., Kamarudin N. et al. 2020: Higher mortality rates for large-and medium-sized mammals on plantation roads compared to highways in Peninsular Malaysia. *Ecol. Evol.* 10: 12049–12058.
- Kabemura Y., Kubota K., Kajiya K. et al. 2010: Monitoring of mammal sightings in research forest in Miyazaki. *Bull. Kyushu Univ. For.* 91: 29–33. (in Japanese)
- Kakuta A. 2006: Nest site selection and growth of Japanese flying squirrel. *Tokyo University of Agriculture, Kanagawa, Japan.* (in Japanese)
- Kawamichi T. 2009: New distributions of *Pteromys momonga* in Tokushima and Kochi Prefectures, Shikoku Island. *Sciurid Information* 22: 9–13. (in Japanese)
- Kawamichi T. 2015: Japanese giant flying squirrel. *Tsukiji Shokan Publishing, Tokyo, Japan.* (in Japanese)
- Kikuchi H. & Izumiyama S. 2020: Faunal survey of small mammals in the Nishikoma Research Forest at Shinshu University. *Bull. Shinshu Univ. AFC* 18: 85–91. (in Japanese)
- Kikuchi H., Izumiyama S. & Oshida T. 2022: Does communal nesting help thermoregulation in Japanese flying squirrels (*Pteromys momonga*) in winter? *Russ. J. Theriol.* 21: 38–44.
- Kobayashi T. 2012a: Some ecological characteristics of small Japanese flying squirrels *Pteromys momonga* in a forest of Tottori Prefecture. *Nat. Environ. Sci. Res.* 25: 15–22.
- Kobayashi T. 2012b: Why do small Japanese flying squirrels, *Pteromys momonga*, prefer to use bark of Japanese cedar as a nest material? *Nat. Environ. Sci. Res.* 25: 23–28.
- Kobayashi T. 2013: An analysis of factors inducing use of a nest by plural individuals in Japanese flying squirrels, *Pteromys momonga*. *Bull. Tottori Prefect. Mus.* 50: 55–59.
- Kobayashi T. 2014a: Studies on habitat preference using nest boxes of the Japanese flying squirrel, the Japanese dormouse and the small Japanese field mouse in the Ashizu Valley, Tottori Prefecture, Japan. *Bull. Tottori Prefect. Mus.* 51: 1–6.
- Kobayashi T. 2014b: Why do small Japanese flying squirrels, *Pteromys momonga*, prefer Japanese cedar bark as a nest material? II. Study of heat-trapping capacity. *Nat. Environ. Sci. Res.* 27: 9–13.
- Kobayashi T. 2019: Professor, Japanese ratsnake is closing in on the Japanese flying squirrel family. *Tsukiji Shokan Publishing Co. Ltd, Tokyo, Japan.* (in Japanese)
- Kobayashi T. 2022: Ethologist gets angry at Japanese flying squirrel. *Yama-Kei Publishers Co. Ltd, Tokyo, Japan.* (in Japanese)
- Kobayashi T. 2023: Young Japanese flying squirrels can distinguish family odors! In: Kibayashi T. (ed.), *Professor, toads can't eat woodlouse because of difficulty in choosing.* *Tsukiji Shokan Publishing Co. Ltd, Tokyo, Japan: 139–162.* (in Japanese)
- Koli V.K. 2016: Biology and conservation status of flying squirrels (Pteromyini, Sciuridae, Rodentia) in India: an update and review. *Proc. Zool. Soc.* 69: 9–21.
- Kurota K. 1941: Mammals in Musashino. In: Tamura T. & Honda S. (eds.), *Musashino. Kagaku Shugi Kougyousya, Tokyo, Japan: 352–393.* (in Japanese)
- Lurz P.W.W., Garson P.J. & Wauters L.A. 2000: Effects of temporal and spatial variations in food supply on the space and habitat use of red squirrels (*Sciurus vulgaris* L.). *J. Zool.* 251: 167–178.
- Maser Z., Maser C. & Trappe J.M. 1985: Food habits of the northern flying squirrel (*Glaucomys sabrinus*) in Oregon. *Can. J. Zool.* 63: 1084–1088.
- Matsubayashi H., Ishizaka S., Nakagawa T. & Nakamura Y. 2009: Middle to large mammalian



- fauna in the headwater of Tamagawa River. *J. Agric. Sci. Tokyo Univ. Agric.* 54: 110–115. (in Japanese with English abstract)
- Matsumura T. 1995: Habitats of *Pteromys momonga* in Fukui Prefecture. *Ciconia* 4: 65–69. (in Japanese)
- Mitsuzuka W. & Oshida T. 2018: Feeding adaptation of alimentary tract length in arboreal squirrels. *Mamm. Study* 43: 125–131.
- Murakami S., Kikuchi H. & Oshida T. 2021: Analysis of feeding habit of small rodent by DNA metabarcoding: seasonal change of feeding habit of *Pteromys volans orii*. *Res. Bull. Obihiro Univ.* 42: 77–86. (in Japanese with English abstract)
- Murayama M. & Nambu H. 1998: Note on small Japanese flying squirrel from Arimine, Toyama Prefecture, central Japan. *Bull. Toyama Sci. Mus.* 21: 127–128. (in Japanese)
- Nambu A. & Yanagawa H. 2010: Food selection of the Siberian flying squirrel, *Pteromys volans orii*, in winter. *J. Jpn. Wildl. Res. Soc.* 35: 22–25. (in Japanese)
- Nandini R. & Parthasarathy N. 2008: Food habits of the Indian giant flying squirrel (*Petaurista philippensis*) in a rain forest fragment, Western Ghats. *J. Mammal.* 89: 1550–1556.
- Nimura K., Nakashima T. & Yamanaka N. 1997: A witness record of wild animals in Ashiu. *Rep. Kyoto Univ. For.* 30: 59–72. (in Japanese)
- Ohdachi S.D., Ishibashi Y., Iwasa M.A. et al. 2009: The wild mammals of Japan. *Shoukadoh Book Sellers, Kyoto, Japan.*
- Ohdachi S.D., Ishibashi Y., Iwasa M.A. et al. 2015: The wild mammals of Japan, 2<sup>nd</sup> ed. *Shoukadoh Book Sellers, Kyoto, Japan.*
- Okazaki H. 2012: *Petaurista leucogenys* and *Pteromys momonga* living together at three shrines in Tokyo. *Sciurid Information* 28: 9–11. (in Japanese)
- Okubo Y., Eto T. & Morita T. 2009: Investigation on the distribution of *Pteromys momonga* in Miyazaki Prefecture, Kyushu Island. *Sciurid Information* 22: 14–15. (in Japanese)
- Okubo Y., Sakamoto S., Kashimura A. & Morita T. 2014: Effects of ambient temperature and daylength on body temperature, body mass and food consumption in the Japanese flying squirrel *Pteromys momonga*. *Jpn. J. Environ. Entomol. Zool.* 25: 1–9. (in Japanese with English abstract)
- Okubo Y., Shichijo H., Watanabe D. et al. 2015: Digestion of dietary fiber by two flying squirrel species, *Petaurista leucogenys* and *Pteromys momonga*: a comparative study. *Jpn. J. Environ. Entomol. Zool.* 26: 29–35. (in Japanese with English abstract)
- Ookubo Y. & Ando M. 2005: Nest box height preference of Japanese flying squirrel. *Sciurid Information* 16: 9–11. (in Japanese)
- Oshida T. 2009a: *Pteromys momonga* Temminck, 1844. In: Satoshi D.O., Yasuyuki I., Masahiro A.I. & Takashi S. (eds.), The wild mammals of Japan. *Shoukadoh Book Sellers, Kyoto, Japan*: 194–195.
- Oshida T. 2009b: *Pteromys volans* (Linnaeus, 1758). In: Satoshi D.O., Yasuyuki I., Masahiro A.I. & Takashi S. (eds.), The wild mammals of Japan. *Shoukadoh Book Sellers, Kyoto, Japan*: 196–197.
- Oshida T., Abramov A., Yanagawa H. & Masuda R. 2005: Phylogeography of the Russian flying squirrel (*Pteromys volans*): implication of refugia theory in arboreal small mammal of Eurasia. *Mol. Ecol.* 14: 1191–1196.
- Pyke G.H. 1984: Optimal foraging theory: a critical review. *Annu. Rev. Ecol. Syst.* 15: 523–575.
- Sakaguchi K. 1957: On three fleas found on Eurasian small flying-squirrel, *Pteromys volans amygdali* (Thomas, 1906). *Med. Entomol. Zool.* 8: 160–166. (in Japanese)
- Sakata T., Nakazono T., Kaoka H. et al. 2009: Confirmation of Japanese flying squirrel (*Pteromys momonga*) with nest boxes in Gokanosho and Naidaijin, Kumamoto Prefecture, Japan. *Bull. Kumamoto Wildl. Soc.* 5: 11–20. (in Japanese)
- Sakata T., Yasuda M. & Nagamine S. 2010: Confirmation of Japanese flying squirrel (*Pteromys momonga*) and Japanese dormouse (*Glirulus japonicus*) in Ookawa, Minamata city, Kumamoto Prefecture, Japan. *Bulletin of the Kumamoto Wildlife Society* 6: 23–28. (in Japanese)
- Sakata T., Yasuda M. & Nakazono T. 2011: Survey of arboreal rodent community using nest boxes and camera traps in Kumamoto Prefecture, Kyushu Island, Japan. *Sciurid Information* 26: 8–12. (in Japanese)
- Santangeli A., Wistbacka R., Hanski I.K. & Laaksonen T. 2013: Ineffective enforced legislation for nature conservation: a case study with Siberian flying squirrel and forestry in a boreal landscape. *Biol. Conserv.* 157: 237–244.
- Sasamori K. & Mineshita K. 2019: New records and literature of Japanese flying squirrel (*Pteromys momonga*) in Aomori Prefecture, Japan. *J. Nat. Hist. Aomori* 24: 94–96.
- Sato Y. 1997: Distribution records of mammals using nest boxes for birds in the Kuriyama district, Tochigi. *Bull. Tochigi Prefect. Mus.* 14: 21–31. (in Japanese)
- Selonen V., Hanski I.K. & Painter J.N. 2010: Gene flow and natal dispersal in the Siberian flying squirrel



- based on direct and indirect data. *Conserv. Genet.* 11: 1257–1264.
- Selonen V., Hanski I.K. & Wistbacka R. 2014: Communal nesting is explained by subsequent mating rather than kinship or thermoregulation in the Siberian flying squirrel. *Behav. Ecol. Sociobiol.* 68: 971–980.
- Selonen V. & Mäkeläinen S.L.M. 2017: Ecology and protection of a flagship species, the Siberian flying squirrel. *Hystrix* 28: 134–146.
- Shimizu Z. 2014: The distribution of Japanese flying squirrel *Pteromys momonga* in Mie Prefecture. *Mie Shizenshi* 14: 72–75. (in Japanese)
- Shirai K. 1963: Stories of Japanese flying squirrels and giant Japanese flying squirrels. In: Hayashi T. (ed.), Boys and girls, Japan animal chronicles. *Maki Shoten, Tokyo, Japan*: 49–67. (in Japanese)
- Smith W.P. 2012: Sentinels of ecological processes: the case of the northern flying squirrel. *Bioscience* 62: 950–961.
- Smith W.P., Gende S.M. & Nichols J.V. 2005: The northern flying squirrel as an indicator species of temperate rain forest: test of a hypothesis. *Ecol. Appl.* 15: 689–700.
- Stapp P. & Mautz W.W. 1991: Breeding habits and postnatal growth of the southern flying squirrel (*Glaucomys volans*) in New Hampshire. *Am. Midl. Nat.* 126: 203–208.
- Suzuki K. 2001: Beautiful day for Japanese flying squirrel. *Seikosha, Tokyo, Japan*. (in Japanese)
- Suzuki K. 2006: Effect of plantation on Japanese flying squirrel (*Pteromys momonga*) inhabiting in the Tanzawa Mountains. *Tokyo University of Agriculture, Kanagawa, Japan*. (in Japanese)
- Suzuki K.K. & Ando M. 2017: Seasonal changes in activity patterns of Japanese flying squirrel *Pteromys momonga*. *Behav. Process.* 143: 13–16.
- Suzuki K.K. & Ando M. 2019: Early and efficient detection of an endangered flying squirrel by arboreal camera trapping. *Mammalia* 83: 372–378.
- Suzuki K., Asari Y. & Yanagawa H. 2012: Gliding locomotion of Siberian flying squirrels in low-canopy forests: the role of energy-inefficient short-distance glides. *Acta Theriol. (Warsz)* 57: 131–135.
- Suzuki K., Mori S. & Yanagawa H. 2011a: Detecting nesting trees of Siberian flying squirrels (*Pteromys volans*) using their feces. *Mamm. Study* 36: 105–108.
- Suzuki K., Ogawa H., Amano T. & Ando M. 2008: Habitat preference and nest box use of the small Japanese flying squirrel *Pteromys momonga* in the Tanzawa Mountains. *J. Agric. Sci. Tokyo Univ. Agric.* 53: 13–18. (in Japanese with English abstract)
- Suzuki K., Sagawa M. & Yanagawa H. 2013: Nest cavity selection by the Siberian flying squirrel *Pteromys volans*. *Hystrix* 24: 187.
- Suzuki K., Shimamoto T., Takizawa Y. et al. 2011b: Nest site characteristics of *Pteromys momonga* in the Tanzawa Mountains. *Mammal. Sci.* 51: 65–69.
- Suzuki K.K. & Yanagawa H. 2019: Gliding patterns of Siberian flying squirrels in relation to forest structure. *iForest* 12: 114–117.
- Takanaka K., Ando M., Ogawa H. et al. 2008: Drowning of small mammals from falling into street gutters with constantly flowing water. *Mammal. Sci.* 48: 1–9. (in Japanese with English abstract)
- Tezuka H. 1959: Notes on some habits, especially the eating method of leaves, of *Pteromys volans amygdali* in captivity. *J. Mamm. Soc. Japan* 1: 132–134. (in Japanese)
- Ueno Y., Ashikaga K., Yasui H. & KuwaBara K. 1996: The mammals in Geihoku-cho, Hiroshima Prefecture. *Nat. Hist. Nishi-Chugoku Mountains* 1: 395–441. (in Japanese)
- Wang C., Zhang W., Li X. & Wu J. 2022: A global meta-analysis of the impacts of tree plantations on biodiversity. *Glob. Ecol. Biogeogr.* 31: 576–587.
- Yamaguchi Y. 1997: Japanese flying squirrel, giant Japanese flying squirrel, Japanese squirrel in the Tanzawa Mountains, Kanagawa Prefecture. *Sciurid Information* 2: 8–9. (in Japanese)
- Yamaguchi Y. & Yanagawa H. 1995: Field observations on circadian activities of the flying squirrel, *Pteromys volans orii*. *Mammal. Sci.* 34: 139–149. (in Japanese with English abstract)
- Yamaguchi Y., Yuzawa Y. & Yuzawa Y. 2004: Animals using one nest box. *Sciurid Information* 14: 8–10. (in Japanese)
- Yanagawa H. 1999: Ecological notes on the Russian flying squirrel (*Pteromys volans orii*) with a video camera. *Mammal. Sci.* 39: 181–183. (in Japanese)
- Yanagawa H., Oshida T., Taniguchi M. & Taketatsu K. 1996: Notes on the Japanese flying squirrel, *Pteromys momonga* captured in Fukui Prefecture. *J. Jpn. Wildl. Res. Soc.* 22: 8–16. (in Japanese)
- Yano S. 2009: Distribution of Japanese flying squirrels *Pteromys momonga* in Ehime Prefecture, Shikoku Island. *Sciurid Information* 22: 2–8. (in Japanese)
- Yasuda M. & Kurihara T. 2009: A record of Japanese flying squirrel with camera traps in Shiiba village, Miyazaki Prefecture, Japan. *Bull. Kumamoto Wildl. Soc.* 5: 31–35. (in Japanese)
- Yasuda M. & Yagihashi T. 2004: Small Japanese flying squirrel (*Pteromys momonga*) caught on barbed-wire fence. *Sciurid Information* 15: 1–3. (in Japanese)