

Third record of the Critically Endangered Brazilian Merganser *Mergus octosetaceus* in São Paulo state, south-east Brazil, after almost two centuries

Authors: Neto, Miguel Nema, Pereira, Fabiana D., Wuo, Ana L. M., and Neto, Carlos Zacchi

Source: Bulletin of the British Ornithologists' Club, 138(2) : 131-134

Published By: British Ornithologists' Club

URL: <https://doi.org/10.25226/bboc.v138i2.2018.a7>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Third record of the Critically Endangered Brazilian Merganser *Mergus octosetaceus* in São Paulo state, south-east Brazil, after almost two centuries

by Miguel Nema Neto, Fabiana D. Pereira, Ana L. M. Wuo & Carlos Zacchi Neto

Received 11 February 2018; revised 24 April 2018; published 22 June 2018

<http://zoobank.org/urn:lsid:zoobank.org:pub:9163F1AD-EC6C-495F-A068-59C2067983CD>

SUMMARY.—Brazilian Merganser *Mergus octosetaceus* is one of the most endangered bird species in the Americas and one of the rarest ducks in the world. We photographed the species in August 2017, at Serra do Mar State Park, Salesópolis, in eastern São Paulo state. This is the third documented record in São Paulo, but the first for approximately 200 years.

Brazilian Merganser *Mergus octosetaceus* is considered one of the rarest bird species in the Americas and among the most threatened waterfowl in the world, being currently considered Critically Endangered (Collar *et al.* 1992, BirdLife International 2018). It is adapted to watercourses in montane regions, inhabiting clear-water rivers especially those with rapids, and the species feeds on fish, aquatic insects and molluscs (Sick 1997, Carboneras *et al.* 2018).

Originally found across south-central Brazil and adjacent Argentina and Paraguay, the species' range is now drastically reduced (Collar *et al.* 1992), despite its more recent discovery further north than previously known (Braz *et al.* 2003). Intolerant of impacts



Figure 1. Male Brazilian Merganser *Mergus octosetaceus*, rio Claro, Salesópolis, São Paulo, Brazil, 25 August 2017 (Fabiana Dias Pereira)



Figure 2. Locality where observations were made in August and November 2017, rio Claro, Salesópolis, São Paulo, Brazil (Miguel Nema Neto)

to its environment and sensitive to human disturbance, activities such as agricultural expansion, pollution, dams and loss of riverine vegetation can all negatively impact the species (Hughes *et al.* 2006). Recent sightings evidence its survival in Argentina (Misiones) and perhaps even in Paraguay, but in Brazil it is considered extinct in the states of Mato Grosso do Sul, São Paulo, Rio de Janeiro and Santa Catarina (Collar *et al.* 1992, Lamas & Lins 2009, Carboneras *et al.* 2018). In Paraná, recent field work has failed to confirm the species' presence and the last record was now more than 20 years ago, in 1995 (Lamas & Lins 2009, Carboneras *et al.* 2018). Indeed, in recent decades *M. octosetaceus* has been recorded at just a few locations in Brazil, mainly within protected areas, especially Serra da Canastra National Park (Minas Gerais), Chapada dos Veadeiros National Park (Goiás) and Jalapão State Park (Tocantins) (Braz *et al.* 2003, Bianchi *et al.* 2005, Lamas 2006). It has also been reported in western Bahia state, although subsequent searches of this area were unsuccessful (Lamas & Lins 2009). Occasional records elsewhere in Minas Gerais perhaps involve wandering or dispersing birds (Lamas & Lins 2009), although to date there is no definitive evidence that the species moves far from its natal areas (Ribeiro *et al.* 2011).

In the state of São Paulo, the only well-documented record of the species involves a female (from a pair) collected by J. Natterer in August 1820, at Itararé (24°07'S, 49°20'W), on the border with Paraná, and now in the Natural History Museum, Tring, UK (Collar *et al.* 1992). In addition Stresemann (1954), repeated by Collar *et al.* (1992), mentioned that in the previous year, 1819, Frederick Sellow took specimens (now in Berlin) during his travels through Rio de Janeiro, Minas Gerais and São Paulo, including in the last-named state, although Partridge (1956) insisted that these were likely to have all emanated from Minas Gerais. Another specimen (in the Museu de Zoologia de São Paulo), until recently often reported in the literature as originating from São Paulo state (e.g. Pinto 1938), collected at either Salto Grande or Fazenda Caiuá, on the rio Paranapanema, on 30 May 1903, is in fact from Paraná (Straube *et al.* 2002).

We made the third record of *M. octosetaceus* in São Paulo, almost two centuries after the specimens mentioned above. On 25 August 2017, a male was photographed (Fig. 1)



Figure 3. Study area of Padre Dória, Serra do Mar State Park, Salesópolis, São Paulo, Brazil.

on the rio Claro, in the Alto Tietê basin, Salesópolis. This river lies within a system of dams established in 1973 that provides water to 1.1 million people in the metropolitan region of São Paulo (SABESP 2017). The location is included within Important Bird Area SP04 (Bencke *et al.* 2006), forms part of Serra do Mar State Park (Fig. 3) and is very close to Boracéia Biological Station, whose avifauna has been studied for >60 years, and in this respect is one of the best-known locations in south-east Brazil (Cavarzere *et al.* 2010). It is also c.110 km from São Paulo city, the most populous urban area in Latin America, while the town of Salesópolis is well known among ornithologists for the recently described São Paulo Antwren *Formicivora paludicola* (Buzzetti *et al.* 2013).

The bird was seen swimming and diving next to the riverbank, in a 10-m stretch of slow-moving river, with clear, cold, very shallow (c.1 m deep) water, and a sandy bottom (Fig. 2). The bird was seen again, but not photographed, in the same place on 23 November 2017, perhaps indicating that it was territorial, although there have been no subsequent records despite regular visits to the area.

As previously mentioned, changes in hydrology caused by damming of rivers represents a significant threat to this globally endangered species. The rio Claro is a tributary of the rio Tietê, and the nearest dam is the Ribeirão do Campo, on the rio do Campo, which flows into the rio Claro. Nevertheless, upstream of the dam, the rio Claro extends for many kilometres through the Serra do Mar State Park, making it easier for the bird to move locally. Further work is needed to determine whether there is a population in this area, or this individual was a wanderer. But, for now it appears that Brazilian Merganser is not regionally extinct.

Acknowledgements

We especially thank the management and staff of Serra do Mar State Park, Núcleo Padre Dória, for logistical support, Jamile El Kassis Marangoni and Marcos Salvador Mathias Júnior for translating the text, researchers at the “Instituto Florestal”, Marina Mitsue Kanashiro for the map, and Alexsander Z. Antunes for suggestions concerning the manuscript. The manager of the North Shore Forestall Foundation, Leandro de Oliveira Caetano, and Larissa Araújo Nema encouraged us to publish this observation.

References:

- Bencke, G. A., Maurício, G. N., Develey, P. F. & Goerck, J. M. (eds.) 2006. *Áreas importantes para a conservação de aves no Brasil*, vol. 1. SAVE Brasil, São Paulo.
- Bianchi, C. A., Brant, S., Brandão, R. A. & Brito, B. F. 2005. New records of Brazilian Merganser *Mergus octosetaceus* in the rio das Pedras, Chapada dos Veadeiros, Brazil. *Cotinga* 24: 72–74.
- BirdLife International. 2018. Species factsheet: Brazilian Merganser *Mergus octosetaceus*. <http://datazone.birdlife.org/species/factsheet/brazilian-merganser-mergus-octosetaceus> (accessed 14 January 2018).
- Braz, V. S., Abreu, T. L. S., Lopes, L. E., Leite, L. O., França, F. G. R., Vasconcellos, M. M. & Balbino, S. F. 2003. Brazilian Merganser *Mergus octosetaceus* discovered in Jalapão State Park, Tocantins, Brazil. *Cotinga* 20: 68–71.
- Buzzetti, D. R. C., Belmonte-Lopes, R., Reinert, B. L., Silveira, L. F. & Bornschein, M. R. 2013. A new species of *Formicivora* Swainson, 1824 (Thamnophilidae) from the state of São Paulo, Brazil. *Rev. Bras. Orn.* 21: 269–291.
- Carboneras, C., Kirwan, G. M. & Sharpe, C. J. 2018. Brazilian Merganser (*Mergus octosetaceus*). In del Hoyo, J., Elliott, A., Sargatal, J., Christie, D. A. & de Juana, E. (eds.) *Handbook of the birds of the world Alive*. Lynx Edicions, Barcelona (retrieved from <https://www.hbw.com/node/52928> on 25 April 2018).
- Cavazere, V., Moraes, G. P. & Silveira, L. F. 2010. Boracéia Biological Station: an ornithological review. *Pap. Avuls. Zool., São Paulo*. 50: 189–201.
- Hughes, B., Dugger, B., Cunha, H. J., Lamas, I., Goerck, J., Lins, L., Silveira, L. F., Andrade, R., Bruno, S. F., Rigueira, S. & Barros, Y. M. 2006. *Action plan for the conservation of the Brazilian Merganser Mergus octosetaceus*. ICMBio, Brasília.
- Lamas, I. R. 2006. Census of Brazilian Merganser *Mergus octosetaceus* in the region of Serra da Canastra National Park, Brazil, with discussion of its threats and conservation. *Bird Conserv. Intern.* 16: 145–154.
- Lamas, I. R. & Lins, L. V. 2009. Brazilian Merganser (*Mergus octosetaceus*). In Schulenberg, T. S. (ed.) *Neotropical birds online*. Cornell Lab of Ornithology, Ithaca, NY. <https://doi.org/10.2173/nb.bramer1.01> (accessed 13 January 2018).
- Partridge, W. H. 1956. Notes on the Brazilian Merganser in Argentina. *Auk* 73: 473–488.
- Pinto, O. M. O. 1938. Catálogo das aves do Brasil e lista dos exemplares que as representam no Museu Paulista, 1a. parte. *Rev. Mus. Paulista* 22: 1–556.
- Ribeiro, F., Lins, L. V., Gomes, V. M., Nery, F. H. & Reis, E. S. 2011. Dispersão e maturidade sexual de *Mergus octosetaceus* Vieillot, 1817 na região da Serra da Canastra, Minas Gerais, Brasil. *Rev. Bras. Orn.* 19: 391–397.
- SABESP (Companhia de Saneamento Básico do Estado de São Paulo). 2017. Acesso em 13 de Janeiro de 2018, disponível em SABESP: <http://site.sabesp.com.br/site/imprensa/noticias-detalle.aspx?secaoId=66&id=7437>.
- Sick, H. 1997. *Ornitologia brasileira*. Ed. Nova Fronteira, Rio de Janeiro.
- Straube, F. C., Willis, E. O. & Oniki, Y. 2002. Aves colecionadas na localidade de Fazenda Caiuá (Paraná, Brasil) por Adolph Hempel, com discussão sobre sua localização exata. *Ararajuba* 10: 167–172.
- Stresemann, E. 1954. Ausgestorbene trod Aussterbende Vogelarten, vertreten im Zoologischen Museum zu Berlin. *Mitt. Zool. Mus. Berlin* 30: 38–53.
- Addresses:* Miguel Nema Neto, Rua Horto Florestal 1200, Rio do Ouro, Caraguatatuba, São Paulo, Brazil, CEP 11675-730, e-mail: miguelpsm@gmail.com. Fabiana Dias Pereira, Rodovia Alfredo Rolim de Moura 882, Vila Bueno, Salesópolis, São Paulo, Brazil, CEP 08970-000, e-mail: tiafabi84@hotmail.com. Ana Lúcia Morais Wuo, Rua 15 de Novembro 569, Centro, Salesópolis, São Paulo, Brazil, CEP 08970-000, e-mail: analuciauwuo@gmail.com. Carlos Zacchi Neto, Rua Horto Florestal 1200, Rio do Ouro, Caraguatatuba, São Paulo, Brazil, CEP 11675-730, e-mail: carlospsm@gmail.com

