



First records of Swallow Tanager *Tersina viridis* in Uruguay

Authors: Castelli, Diego, Muñoz, Joaquín, and Azpiroz, Adrián B.

Source: Bulletin of the British Ornithologists' Club, 142(4) : 487-490

Published By: British Ornithologists' Club

URL: <https://doi.org/10.25226/bboc.v142i4.2022.a8>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First records of Swallow Tanager *Tersina viridis* in Uruguay

by Diego Castelli, Joaquín Muñoz & Adrián B. Azpiroz

Received 29 June 2022; revised 25 September 2022; published 7 December 2022

<http://zoobank.org/urn:lsid:zoobank.org:pub:4571E64B-4537-43AE-BC06-FB446C3FAAE8>

SUMMARY.—The first records of Swallow Tanager *Tersina viridis* in Uruguay were made in 2012 and 2022 in different regions of the country. Together with other observations in central-south South America outside its usual distribution, these data appear to support the hypothesis that the species makes erratic movements in search of food during the austral winter non-breeding period. Alternatively, the new records could reflect an expansion of the species' distribution.

Swallow Tanager *Tersina viridis* is a member of the Thraupidae. It is both insectivorous and frugivorous, with the proportion of these dietary components varying seasonally (Schaefer 1953, Hilty 2020). The species is widely distributed in the Neotropics, from southern Panama to north-east Argentina (Misiones) and southern Brazil (Ridgely & Tudor 2009; Fig. 1). Migratory movements have been documented (Schaefer 1953, Schulenberg *et al.* 2007) and presence in the south of its distribution appears seasonal, with the species moving north in the austral winter (Hilty 2020). Some authors consider *Tersina* to be nomadic because, outside the breeding season, it may make irregular and erratic movements (Zelaya *et al.* 2013, Dardanelli *et al.* 2020, Hilty 2020), with exceptional records of vagrancy to the Cayman Islands and Bonaire (Prins *et al.* 2009, Kirwan *et al.* 2019) as well as in the south of its range. In Chile there are just two records, in Antofagasta and Valparaiso (Barros & la Red de Observadores 2016, 2021). In Argentina vagrants have been recorded in the provinces of Buenos Aires (Bremer & Bremer 1987, Narosky & Di Giacomo 1993, Bodrati & Bodrati 2017), La Pampa (Darrieu 1994), Salta (Di Giacomo *et al.* 1995), Tucumán (Ortiz 2009), San Juan (Lucero 2013), Córdoba, Formosa (Zelaya *et al.* 2013), Entre Ríos (Dardanelli *et al.* 2020), and most recently in Jujuy in 2021 (L. Fernández, <https://ebird.org/ebird/view/checklist/S87846418>) and Mendoza in 2022 (A. Zarco & Q. Vandemoortele, <https://ebird.org/checklist/S115612948>). In southern Brazil there are also records outside its main range. In July 2008 the species was observed for the first time in Pelotas, Rio Grande do Sul (P. R. Santos, <http://www.wikiaves.com/1343704>). During May–August 2021, several observations were made in the same municipality, with a max. of 30–40 individuals (eBird 2022; F. Jacobs, <http://www.wikiaves.com/4380737>). Some were feeding on the fruits of *Myrsine* sp., as well as *Tripodanthus* sp. and *Schinus terebinthifolius* (F. Jacobs pers. comm.).

In Uruguay there are no published records of Swallow Tanager. However, an oral communication made on 20–21 October 2012 (II Jornadas sobre 'Investigación y Manejo de Fauna Silvestre', Córdoba, Argentina) reported the first record of the species in the country (J. Villalba Macías pers. comm.). In addition, a few records have been reported in a national ornithology forum ('Ornitologiauy'). Below we provide details of several documented records, plus other recent reports, in Uruguay.

On 9 June 2012, E. Elgue observed an immature male Swallow Tanager (blotched green and blue; Fig. 2A) at Las Grutas, dpto. Maldonado (34°54'12"S, 55°02'10"W; Fig. 1). It was found in a small native forest feeding on fruits of *Myrsine laetevirens* (Primulaceae). The observation lasted a few minutes and the bird was very tame.




Figure 1. Distribution of Swallow Tanager *Tersina viridis* (based on Hilty 2020). Black dots indicate the records reported herein for Uruguay: (1) Las Grutas, dpto. Maldonado, 9 June 2012; (2) M' Bopicuá Biopark, dpto. Río Negro, 1 October 2012; (3) Botanic Garden, dpto. Montevideo, 20 April 2022; (4) Costa Azul, dpto. Rocha, 14 May 2022.

On 1 October 2012, J. Villalba Macías photographed an immature male (Fig. 2B) at M' Bopicuá Biopark, near the Uruguay River, dpto. Río Negro ($33^{\circ}06'48''\text{S}$, $58^{\circ}12'21''\text{W}$; Fig. 1). It was observed on a *Ficus luschnathiana* (Moraceae) and a *Prosopis nigra* (Fabaceae), and remained at the site for at least six days.

On 20 April 2022, L. Piñeyrúa, C. Croce & E. Muñoz observed a female-plumaged individual (Fig. 2C) at the Botanic Garden, dpto. Montevideo ($34^{\circ}51'33''\text{S}$, $56^{\circ}12'08''\text{W}$; Fig. 1). It was found in a sector of native flora with some fruiting trees that are often visited by other Thraupidae and *Turdus* spp.

On 14 May 2022 JM observed a female-plumaged individual (Fig. 2D) in a patch of psammophilous forest at Costa Azul, dpto. Rocha ($34^{\circ}37'45''\text{S}$, $54^{\circ}09'32''\text{W}$; Fig. 1). It was seen for four minutes atop a *Myrsine laetevirens*, feeding on its fruits and vocalising.

There are other undocumented records. In winter 2010, a female-plumaged bird was observed at Barrio Prado, dpto. Montevideo (J. Mazzula pers. comm.) and in 2012, another female-plumaged individual was seen at Santa Teresa, dpto. Rocha (M. Rodríguez-Cajaraville pers. comm.).

The observations detailed here represent the first records of Swallow Tanager in Uruguay. The species seems to be an occasional visitor attracted to fruiting trees during the austral autumn and winter. The records in Uruguay appear to support the hypothesis that it makes erratic movements in search of food, mainly fruit (Zelaya *et al.* 2013, Dardanelli *et al.* 2020, Hilty 2020). More than half of the records involved individuals taking fruits,




Figure 2. Swallow Tanager *Tersina viridis* records in Uruguay. (A) Immature male, Las Grutas, dpto. Maldonado, 9 June 2012 (E. Elgue). (B) Immature male, M'Bopicuá Biopark, dpto. Río Negro, 1 October 2012 (J. Villalba Macías). (C) Female-plumaged individual, Botanic Garden, dpto. Montevideo, 20 April 2022 (L. Piñeyrúa). (D) Female-plumaged individual, Costa Azul, dpto. Rocha, 14 May 2022 (J. Muñoz)

especially *Myrsine laetevirens*. Alternatively, the records could reflect an expansion of the species' range. The recent establishment of populations of numerous 'subtropical' bird species in Uruguay supports this hypothesis (Azpiroz *et al.* 2012, Menéndez 2020, Campaña *et al.* 2021), although more data are needed.

Acknowledgements

We are grateful to Ernesto Elgue, Juan Villalba Macías, Luis Piñeyrúa, Carlos Croce and Eduardo Muñoz who kindly provided their records. We thank Fernando Jacobs for information on the species in Rio Grande do Sul, Brazil; Ariel de León Mackey for comments on the text; and Guy Kirwan and Chris Sharpe for their valuable comments and helpful suggestions that improved the submitted manuscript.

References:

- Azpiroz, A. B., Menéndez, J. L., Jaramillo, A., Presa, D., Calimares, C., Saralegui, A. & Abente, J. S. 2012. New information on the distribution and status of birds in Uruguay. *Bull. Brit. Orn. Cl.* 132: 46–54.
- Barros, R. & la Red de Observadores de Aves. 2016. Resumen de avistamientos. Enero – diciembre 2015. *La Chiricoca* 21: 35–53.
- Barros, R. & la Red de Observadores de Aves. 2021. Resumen de avistamientos. Enero – diciembre 2020. *La Chiricoca* 27: 35–53.
- Bodrati, A. & Bodrati, G. 2017. Dos observaciones de Tersina o Golondrina naranjera (*Tersina viridis*) en el norte del conurbano bonaerense. *Nuestras Aves* 62: 66–67.
- Bremer, E. & Bremer, P. 1987. En los alrededores de Punta Rasa. *Nuestras Aves* 5: 21.
- Campaña, G., Rabau, T. & Velli, R. 2021. Primeros registros de Picaflor Tijereta *Eupetomena macroura* (Gmelin, 1788) en Uruguay. *Achará, Segunda época* 8: 8–15.

- Dardanelli, S., Reales, F., Ali, S. & Bisheimer, J. 2020. Primer registro de Tersina (*Tersina viridis*) para la provincia de Entre Ríos, Argentina. *Nuestras Aves* 65: 88–89.
- Darrieu, C. 1994. Dos nuevos registros para la avifauna pampeana *Tersina viridis viridis* (Emberizidae: Thraupidae) y *Columbina talpacoti talpacoti* (Columbidae). *Neotropica* 40: 56.
- Di Giacomo, A. G., Di Giacomo, A. S., López-Lanús, B. & Caradona, A. 1995. Nuevos registros de aves en el noroeste argentino. *El Hornero* 14: 67–68.
- eBird. 2022. eBird: an online database of bird distribution and abundance. Cornell Lab of Ornithology, Ithaca, NY. <http://www.ebird.org> (accessed 20 June 2022).
- Hilty, S. 2020. Swallow Tanager (*Tersina viridis*), version 1.0. In del Hoyo, J., Elliott, A., Sargatal, J., Christie, D. A. & de Juana E. (eds.) *Birds of the world*. Cornell Lab of Ornithology, Ithaca, NY. <https://doi.org/10.2173/bow.swatan1.01> (accessed 20 June 2022).
- Jacobs, F. & Fenalti, P. 2020. *Guia de identificação: aves do Rio Grande do Sul*. Ed. Aratinga, Pelotas.
- Kirwan, G. M., Levesque, A., Oberle, M. & Sharpe, C. J. 2019. *Birds of the West Indies*. Lynx Edicions, Barcelona.
- Lucero, F. 2013. Nuevas aves, primeras evidencias y localidades para las provincias de San Juan y Catamarca, Argentina. *EcoRegistros Rev.* 3: 52–63.
- Menéndez, J. 2020. Primer registro de mosqueta corona oliva *Phyllomyias virescens* (Temminck, 1824) para Uruguay. *Achará, Segunda época* 7: 1–3.
- Narosky, T. & Di Giacomo, A. D. 1993. *Las aves de la provincia de Buenos Aires: distribución y estatus*. Asociación Ornitológica del Plata & Ed. Vásquez Mazzini, Buenos Aires.
- Ortiz, D. 2009. Un nuevo registro del Tersina o Azulejo Golondrina (*Tersina viridis viridis*) para el noroeste argentino. *Nótulas Faunísticas (Segunda Ser.)* 38: 1–3.
- Prins, T. G., Reuter, J. H. H., Debrot, A. O., Wattle, J. & Nijman, V. 2009. Checklist of the birds of Aruba, Curaçao and Bonaire, south Caribbean. *Ardea* 97: 137–268.
- Ridgely, R. S. & Tudor, G. 2009. *Field guide to the songbirds of South America: the passerines*. Univ. of Texas Press, Austin.
- Schaefer, E. 1953. Contribution to the life history of the Swallow-Tanager. *Auk* 70: 403–470.
- Schulenberg, T. S., Stotz, D. F., Lane, D. F., O'Neill, J. P. & Parker, T. A. 2007. *Birds of Peru*. Revised and updated edn. Princeton Univ. Press.
- Zelaya, P. V., Salvador, S. A., Giraudo, A. & Klavins, J. 2013. Primeros registros de Tersina (*Tersina viridis*) para la ecorregión chaqueña de Argentina. *Nuestras Aves* 58: 52–54.
- Addresses:* Diego Castelli & Joaquín Muñoz, Facultad de Ciencias, Universidad de la República, Uruguay, Iguá 4225, Montevideo, Uruguay, e-mail: diegocastelli.22@gmail.com. Adrián B. Azpiroz, Departamento de Biodiversidad y Genética, Instituto de Investigaciones Biológicas Clemente Estable, Av. Italia 3318, 11600 Montevideo, Uruguay.