

Behavioural Responses of GPS-Collared Female Red Deer Cervus elaphus to Driven Hunts

Authors: Sunde, Peter, Olesen, Carsten R., Madsen, Torben L., and Haugaard, Lars

Source: Wildlife Biology, 15(4) : 454-460

Published By: Nordic Board for Wildlife Research

URL: https://doi.org/10.2981/09-012

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Wildl. Biol. 15: 454-460 (2009) DOI: 10.2981/09-012 © Wildlife Biology, NKV www.wildlifebiology.com

Behavioural responses of GPS-collared female red deer Cervus elaphus to driven hunts

Peter Sunde, Carsten R. Olesen, Torben L. Madsen & Lars Haugaard

Precise knowledge of how game species react to different hunting practices is a prerequisite for sound management of intensively hunted populations. We compared behavioural and spatial behaviour of five GPS-collared female red deer Cervus elaphus in Denmark before, during and after exposure to 21 driven hunts (2-5 times each). In 53% of all hunts, deer left their normal home ranges within 24 hours, moving on average 4 km and remaining away for an average of six days. Compared to pre-hunt values, deer moved longer distances per unit time on the day of the hunt and during the following two nights. Diurnal activity (based on motion sensors) did not increase significantly on the hunting day, but was lower than normal the day after the hunt. Nocturnal activity was equal before and after hunts. Deer spent 96% of their time in (safer) forest habitats by day and 43% by night before and after hunts. No induced responses were conditional on distance to the hunters (0-1.5 km), hunt duration (1.3-6.4 hours) or the time elapsed since previous hunts (4 to $>$ 30 days). The inclination of deer to flee from areas following hunts might complicate attempts to optimise harvesting policies in landscapes with many landowners within a typical flight range.

Key words: activity, Cervus elaphus, disturbance, habitat use, home range, movements, spatial behaviour

Peter Sunde, Carsten R. Olesen & Lars Haugaard, Aarhus University, National Environmental Research Institute, Department of Wildlife Ecology and Biodiversity, Grenaavej 14, DK-8410 Rønde, Denmark - e-mail addresses: psu@dmu.dk (Peter Sunde); cro@dmu.dk (Carsten R. Olesen); laha@dmu.dk (Lars Haugaard) Torben L. Madsen, St. Hjøllund Plantage, Faurholtvej 12, DK-7362 Hampen, Denmark - e-mail: tlmvrads@post.tele.dk

Corresponding author: Peter Sunde

Received 3 February 2009, accepted 18 May 2009

Associate Editor: Leif Egil Loe

Human activities in general, and hunting actions in particular, often have significant impacts on behaviour and spatial distribution of wildlife through the anti-predator behaviours they evoke (Frid & Dill 2002). This applies particularly to large-bodied, long-lived species with a long evolutionary history of human persecution such as the red deer Cervus elaphus (e.g. Jeppesen 1987a, Cole et al. 1997, Burcham et al. 1999, Conner et al. 2001, Vieira et al. 2003). Managers of intensively hunted deer populations should therefore not only address the impacts of harvest pressure, but also disturbance-mediated impactsonindividuals andpopulations (e.g.Phillips & Alldredge 2000, Jayakody et al. 2008).

Spatial reactions of red deer to recreational activities and hunting have mainly been studied in North American landscapes with continuous forests (e.g. Cole et al. 1997, Burcham et al. 1999, Conner et al. 2001,Vieiraetal.2003).Lessinformationexists from the fragmented forestlandscapes ofWesternEurope (Jeppesen 1987a, Jayakody et al. 2008) where escape possibilities are constrained by agricultural areas. Studies show that red deer react to hunters by leaving disturbed areas (Jeppesen 1987a, Cole et al. 1997, Burcham et al. 1999, Conner et al. 2001, Vieira et al. 2003), but little is known about the short- and longterm impacts on movements, activity and habitat use.

In this paper, we quantify and test changes in spatial behaviour, habitat use and activity levels of female red deer taggedwithGPS- and activity-loggers, and exposed to driven hunts within their home

WILDLIFE BIOLOGY 15:4 (2009)

	Age in years	With calf at					No. of
					95% kernel in		hunting
Deer	at hunting	hunting?	Survey period	Reason for termination	ha (Sep.-Oct.)	Data	events
A1	$1^{1/2}$	N ₀	6 Mar.07 - 14 Dec.07	Shot/wounded on 15 Dec.07	415	P.A	
A ₂	$2^{1/2}$	N ₀	15 Mar.06 - 26 Mar.07	Data storage filled	711	P.A	
A ₃	\geq 3 ¹ / ₂	Yes	15 Mar.06 - 23 Jan.07	Shot on 24 Jan.07	287	P.A	
A ₅	$2^{1/2}$	Yes	12 Apr.06 - 21 Mar.07	Data storage filled	330	P.A	
A7		No.	6 Mar.07 - 20 Jan.08	Data storage filled	360	D	4

Table 1. Data on the five female red deer exposed to hunting events. P indicates positioning data based on hourly GPS positions, and A indicates activity data based on movement sensor information logged every five minutes.

ranges. Danish red deer should be a good subject for studies of human disturbance effects on heavily exploited deer populations, as man has been the sole population regulating agent for >200 years. The hunting season extends from November to January for hinds and calves, and from September to January for stags. Owners of properties >1 ha and renters of properties > 5 ha are free to shoot as many deer as they like from sunrise to sunset.Danishlandscapes with forests and plantations (which provide cover during daylight; Mysterud & Østbye 1999) surrounded by agriculturallandscapes (used for foraging at night), also provide an excellent opportunity to study disturbance responses of deer whose spatial escape possibilities are constrained by fragmentation of protective vegetation cover.

Material and methods

Red deer were captured, and all driven hunts took place in the privately-owned plantation of 'St. Hjøllund Plantage', which covers 14.0 km² (98% conifers, consisting mainly of Norway spruce Picea *abies*) in central Jutland, Denmark (56 \degree 05'N, 9 \degree 25'E). St. Hjøllund Plantage is situated in a lowland (0-170 m a.s.l.) region with amaritime climate (average annual precipitation for mid and western Jutland, 1960-90: 781 mm spread over 131 days; monthly mean temperatures ranging from -0.2 in January to 15.4° C in July) and sandy soils. The numbers of reddeerwithin theplantationvaried seasonally from ca 100 (in March) to ca 200 (in September-October; i.e. $7-14 \text{ km}^{-2}$ of which 20-24 are shot annually. Herds appear to be diffusely organised into groups of usually 3-8 and occasionally 80-90 red deer. St. Hjøllund Plantage is surrounded by agricultural land, farms, villages and coniferous plantations (0.1-30 km2). During our two-year study, driven hunts in St. Hjøllund Plantage were carried out in the same way andintensity asin all other years (i.e. 5- 6 hunts per season, seeTable 2). In addition to driven

- WILDLIFE BIOLOGY 15:4 (2009) 455

deer hunts, 1-2 small-game hunts per year were the only other disturbance activities which occurred in the plantation during the hunting season. Although the plantation is open to foot-access by the general public, visitors were rarely encountered. No information was available on hunting pressure on neighbouring properties, although deer were intensively hunted everywhere in the neighbourhood throughout the hunting season.

During March-April, we captured five female red deer (Table 1) which maintained home ranges within the St. Hjøllund estate during the following hunting season (November-January). We caught the deer in 300 m^2 traps established in the plantation which were baited with sugar beet Beta vulgaris during late winter. After being immobilised by a mixture of the drugs Etorfine and Rompun (Diprenofine and Antisedanwere used as antidotes), all deerwere equipped with a LotekWildcell GPS-GSM 4400M collar with a storage capacity of 7,000 positions, registered at minimum intervals of one hour and automatically transferred to the base station viaSMS.This enabled us to register hourly positions for most of the deer throughout the following hunting season. The mean fix success rate of the GPS collars was 94% (range: 92-96%). We retrieved activity information from motion sensors (arbitrary value ranging from 0 (immobile) to 100 (maximum activity), logged every 5th minute) from four deer after collar recovery (see Table 1).

Quantification of deer behaviour and habitat use

Hourly behavioural and spatial parameters were averaged for the diurnal (sun angle $>0^{\circ}$) and nocturnal (sun angle $\langle 0^\circ \rangle$ phases of each day for each individual deer. Response variables were quantified as 1) mean hourly displacement distances, 2) mean motion score (activity censor values), and 3)whether allfixeswerelocatedoutside anindividualdeer'sprehunting (September-October) home range and (if being outside for an entire day or night) 4) its maximum distance to the nearest home-range border.

The pre-hunting home range was defined as a 95% fixed kernel isopleth calculated with the Animal Movement extension for ArcView 3.2 (Hooge & Eichenlaub 1997) using the ad hoc option to select smoothing parameter (H). Each home range was based on ca $24 \times 60 = 1,440$ locations, sampled with 1-hour intervals throughout September-October. Finally, we measured 5) habitat use, defined as the proportion of time spent in forest vegetation (in this area almost entirely spruce) vs other habitats as measured from the digital image TOP10DK.

Quantification of hunting disturbances

We considered a deer to be potentially exposed to a hunting event if it was located within a distance of 1.5 km of a hunted area. All hunting took place within the plantation where deer remained during daytime. Organised hunting sessions (Table 2) were registered as the mapped zones along which hunters were posted and in which beaters and dogs systematically searched the area to drive out deer towards a line of hunters. As a rule, 2-4 areas covering 50- 300 ha each were searched from dawn (09:00-10:00) to 1-hour before dusk (15:00). Movements of beaters and dogs were registered using GPS-loggers. Positions of the surveyed deer in relation to the hunting activities were obtained from hourlyGPS-positions.

The type and strength of the stress stimulus on individuals during a given hunting session (see Table 2) was quantified as 1) the duration of the total hunting period, 2) the minimum recorded distance of an individual deer to a hunting zone, 3) the type of huntingpractice (traditionaldrivenhuntswithlarge, free-ranging dogs vs a presumably more gentle practice using smaller dogs called 'motion hunts'), and 4) the (log-transformed) number of days which had passed since the previous hunting disturbance (see Table 2).

Statistical analyses

Because deer in diurnally hunted populations behave differently during daylight and darkness hours (being less active in daylight), we analysed day and night values separately.We used general linear mixed models (PROCMIXEDin SAS 9.1) and generalised linear mixed models (PROC GLIMMIX in SAS 9.1)

Table 2. The 10 hunts (A-J; where type D indicates traditional driven hunts, and M motion hunts) and 21 deer-specific hunting events with information on the deer's shortest distance (in m) to the hunting zone (DIST), days elapsed since the deer experienced the previous hunt (DSPH), and duration of the survey period in days before and after the actual hunt (lasting until the next hunt was initiated or termination of data collection). Spatial reactions to hunts are categorised as 'S': stayed (located in the home range the day after the hunt), 'L-R': left home range within one day, returning during the survey period, 'L-C': left range within one day without returning within the survey period ('censored'). 'S^{*'} indicates a case in which a deer stayed within its home range until the second day after the hunt, and was shot the day after. MaxDIST indicates the maximum distance (in km) of a deer from its home range during the pre-hunt period and the day after the hunt.

 456 \circ

WILDLIFE BIOLOGY 15:4 (2009)

to contrast the deers' reactions (spatially, behaviourally and concerning habitat choice) to hunting with the pre-hunt baseline values (last three days before hunts) across each individual hunting experience (hereafter 'deer-specific hunting events'; see Table 1). Time stage (categorised as 'before hunts' and day 0, 1, 2, .. post-hunt) was entered as a fixed class variable. As experimental blocking units, we entered deer identity, the deer-by-time stage interactionanddeer-specifichuntingeventsnestedwithin deer as random effects. Variance heteroscedasticity between time stages was included in all models. We used least-square means tests to test for differences between the pre-hunt situation and the later stages.

Binary response variables (presence/absence in the home range during an entire diurnal ornocturnal phase or whether a location was situated inside or outside forest) were modelled with a logit link functioninGLIMMIX,estimatingvariancecomponents and denominator degrees of freedom with Saittertwaitte's approximation (Littell et al. 2006), adjusting for over-dispersion ("Random residual;" statement).

Normally distributed response variables (logtransformed if necessary, including distance from home range of migrating deer, mean hourly movement distance, and mean activity score) were modelled in MIXED, estimating variance components and denominator degrees of freedom with Kenward-Roger's approximation (Littell et al. 2006).

Durations of extra home-range excursions (see Table2)weremodelledusingKaplan-Meier analysis (PROC LIFETEST in SAS, setting 'survival time'= duration of excursion to 0 for non-migrating deer), as this approach enabled proper handling of censoredcases (excursionslastingbeyond thedateof the next hunt or termination of the survey period: see Tables 1 and 2).

Results

Spatial reactions to hunting

After 53% of all hunts, female deer left their home range during the subsequent night, being significantly more likely to be away from their home range compared to the pre-hunt baseline for up to a week (Fig. 1A). Those deer that left their home range stayed away for an average of 4.7 days $(SE=1.04)$ and a maximum of 9.5 days (see Table 2). Despite individual variation (log-rank test of equality over strata: $\chi^2_4 = 9.83$, P = 0.043), responses were not con-

© WILDLIFE BIOLOGY 15:4 (2009) 457

Figure 1. Spatial reactions of GPS-collared female red deer to hunting events carried out during the diurnal period of day 0 measured as the probability of deer migrating out their home range, for hinds which migrate (A), and the harmonic mean migration distance (in m) from the nearest home-range border (B). Least square mean estimates (95% CI) for each day (with separate analyses for diurnal and nocturnal periods) are based on generalised linear mixed models, accounting for variation across individuals and hunting events, incorporating variance heteroscedasticity between time intervals. Significant deviations from pre-hunt estimates are shown as $\cdot P < 0.05$, $\cdot\cdot P < 0.01$ and $\cdot\cdot\cdot$. $P < 0.001$.

ditional on year, type or duration of hunt, the initial distance to the hunting zone nor the time elapsed since the previous hunt (log-rank tests: all $Ps > 0.5$). Deer leaving their normal home range after a hunting event migrated, on average, 4 km (maximum: 17 km), which is significantly higher than the mean excursion distances before hunts (Fig. 1B).

Behavioural reactions to hunting

Motion sensor information showed that deer were considerably more active at night than by day (Fig. 2A). Diurnal activity was not significantly different on the day of the hunt compared to pre-hunt levels, but there was a significant drop on the day after the hunt (see Fig. 2A). Nightly activity levels were identical before and after hunts (see Fig. 2A).

Before hunts, deer moved 2.2 times longer distancesperhourduring thenight thanduringdaytime hours (Fig. 2B). Compared to the pre-hunt daylight baseline, deer increased their mean hourly displacement distances on the day of the hunt, returning to

Figure 2. Least square means (95% CI) of activity (A; based on motion sensors with daily means of 17 hunting events from four deer; log-transformed in analysis) and hourly displacement distances (B; with daily means from 21 hunting events from five deer using diurnal values log-transformed in analysis) of GPScollared female red deer before, during (diurnal period of day 0) and after the hunting events. Separate analyses were run for diurnal and nocturnal situations. Estimates are based on mixed models (see text), and significant differencesin least squaremeans from the before-hunt situation (last three days before a hunt) are marked as $*: P < 0.05$ and $***: P < 0.001$.

normal values the day after the hunt. Relative to the nocturnal pre-hunt baseline, deer increased their mean hourly displacement distances significantly on the first and second nights after hunts (see Fig. 2B).

Further statistical modelling revealed that the increased movement rates recorded during the first night following hunts were caused by individuals which moved out of their home range $(F_{1, 116}=31.62)$, $P < 0.0001$), whereas deer which remained in their home range did not move significantly more the first nightafter thehunt thanin theprecedingnights (least square mean comparison: $t_{55.6} = 0.49$, $P = 0.62$).

Activity responses on the day of the hunt and the following day were independent of the type of hunt, duration of hunting activities, the deer's shortest distance to the hunted area and the elapsed time since the preceding hunt (48 tests: all $Ps > 0.09$).

Time spent in forest cover before and after hunts

Prior to hunts, deer spent 96% (95% CI: 83-99) of their time in forest during the daytime, compared to

43% (27-61) at night. Deer spent equal amounts of time in forest before hunts, and the first and second day during and after hunting (day: $F_{2, 8.67}=0.33$, P=0.73; night: $F_{2,7,40}$ =0.13, P=0.88).

Discussion

Inabouthalfofallourcases, traditionaldrivenhunts and thelessintensivemotionhuntscaused female red deer located within a distance of 1.5 km from hunted areas to leave their home ranges the following night, and to stay away for an averageof sixdays.Our study clearly shows that deer postponed their emigration response until after dark, i.e. after hunters, dogs and beaters hadleft the plantation.Thus, the deer did not shift as an immediate flight response, but because they assessed the hunted area as being unsafe for the subsequent days. Even though hunting-induced migration responses varied amongst individuals, all marked deer left their range at least once, suggesting that migration was a general response-type to hunting disturbance in female red deer. This reaction pattern agrees well with anti-hunter responses of red deer previously reported from elsewhere in Europe (Jeppesen 1987a) and North America (Cole et al. 1997, Burcham et al. 1999, Conner et al. 2001, Vieira et al. 2003) and is basically similar to responses to natural predators which are also avoided on a scale above the home-range level (e.g. Anderson et al. 2005). Temporary avoidance of areas following human disturbance is also known from other groupliving cervids such as reindeer Rangifer tarandus (e.g. Reimers et al. 2003, Seip et al. 2007, Vistnes & Nellemann 2008) adapted to escape natural predators through migration (Fryxell & Sinclair 1988). By contrast, other European, forest-dwelling but less social cervids such as moose Alces alces (Neumann et al. 2009) or roe deer Capreolus capreolus (Jeppesen 1987b) do not appear to leave their home ranges as a delayed response to hunting activities.

Increased hourly displacement distances on the day of hunts and lowered diurnal motion indices on the day after hunts were the only detectable behavioural response to hunting apart from derived emigration responses (significantly increased hourly !displacement distances the first two nights after hunts). As increased hourly displacement distances of migrating individuals might be a mere result of straighter path lines rather than increased movement speeds, energy expenditure and foraging efficiency may therefore not necessarily have been

WILDLIFE BIOLOGY 15:4 (2009)

severely affected after hunts, although no definite conclusions can be drawn from activity and spatial data alone, as presented here.

Lack of change in usage of forest and non-forest vegetation after hunts also suggests that hunting activities did not severely distort basic foraging routines. This is probably because prior to hunts deer already showed anti-hunter avoidance responses, expressed by their habitat use during daylight, when they almost exclusively used forested habitats which deer are known to perceive as safer (Mysterud & Østbye 1999), as opposed to during the night when they spent less than half of their time in forests. Lack of response to hunting during daytime habitat use was therefore a trivial result of the fact that this antipredator response could not be heightened. Unaltered use of non-forest habitats after dark before, during and after hunts might reflect an adaptive response to the actual hazard situation, as no shot must be fired at deer between sundown and sunrise.

In highly cultivated landscapes with no natural predators and with multiple landowners within the radius of a hunting-induced migration distance, the observeddelayedmigratory reaction tohuntingmay not be adaptive if deer thereby expose themselves to new hunters. In our study this was exemplified by two of five hinds being shot during hunting-induced migrations to neighbouring estates. In Denmark at least, unsustainably high hunting pressure in areas with high landowner densities is the likely reason for red deer populations being much more aggregated than habitat composition would predict (Sunde et al. 2008). With the caveat that disturbance-induced migration responses might vary across habitats (Jayakody et al. 2008) and between countries with different hunting practices (see Milner et al. 2006), our results seem to suggest that even in areas where hinds are seasonally stationary, hunting should be coordinated within units with a spatial resolution of at least 10×10 km to avoid local overexploitation of hinds. As hunting activities also appeared to elicit emigration responses in deer located adjacent to the hunted areas, organisers of deer driven hunts should furthermore be aware that hunting activities might affect deer even when not directly encountered by beaters, dogs or hunters.

Acknowledgements - our grateful thanks go to the owners of St. Hjøllund Plantage who permitted us to study deer and hunter behaviour on their estate. Our study was financed by the Danish Forest and Nature Agency (Wildlife & Landscape Research Package - project 2B). We are also grateful for the thorough comments on thepaper from two

© WILDLIFE BIOLOGY 15:4 (2009) 459

anonymous reviewers and to Tony D. Fox for polishing the English language.

References

- Anderson, D.P., Turner, M.G., Forester, J.D., Zhu, J., Boyce, M.S., Beyer, H. & Stowel, L. 2005: Scale-Dependent Summer Resource Selection by Reintroduced Elk in Wisconsin, USA. - Journal of Wildlife Management 69: 298-310.
- Burcham, M., Edge, W.D. & Marcum, C.L. 1999: Elk use of private land refuges. - Wildlife Society Bulletin 27: 833-839.
- Cole, E.K., Pope,M.D. & Anthony, R.G. 1997: Effects of road management on movement and survival of Roosevelt elk. - Journal of Wildlife Management 61: 1115- 1126.
- Conner, M., White, G.C. & Freddy, D.J. 2001: Elk movement in response to early-season hunting in Northwest Colorado. - Journal of Wildlife Management 65: 926- 940.
- Frid, A. & Dill, L. 2002: Human-caused disturbance stimuli as a form of predation risk. - Ecology & Society 6(1): article 11. Available at: http://www.ecologyand society.org/vol6/iss1/art11/print.pdf (Last accessed on 14 September 2009).
- Fryxell, J.M. & Sinclair, A.R.E. 1988: Causes and consequences of migration of large herbivores. - Trends in Ecology and Evolution 3: 237-241.
- Hooge, P.N. & Eichenlaub, B. 1997: Animal movement extension to ArcView. ver 1.1. Alaska Biological Science Center, U.S. Geological Survey, Anchorage. Available at: http://www.absc.usgs.gov/glba/gistools/ animal_mvmt.htm (Last accessed on 14 September 2009).
- Jayakody, S., Sibbald, A.M., Gordon, I.J. & Lambin, X. 2008: Red deer Cervus elaphus vigilance behaviour differs with habitat type and human disturbance. -Wildlife Biology 14(1): 81-91.
- Jeppesen, J.L. 1987a: Impact of human disturbance on home range, movements and activity of red deer (Cervus elaphus) in a Danish environment. - Danish Review of Game Biology 13(2): 1-38.
- Jeppesen, J.L. 1987b: The disturbing effects of orienteering and hunting on roe deer (Capreolus capreolus) -Danish Review of Game Biology 13(3): 1-24.
- Littell, R.C., Milliken, G.A., Stroup, W.V., Wolfinger, R.D. & Shabenberger, O. 2006: SAS® for Mixed Models. 2nd edition. - Cary, NC, USA: SAS Institute Inc., 814 pp.
- Milner, J.M., Bonenfant, C., Mysterud, A., Gaillard, J.M., Csányi, S. & Stenseth, N.C. 2006: Temporal and spatial development of red deer harvesting in Europe: biological and cultural factors. - Journal of Applied Ecology 43: 721-734.
- Mysterud, A. & Østbye, E. 1999: Cover as a habitat element for temperate ungulates: effects on habitat selection and demography. - Wildlife Society Bulletin 27: 385-394.
- Neumann, W., Ericsson, G. & Dettki, H. 2009: The nonimpact of hunting on moose *Alces alces* movement, diurnal activity, and activity range. - European Journal of Wildlife Research 55: 255-265.
- Phillips, G.E. & Alldredge, A.W. 2000: Reproductive success of elk following disturbance by humans during calving. - Journal ofWildlifeManagement 64: 521-530.
- Reimers, E., Eftestøl, S. & Colman, J.E. 2003: Behavior responses of wild reindeer to direct provocation by a snowmobile or skier. - Journal of Wildlife Management 67: 747-754.
- Seip, D., Johnson, C.J. & Watts, G.S. 2007: Displacement of mountain caribou from winter habitat by snow-

mobiles. - Journal of Wildlife Management 71: 1539- 1544.

- Sunde,P.,Asferg,T.,Andersen,P.N.&Olesen,C.R. 2008: Hvor nedlægges krondyrene - og hvorfor? Betydningen af landskab, urbanisering og tidligere udbredelse for det lokale jagtudbytte af krondyr i Jylland i jagtsæsonen 2001/02. - NERI Technical report 690, 38 pp. (In Danish). Available at: http://www2.dmu.dk/Pub/ FR690.pdf (Last accessed on 14 September 2009).
- Vieira, M.E., Conner, M.M., White, G.C. & Freddy, D.J. 2003: Effects of archery hunter numbers and opening dates on elk movement. - Journal of Wildlife Management 67: 717-728.
- Vistnes, I. & Nellemann, C. 2008: The matter of spatial and temporal scales: a review of reindeer and caribou response to human activity. - Polar Biology 31: 399-407.