

## **Cassin's Hawk-Eagle *Spizaetus africanus* in Ndundulu Forest: a First Record for Tanzania, with Biogeographical Implications**

Author: Jones, Trevor

Source: Journal of East African Natural History, 96(2) : 187-192

Published By: Nature Kenya/East African Natural History Society

URL: [https://doi.org/10.2982/0012-8317\(2007\)96\[187:CHSAIN\]2.0.CO;2](https://doi.org/10.2982/0012-8317(2007)96[187:CHSAIN]2.0.CO;2)

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

**CASSIN'S HAWK-EAGLE *SPIZAETUS AFRICANUS* IN NDUUNDULU  
FOREST: A FIRST RECORD FOR TANZANIA, WITH  
BIOGEOGRAPHICAL IMPLICATIONS**

**Trevor Jones**

Environmental Sciences Research Centre, Department of Life Sciences  
Anglia Ruskin University, East Road, Cambridge, CB1 1PT, UK  
Udzungwa Ecological Monitoring Centre  
Box 99, Mang'ula, Morogoro, Tanzania  
tembomkubwa@gmail.com

**ABSTRACT**

A single adult Cassin's hawk-eagle *Spizaetus africanus* was sighted on five occasions over three years in a highland forest in the Udzungwa Mountains, the first ever record of this species in Tanzania. This discovery has potentially significant biogeographical implications, strengthening ancient links between the forests of the Udzungwa Mountains and the Guineo-Congolian forests of central Africa.

**Keywords:** Cassin's, hawk-eagle, *Spizaetus*, Nduundulu, Udzungwa Mountains.

Cassin's hawk-eagles are the only African representative of the tropical genus *Spizaetus* Vieillot, 1816, a group of small to medium sized eagles with short, rounded wings and long tails (Brown *et al.*, 1982; Ferguson-Lees & Christie, 2001; though for current taxonomic debate based on recent molecular analyses see Helbig *et al.*, 2005; Lerner & Mindell, 2005; Haring *et al.*, 2007). *Spizaetus africanus* Cassin, 1865 is a forest eagle with a black-and-white appearance (wing length, male 330–341 mm, female 381 mm; tail length, male 211–234 mm, female 266 mm) (Ferguson-Lees & Christie, 2001). The adult is mainly white below with black patches on the side of the breast, and blackish axillaries and underwing-coverts. Above it is mainly black, with black-tipped primaries. The tail is banded dark and light brown; cere and feet are pale yellow. It is known to be resident in dense tropical forests across Equatorial Africa, from Togo east to Gabon, Cameroon and the Democratic Republic of Congo. The easternmost known site for this species is the Mabira Forest Reserve of central Uganda (Fishpool & Evans, 2001), though a dead juvenile was found on Mount Elgon in western Kenya in 1926 (Clark & Edelstam, 2001) (figure 1).

A single adult Cassin's hawk-eagle was observed on five occasions between September 2004 and November 2007 (11 September 2004, 13 September 2004, 19 February 2006, 29 August 2006, 2 November 2007) in Nduundulu Forest (7°45'S, 36°30'E; 1300–2000 m) in the Udzungwa Mountains of southern Tanzania (figure 1). The plumage pattern and feathered tarsi of the observed and photographed bird (figure 2) are diagnostic. The amount of white on the underparts is consistent with Cassin's hawk-eagle and distinguishes it from African hawk-eagle *Aquila spilogaster* Bonaparte, 1850 (Helbig *et al.*, 2005; formerly *Hieraaetus*

*spilogaster*: Brown *et al.*, 1982; Ferguson-Lees & Christie, 2001) and Ayres' eagle *Hieraaetus ayresii* Gurney, 1862. The white feathered tarsi, and black underwing coverts observed in flight, distinguish it from the Black sparrowhawk *Accipiter melanoleucus* Smith, 1830. Colour and length of the tail distinguish it from the Augur buzzard *Buteo augur*




Figure 1. Map showing the currently known distribution of Cassin's hawk-eagle *Spizaetus africanus* in East Africa, including Ndundulu Forest, Tanzania. Mount Elgon, Kenya, from where a juvenile specimen was collected in 1926, is also shown.

Rüppell, 1836. Single birds were spotted on two occasions sitting in the tall upper canopy of the forest (which reaches over 50 m in height), and on three occasions soaring in circles low over the forest canopy. On two occasions a bird was heard calling while in flight, a short high-pitched call repeated several times, which was later recorded and compared with and noted to sound identical to an audio recording of Cassin's hawk-eagle. It is not possible to confirm whether all encounters have been with the same or different individuals.



Figure 2. An adult Cassin's hawk-eagle *Spizaetus africanus*, photographed in Ndundulu Forest, Tanzania, August 2006. Photograph by Anthony Jarrett.

Ndundulu and the contiguous Luhomero Forest (to 2500 m) cover a total area of 250 km<sup>2</sup> comprising closed canopy forest with a few open clearings, and is one of the richest and most biodiverse of all forests in Tanzania (Dinesen *et al.*, 2001; Burgess *et al.*, 2007). As an indication of its extraordinary richness across all taxa, it is home to 22 restricted-range bird species (Marshall *et al.*, 2001; T. Jones, unpubl. data) and at least three restricted-range primate species, including the recently discovered kipunji *Rungwecebus kipunji* Davenport *et al.*, 2006 (Jones *et al.*, 2005; Davenport *et al.*, 2006). There have also been discoveries in recent years of a new genus and species of bird (Dinesen *et al.*, 1994), a new species of giant sengi or elephant shrew (Rovero & Rathbun, 2006; Rovero *et al.*, in press), and a new species of shrew (Stanley *et al.*, 2005). The discovery of Cassin's hawk-eagle here further reinforces the perception of Ndundulu as an outstanding refuge for relic faunal populations.

All five sightings to date have been made within an area of 7 km<sup>2</sup> in southern Ndundulu, although several weeks of survey effort were completed during 2005–2006 in other parts of this forest (Jones, 2006). However, these surveys were primarily focused on primates and the apparent tendency of these birds to spend most of the time in or above the forest canopy makes them very difficult to detect (Friedmann & Williams, 1970). A failure of previous surveys to detect this species does not prove that it was absent (Dinesen *et al.*, 1993; Hunter *et al.*, 1996;

Butynski & Ehardt, 2003). Moreover, there are large areas of the surrounding forest which have not yet been surveyed. The minimum 250 km<sup>2</sup> of available potential habitat in Ndundulu/Luhomero raises the possibility of a small viable population of resident Cassin's hawk-eagles, and it is not inconceivable that they could also have gone undetected by researchers in neighbouring forests such as Nyumbanitu (49 km<sup>2</sup>) (e.g. Dinesen *et al.*, 1993, 2001; Marshall *et al.*, 2001). It is now important that further investigations focused on this species are carried out to determine distribution, abundance and breeding status in the Udzungwa Mountains.

The nearest reported site of Cassin's hawk-eagle to Ndundulu is in Mabira Forest Reserve, south-central Uganda, 900 km away (figure 1). The discovery of this bird in Ndundulu has interesting implications for the debate over the presence and timing of an ancient continuous forest between the Udzungwa Mountains (and other forested highlands of Tanzania) and the Guineo-Congolian forests of central and western Africa (Axelrod & Raven, 1978; White, 1981; Lovett, 1993a; Burgess *et al.*, 2007). Although the Tanzanian highland forests have been isolated for several million years, recent discoveries of organisms endemic to these mountains, but having Guineo-Congolian affinities, have argued for a more contiguous historical distribution of species, maybe as recently as 10 million years ago. For example, the Udzungwa-endemic gecko *Urocotyledon rasmusseni* Bauer & Menegon, 2006 is more closely related to *U. weileri* Müller, 1909 of West Africa than to its Tanzanian relative *U. wolterstorffi* Tornier, 1900 (Bauer & Menegon, 2006). There are butterflies (De Jong & Congdon, 1993), reptiles (Howell, 1993) and trees (Lovett, 1993b) in the Udzungwas with close Guineo-Congolian affinities, and the recent discovery of the shrew *Congosorex phillipsorum* in Ndundulu marked the first known record of a mammalian genus with disjunct species in these mountains and in central Africa (Stanley *et al.*, 2005). Among birds, molecular analyses of some isolated eastern Tanzanian montane subspecies have shown them to be divergent from relatives in the Central Congolian forests (Roy, 1997; Roy *et al.*, 1997; Beresford, 2003). Although birds are generally more mobile dispersers across different habitats than other vertebrates, the unexpected presence of the forest-dependent Cassin's hawk-eagle in Ndundulu, if it is indeed indicative of a relic population, lends further support to the theory of more contiguous forest between the Congo Basin and southern Tanzania than is found today.

## ACKNOWLEDGMENTS

I am grateful to Simon Thomsett, Munir Virani, Simon Mann, Neil Baker and other contributors to the Tanzania Birds email group for assisting with species identification; to Pete Leonard for providing an audio recording of Cassin's hawk-eagle; and to Bill Stanley and Rebecca Banasiak (Field Museum of Natural History, Chicago) for helping with the map. Thanks to my field assistant Athumani Mndeme and to many people from Udekwa who have helped in the forest. Special thanks to Anthony Jarrett for the photograph. All observations were made during fieldwork funded by the Critical Ecosystem Partnership Fund, Wildlife Conservation Society, Fauna and Flora International, and the Primate Society of Great Britain. Research permission was granted by the Tanzania Wildlife Research Institute, the Commission for Science and Technology, and the Tanzania Forestry and Beekeeping Division. The manuscript was improved by comments from Simon Thomsett and two anonymous reviewers.

## REFERENCES

- Axelrod, D.I. & P.H. Raven (1978). Late Cretaceous and Tertiary vegetation history of Africa. In M.J.A. Werger (ed.), *Biogeography and Ecology of Southern Africa*. Junk, The Hague. Pp. 77–130.
- Bauer, A.M. & M. Menegon (2006). A new species of Prehensile-tailed Gecko, *Urocotyledon* (Squamata: Gekkonidae), from the Udzungwa Mountains, Tanzania. *African Journal of Herpetology* **55**: 13–22.
- Beresford, P. (2003). Molecular systematics of *Alethe*, *Sheppardia* and some other African robins (Muscicapidae). *Ostrich* **74**: 58–73.
- Brown L.H., E.K. Urban & K. Newman (1982). *The Birds of Africa, Vol. I*. Academic Press, London.
- Burgess, N.D., T.M. Butynski, N.J. Cordeiro, N. Doggart, J. Fjeldså, K.M. Howell, F. Kilahama, S.P. Loader, J.C. Lovett, B. Mbilinyi, M. Menegon, D.C. Moyer, E. Nashanda, A. Perkin, F. Rovero, W.T. Stanley & S.N. Stuart (2007). The biological importance of the Eastern Arc Mountains of Tanzania and Kenya. *Biological Conservation* **134**: 209–231.
- Butynski, T.M. & C.L. Ehardt (2003). Notes on ten restricted-range birds in the Udzungwa Mountains, Tanzania. *Scopus* **23**: 12–27.
- Clark, W.S. & C. Edelstam. (2001). First record of Cassin's Hawk Eagle *Spizaetus africanus* for Kenya. *Bulletin of the African Bird Club* **8**: 138–139.
- Davenport, T.R.B., W.T. Stanley, E.J. Sargis, D.W. De Luca, N.E. Mpunga, S.J. Machaga & L.E. Olson (2006). A new genus of African monkey, *Rungwecebus*: morphology, ecology, and molecular phylogenetics. *Science* **312**: 1378–1381.
- De Jong, R. & T.C.E. Congdon (1993). The montane butterflies of the eastern Afrotropics. In J.C. Lovett & S.K. Wasser (eds), *Biogeography and Ecology of the Rain Forests of Eastern Africa*. Cambridge University Press, Cambridge. Pp. 133–172.
- Dinesen, L., T. Lehmberg, J.O. Svendsen & L.A. Hansen (1993). Range extension and other notes on some restricted-range forest birds from West Kilombero in the Udzungwa Mountains, Tanzania. *Scopus* **17**: 48–59.
- Dinesen, L., T. Lehmberg, J.O. Svendsen, L.A. Hansen & J. Fjeldså (1994). A new genus and species of perdicine bird (Phasianidae, Perdicini) from Tanzania: a relict form with Indo-Malayan affinities. *Ibis* **136**: 2–11.
- Dinesen, L., T. Lehmberg, M. Rahner & J. Fjeldså (2001). Conservation priorities for the forests of the Udzungwa Mountains, Tanzania, based on primates, duikers and birds. *Biological Conservation* **99**: 223–226.
- Ferguson-Lees, J. & D.A. Christie (2001). *Raptors of the world*. Christopher Helm, London.
- Fishpool, L.D.C. & Evans, M.I., eds (2001). *Important Bird Areas in Africa and Associated Islands: Priority Sites for Conservation*. Pisces Publications and Birdlife International, Newbury and Cambridge.
- Friedmann, H. & J.G. Williams (1970). *The Birds of the Kalinzu Forest, Southwestern Ankole, Uganda*. Los Angeles County Museum Contributions in Science **195**.
- Haring, E., K. Kvaløy, J.-O. Gjershaug, N. Røv & A. Gamauf (2007). Convergent evolution and paraphyly of the hawk-eagles of the genus *Spizaetus* (Aves, Accipitridae) - phylogenetic analyses based on mitochondrial markers. *Journal of Zoological Systematics and Evolutionary Research* **45**: 353–365.

- Helbig, A.J., A. Kocum, I. Seibold & M.J. Braun (2005). A multi-gene phylogeny of accipitrine eagles (Aves: Accipitriformes) reveals extensive paraphyly at the genus level. *Molecular Phylogenetics and Evolution* **35**: 147–164.
- Howell, K.M. (1993). Herpetofauna of the eastern African forests. In J.C. Lovett & S.K. Wasser (eds), *Biogeography and Ecology of the Rain Forests of Eastern Africa*. Cambridge University Press, Cambridge. Pp. 173–201.
- Hunter, N., C. Carter & E. Mlungu (1996). Recent observations in Udzungwa and Uluguru Mountains, Central Tanzania. *African Bird Club* **3**(2): 96–98.
- Jones, T. (2006). Kipunji in Ndundulu Forest, Tanzania: Distribution, Abundance and Conservation Status. Unpublished report for the Critical Ecosystem Partnership Fund, Fauna and Flora International and the Wildlife Conservation Society. 28pp. [www.cepf.net/xp/cepf/static/pdfs/Jones\\_kipunji\\_report\\_2006\\_website.pdf](http://www.cepf.net/xp/cepf/static/pdfs/Jones_kipunji_report_2006_website.pdf) [accessed 6 August 2007].
- Jones, T., C.L. Ehardt, T.M. Butynski, T.R.B. Davenport, N.E. Mpunga, S.J. Mchaga & D.W. DeLuca (2005). The highland mangabey *Lophocebus kipunji*: A new species of African monkey. *Science* **308**: 1161–4.
- Lerner, H.R.L. & D.P. Mindell (2005). Phylogeny of eagles, Old World vultures, and other Accipitridae based on nuclear and mitochondrial DNA. *Molecular Phylogenetics and Evolution* **37**: 327–346.
- Lovett, J.C. (1993a) Climatic history and forest distribution in eastern Africa. In J.C. Lovett & S.K. Wasser (eds), *Biogeography and Ecology of the Rain Forests of Eastern Africa*. Cambridge University Press, Cambridge. Pp. 23–33.
- Lovett, J.C. (1993b). Temperate and tropical floras in the mountains of eastern Tanzania. *Opera Botanica* **121**: 217–227.
- Marshall, A.R., J.E. Topp-Jørgensen & H. Brink (2001). Bird observations from West Kilombero Scarp Forest Reserve. In K.Z. Doody, K.M. Howell & E. Fanning (eds), *West Kilombero Scarp Forest Reserve – Zoological Report*. Report for the Udzungwa Mountains Forest Management and Biodiversity Conservation Project, MEMA, Iringa, Tanzania. Pp. 86–92. [www.easternarc.or.tz/udzungwa#dl](http://www.easternarc.or.tz/udzungwa#dl) [accessed 6 August 2007].
- Rovero, F. & G. Rathbun (2006). A potentially new giant sengi (elephant-shrew) from the Udzungwa Mountains, Tanzania. *Journal of East African Natural History* **95**: 111–115.
- Rovero, F., G.B. Rathbun, A. Perkin, T. Jones, D.O. Ribble, C. Leonard, R.C. Mwakisoma & N. Doggart (in press). A new species of giant sengi or elephant-shrew (genus *Rhynchocyon*) highlights the exceptional biodiversity of the Udzungwa Mountains of Tanzania. *Journal of Zoology*.
- Roy, M.S. (1997). Recent diversification in African greenbuls (Pycnonotidae: Andropadus) supports a montane speciation model. *Proceedings of the Royal Society of London B* **264**: 1337–1344.
- Roy, M.S., J.C. da Silva, P. Arctander, J. Garcia-Moreno & J. Fjeldså (1997). The role of montane regions in the speciation of South American and African birds. In D.P. Mindell (ed.), *Avian Molecular Evolution and Systematics*. Academic Press, London. Pp. 325–343.
- Stanley, W.T., M.A. Rogers & R. Hutterer (2005). A new species of *Congosorex* from the Eastern Arc Mountains, Tanzania, with significant biogeographical implications. *Journal of Zoology* **265**: 269–280.
- White, F. (1981). The history of the Afromontane archipelago and the scientific need for its conservation. *African Journal of Ecology* **19**: 33–54.