

Distribution of the genus *Anyphaena* in the Western Mediterranean region, with the first record of *Anyphaena alboirrorata* in the Maghreb (Araneae: Anyphaenidae)

Authors: Habita, Aicha, Benhalima, Souâd, Kherbouche-Abrous, Ourida, Bosmans, Robert, Brague-Bouragba, Nadia, et al.

Source: Arachnologische Mitteilungen: Arachnology Letters, 63(1) : 19-23

Published By: Arachnologische Gesellschaft e.V.

URL: <https://doi.org/10.30963/aramit6306>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Distribution of the genus *Anyphaena* in the Western Mediterranean region, with the first record of *Anyphaena alboirrorata* in the Maghreb (Araneae: Anyphaenidae)

Aicha Habita, Souâd Benhalima, Ourida Kherbouche-Abrous, Robert Bosmans,
Nadia Brague-Bouragba & Omar Guezoul



doi: 10.30963/aramit6306

Abstract. New data on the distribution of the genus *Anyphaena* Sundevall, 1833 in Mediterranean Europe and North Africa are given. *Anyphaena alboirrorata* Simon, 1878 is newly recorded in the Maghreb. The species was found in the Saharan and the Tell Atlas in Algeria, in one locality in Morocco and three localities in central north Tunisia. In addition, *Anyphaena numida* Simon, 1897 is presented as new to Morocco, together with a further record from Algeria and Tunisia. Supplementary material from Spain was examined and the data are also given. *Anyphaena sabina* L. Koch, 1866 is recorded for the first time in Algeria since records by Denis in 1937.

Key Words: Algeria, Morocco, Saharan Atlas, species range, spiders, Tunisia

Zusammenfassung. Verbreitung der Gattung *Anyphaena* im westlichen Mittelmeerraum, mit dem Erstnachweis von *Anyphaena alboirrorata* im Maghreb. Neue Daten zur Verbreitung der Gattung *Anyphaena* im mediterranen Europa und Nordafrika werden präsentiert. *Anyphaena alboirrorata* Simon, 1878 wurde erstmals im Maghreb nachgewiesen. Die Art wurde im Sahara- und Tellatlas in Algerien, von einer Lokalität in Marokko sowie drei Lokalitäten im nördlichen Bereich von Zentraltunesien gefunden. Zusätzlich wird der Erstnachweis von *Anyphaena numida* Simon, 1897 für Tunesien sowie ein weiterer Nachweis aus Algerien vorgestellt. Ergänzendes Material aus Spanien wurde untersucht und wird ebenfalls präsentiert. *Anyphaena sabina* L. Koch, 1866 ist zum ersten Mal seit dem Fund von Denis im Jahr 1937 in Algerien nachgewiesen.

The spider family Anyphaenidae, commonly called ghost spiders, has long been considered a subfamily of the Clubionidae and the former Drassidae (today Gnaphosidae), but they are sufficiently distinctive to be ranked as an independent family (e.g. Platnick 1974, Ramirez 2003). According to Urones et al. (1995) the position of the tracheal stigma, the structure of lamelliform claw tufts and the extension of its tracheal system seem to be determining features for the whole family. These spiders are mostly nocturnal, wandering hunters.

Anyphaenidae include 57 genera and 614 species worldwide (World Spider Catalog 2022), but only a few species live in the western Palaearctic region (Nentwig et al. 2022). The genus *Anyphaena* is the largest and most unique genus of this family, distributed in Europe and North Africa. From a total of 87 species (World Spider Catalog 2022), 61 are known from North and central America (70%), 18 from Asia (21%), six from Europe (7%) and only two (2%) from Africa: *Anyphaena numida* Simon, 1897 and *Anyphaena sabina* L. Koch, 1866 (Nentwig et al. 2022). In addition to these two species, we add a further one, *Anyphaena alboirrorata* Simon, 1878, which was newly collected in Algeria, Morocco and Tunisia and, which is therefore also new to the Maghreb region.

Material and methods

Study area. The research was conducted in three countries: Algeria, Morocco and Tunisia. In Algeria, three regions were investigated. The first one was the Ouled Nail Range in the Saharan Atlas, on the edge of the desert. This dryland is becoming presently heavily degraded in the process of desertification (Benderradji et al. 2006, Guit & Nedjimi 2020). Two sites were sampled in this region. Djellal Chergui, naturally occurring Aleppo pine trees, 12 km south-east of Djelfa and Sahary Guebli (Fig. 1), 300 km south of Algiers, which is considered as a one of the most important natural stands of Aleppo pine forests in the arid zones of the Algerian Saharan Atlas. Both sites have a semi-arid, cold winter climate, with an annual average temperature of 21.6°C. The average precipitation is estimated at 300 mm/year. The second investigated area in Algeria was Aïn Yagout, located 35 km north-east of Batna in the Aurès region, which is a part of the Saharan Atlas. The third studied region was located in the forest of Djebel Babor in Sétif, a part of the Tell Atlas massif, located just 15 km from the Mediterranean Sea. The precipitation can reach over 2000 mm/year in this area.

In Morocco, the study sites were situated in the two northern provinces: the region of Larache in the province of Tangier and the Maâmora Forest in the province of Rabat. The spiders were sampled here in an Atlantic cork oak (*Quercus suber* L.) forest. In Tunisia, the material was collected from the northern governorates: Kasserine, El Kef and Zaghuan. The map was created with d-maps (2021).

Sampling. Material was collected using pitfall traps or by hand. Specimens were preserved in 70% ethanol. A stereomicroscope (Nikon SMA 1270) was used for the specimens' examination and a Moticam camera mounted on a Realux microscope and Olympus SZX7 stereomicroscope to take photographs. Material was collected in Algeria between 1989 and 2003 by the fourth author and in 2018 by the first author from natural Aleppo pine stands. In Morocco, individuals of *Anyphaena* were collected by the second author during 1989.

Aicha HABITA, Department of Biology, Faculty of Natural and Life Sciences, University of Djelfa, BP 3117, 17000 Djelfa; Department of Agronomy, Faculty of Natural and Life Sciences, University of Kasdi-Merbah, 30000 Ouargla, Algeria; E-mail: habita_a@yahoo.fr
Souâd BENHALIMA, Geo-Biodiversity and Natural Patrimony Laboratory (GEOBIO), Scientific Institute, Research Centre (GEOPAC), Mohammed V University in Rabat, BP 703, Agdal, 10090, Rabat, Morocco; E-mail: souad.benhalima@is.um5.ac.ma
Ourida KHERBOUCHE-ABROUS, Laboratory of Dynamics and Biodiversity, Faculty of Biological Sciences, University of Sciences and Technology Houari Boumediene BP 32 El Alia, Bab Ezzouar, Algiers, Algeria; E-mail: ouridakherbouche@yahoo.fr
Robert BOSMANS, Terrestrial Ecology Unit, Ledeganckstraat 35, B-9000 Gent, Belgium; E-mail: rop_bosmans@telenet.be
Nadia BRAGUE-BOURAGBA, National Forest Research Institute, 17000 Djelfa, Algeria; E-mail: bouragbanadia@yahoo.fr
Omar GUEZOU, Department of Agronomy, Faculty of Natural and Life Sciences, University of Kasdi-Merbah, 30000 Ouargla, Algeria; E-mail: oguezoul@yahoo.fr

Academic editor: Konrad Wiśniewski

submitted: 18.2.2022, accepted: 17.6.2022, online: 26.8.2022



Fig. 1: Sampling area for *Anyphaena alboirrorata*: natural strips of forests in Djebel Bahrara (Sahary Guebli)

In Tunisia, spiders were collected in 2003 and in Spain between 1996 and 1999 by the fourth author.

Abbreviations

CRB: Collection Robert Bosmans, CAH: Collection Aicha Habita, CYA: Collection Youcef Alioua, CSB: Collection Souâd Benhalima.

Results

Anyphaena Sundevall, 1833

Anyphaena alboirrorata Simon, 1878 (Figs 2–4a, 5a)

For detailed taxonomic references, see the World Spider Catalog (2022).

Previous records in the Western Mediterranean. FRANCE: Marseille, Toulon (Simon 1878); Marseille, Nice, Perpignan, Toulon (Simon 1932). SPAIN: Balearic Islands, Cáceres, Cádiz, Huelva, Huesca, Jaén, Madrid, Navarra, Salamanca, Zaragoza (Branco et al. 2019, De Biurrun et al. 2019); Valladolid (Urones 1996). PORTUGAL: Bragança (Morano et al. 2019), Guarda (Machado 1949). ITALY: Bari, Bologna, Forlì-Cesena, Perugia, Trieste, Udine (Pantini & Isaia 2019). **New records.** ALGERIA: Sétif: Djebel Babor (31.50972°N, 5.48750°E), 1750 m a.s.l., pitfall trap in a *Cedrus* forest, 22. Oct. 1989, 1 ♀ (CRB). Batna: Ain Yagout (35.77444°N, 6.41638°E), 950 m a.s.l., 27. Dec. 2016, 1 ♀ (CYA). Djelfa: Djebel Bahrara (34.79944°N, 3.29055°E), 1170 m a.s.l., pitfall trap in a natural forest, 20. Dec. 2018, 1 ♀ (CAH); Djebel Djellal (34.57152°N, 3.39027°E), 1356 m a.s.l., pitfall trap in a natural forest, 20. Dec. 2018, 1 ♀ (CAH). MOROCCO: Tanger-Tétouan-El Hoceima: Larache, Rhaba El Khalifa (35.12138°N, 6.71972°W), 33 m a.s.l., pitfall trap in litter layer of *Quercus suber* forest, 12.–19. Jan. 1989, 1 ♀ (CSB). TUNISIA: Kasserine: Haidra S. (35.54472°N, 8.45750°E), 950 m a.s.l., under stones in a *Pinus* forest, 4. Mar. 2005, 4 ♀♀ (CRB). Le Kef: Hammam Mellègue (36.11638°N, 8.48972°E), 900 m a.s.l., under stones in a *Pinus* forest, 4. Mar. 2005, 1 ♀ (CRB). Zaghuan: El Fahs S. (36.37416°N, 9.90638°E), 195 m a.s.l., under stones in a *Pinus* forest, 27. Jan. 2003, 1 ♀ (CRB). SPAIN: Cuenca: Casillas de Ranera (39.78333°N, 1.25000°W), 800 m a.s.l., under stones in a *Pinus* forest, 8. Apr. 1997, 1 ♀ (CRB); Molina de Aragón

(40.86083°N, 1.83277°W), 1050 m a.s.l., stones in a *Pinus* forest, 14. Apr. 1998, 2 ♀♀ (CRB). Granada: El Baúl (37.42805°N, 2.92916°W), 900 m a.s.l., under stones in a *Quercus ilex* L. forest, 11. Apr. 1999, 2 ♀♀ (CRB). Jaén: Jabalcuz (37.73111°N, 3.80916°W), 700 m a.s.l., under stones in *Pinus* forest, 12. Apr. 1998, 2 ♀♀ (CRB). Teruel: Agua-viva (40.80166°N, 0.15277°W), 540 m a.s.l., under stones in *Pinus* forest, 2. Apr. 1996, 2 ♀♀ (CRB). idem, 8. Apr. 1997, 1 ♀ (CRB). Valladolid: Tordesillas, SW (41.48055°N, 5.05361°W), 670 m a.s.l., litter of a *Pinus* forest, 10. Apr. 1996, 4 ♀♀ (CRB). Zaragoza: Daroca (41.11444°N, 1.41444°W), 775 m a.s.l., under stones in *Pinus* plantation, 14. Apr. 1998, 2 ♀♀ (CRB).

Distribution. Southern and western part of the Mediterranean region (World Spider Catalog 2022). In Morocco, *Any-*



Fig. 2: *Anyphaena alboirrorata* from Djelfa (Algeria). Female, ventral view (Oger 2021)




Fig. 3: Male palp of *Anyphaena alboirrorata* from Djelfa (Algeria). **a.** ventral view; **b.** prolateral view (Oger 2021)




Fig. 4: Male palps in retrolateral view. **a.** *Anyphaena alboirrorata* from Djelfa (Algeria); **b.** *Anyphaena numida* from France; **c.** *Anyphaena sabina* from France (Oger 2021)

phaena alboirrorata was collected in the northern part, while in Algeria, the species occupies a steppe region (Djelfa). For Tunisia, the species was collected in the west-central part. These are the first records for the three countries.

***Anyphaena numida* Simon, 1897** (Figs 4b, 5b)

For detailed taxonomic references, see the World Spider Catalog (2022).

Previous records in the Western Mediterranean. FRANCE: Pyrennees-Orientales, 1000 m (Simon 1932). ALGERIA: Algiers: Algiers, 36.73222°N, 3.08750°E (Simon 1896). El Harrach (Maison Carré), 36.71277°N, 3.14888°E (Simon 1896). Blida: Gorges de La Chiffa, 36.40000°N, 2.76666°E (Simon 1896). PORTUGAL: Bragança, Portalegre (Morano et al. 2019), Braga (Machado 1937), Guarda, Porto (Simon 1897). SPAIN: Asturias, Ávila, Andorra, Barcelona, Cáceres, Cantabria, Gipuzkoa, Madrid, Salamanca, Segovia (De Biurrun et al. 2019, Morano et al. 2019), A Coruña (Simon 1897), Zamora (Urones 1987).

New records. ALGERIA: Tizi Ouzou: Djurdjura mountains, Tala Guilef (36.47666°N, 3.99666°E), 1400 m a.s.l., litter in *Cedrus* forest, 25. Nov. 1985, 1 ♀ (CRB). MOROCCO: Rabat-Salé-Kénitra: Maâmora cork forest, Sidi Amira (34.05166°N, 6.15083°W), 129 m a.s.l., pitfall trap in litter layer of *Quercus suber* forest, 1.–6. Nov. 1989, 2 ♀♀ (CSB).

Distribution. Portugal, Spain and France (World Spider Catalog 2022) and United Kingdom (Harvey 2017). In the Maghreb the species occurs only in Algeria and Morocco. It is the first record for Morocco.

***Anyphaena sabina* L. Koch, 1866** (Figs 4c, 5c)

For detailed taxonomic references, see World Spider Catalog (2022).

Previous records in the Western Mediterranean. FRANCE: Corsica (Simon 1932), Bouches-du-Rhône, Pyrénées-Orientales, Midi de la France (Le Peru 2007). ITALY: Cagliari, Carbonia-Iglesias Catania, Catanzaro, Cosenza, Firenze, Foggia, Forlì-Cesena, Grosseto Lecce, Medio Campidano Nuoro Olbia-Tempio, Oristano, Perugia, Roma, Salerno, Sassari, Siracusa, Udine, Venezia (Pantini & Isaia 2019). Isola del Giglio (Simon 1932). SPAIN: Andorra, Ávila, Barcelona, Cáceres, Cádiz, Granada, La Rioja, Salamanca, Zaragoza (Morano et al. 2019; Pantini & Isaia 2019). PORTUGAL: Bragança, Faro, Portalegre, Porto, Setúbal (Morano et al. 2019). ALGERIA: Mila: Djebel Daya (Denis 1937).

New records. ALGERIA: Tissemsilt: Theniet-el-Had National Park, Rond Point des Cèdres (35°52'15"N 1°56'41"E), 1550 m a.s.l., *Cedrus* forest, 23. May 1987, 1 ♀ (CRB). Blida: Chréa National Park (36.42916°N, 2.88055°E), 1520 m a.s.l., *Cedrus* forest, 20. May 1988, 1 ♂ (CRB).

Distribution. Southern Europe, United Kingdom, Turkey, Caucasus (World Spider Catalog 2022). In the Maghreb, the species is found only in Algeria.

Distinguishing the three species. Males are easily distinguished by the shape of the tibial retrolateral apophyses (Fig. 4a–c). The females differ in the shape of their epigyne: *A. alboirrorata* lacks an anterior hood (Fig. 5a), *A. numida* has a narrow (Fig. 5b) and *A. sabina* has a very wide one (Fig. 5c).




Fig. 5: Ventral view of epigynes. **a.** *Anyphaena alboirrorata* from Djelfa (Algeria); **b.** *Anyphaena numida* from France; **c.** *Anyphaena sabina* from France (Oger 2021)




Fig. 6: Distribution of *Anyphaena alboirrorata* in the Western Mediterranean region. Circles: previous records (Simon 1878, 1932, Machado 1949, Branco et al. 2019, De Biurrun et al. 2019, Morano et al. 2019, Pantini & Isaia 2019), triangles: new records

Discussion

Three *A. alboirrorata* females were collected in the canopy of naturally occurring Aleppo pine trees in Djelfa, where they were living in tree foliage. According to Simon (1878), this *Anyphaena* species lives on tall plants and shrubs. Soyer (1963) collected it on a hill and around salt flats (France), Urones et al. (1995) on vegetation in dry and open areas with a predominance of heather (Portugal and Spain). Urones (1996) also found it on vegetation in open or dry areas such as moors, open pine forests and oak forests (Portugal, Spain and Balearic Islands). Our result is in accordance with these previously published data, since our localities consisted of dry habitats. Apparently, *A. alboirrorata* prefers dry and hot areas with dense vegetation providing support for silken threads of their retreat.

The new records in Algeria, Tunisia and Morocco are the most southern points in the known distribution of *A. alboirrorata*. It was captured at several localities in the northernmost parts of the countries (Fig. 6), from the coast to the steppe region. This suggests that the species is fairly widespread in the region and occupies a large spectrum of different habitats. Soyer (1963) showed that adults are found late and early in the year. In Spain, the females were collected in spring (Urones 1996), however, in Algeria and Morocco adults were also found during the winter.

Anyphaena numida was collected from a cork oak forest in Tunisia. This is in accordance with Urones et al. (1995) who mentioned different Mediterranean forest types as a typical habitat for the species.

The here presented data show that the Maghreb region is still largely unexplored, and more efforts are needed to uncover the occurrence and distribution of spiders and other arachnids.

Acknowledgements

We would like to thank Dr. Lynda Beladjal and Prof. Johan Mertens (University of Ghent) for their kind, warm welcome and support during our visit in Belgium and Pierre Oger for the excellent photos. Moreover, the authors express their sincere thanks to Dr. Alioua Youcef (University of Ghardaia) for providing some of the material as well as the reviewers for their careful reading of our manuscript and their many insightful comments and suggestions.

References

- Branco V, Morano E & Cardoso P 2019 An update to the Iberian spider checklist (Araneae). – *Zootaxa* 4614: 201–254 – doi: [10.11646/zootaxa.4614.2.1](https://doi.org/10.11646/zootaxa.4614.2.1)
- Benderradji MEH, Alatou D, Arfa AMT & Benachour K 2006 Problemes de degradation de l'environnement par la desertification et la deforestation. Impact du phenomene en Algerie. – *New Medit* 5 (4): 15–22
- Denis J 1937 On a collection of spiders from Algeria. – *Proceedings of the Zoological Society of London* 106: 1027–1060 – doi: [10.1111/j.1469-7998.1936.tb06301.x](https://doi.org/10.1111/j.1469-7998.1936.tb06301.x)
- De Biurrun G, Prieto C & Baquero E 2019 Iberian Spider Catalog. Actualización del mapa web y sus funciones. Arachnomap 2019. – Internet: <http://sea-entomologia.org/gia/map> (1. Jun. 2022)
- d-maps 2021 d-maps.com free maps. – Internet: <http://d-maps.com> (17. Apr. 2021)
- Guit B & Nedjimi B 2020 Croissance radiale du pin d'Alep (*Pinus halepensis* Mill.) en fonction des paramètres stationnels dans les massifs forestiers naturels de l'Atlas saharien algérien. – *Bois et Forêts des Tropiques* 345: 3–11 – doi: [10.19182/bft2020.345.a31803](https://doi.org/10.19182/bft2020.345.a31803)
- Harvey P 2017 The buzzing spider *Anyphaena numida* Simon, new to Britain. – *Newsletter of the British Arachnological Society* 140: 18–20
- Le Peru B 2007 Catalogue et répartition des araignées de France. – *Revue Arachnologique* 16: 1–468
- Machado A de B 1937 Aranhas novas para a fauna portuguesa (I). – *Memórias e Estudos do Museu zoológico da Universidade de Coimbra. Seria I*, 107: 1–7
- Machado A de B 1949 Araignées nouvelles pour la faune portugaise (III). – *Memórias e Estudos do Museu zoológico da Universidade de Coimbra* 191: 1–69
- Morano E, Branco V, Carrillo J & Cardoso P 2019 Iberian Arachnology Group (GIA) version III – Internet: <http://sea-entomologia.org/gia/map/index.html> (19. Dec. 2021)
- Nentwig W, Blick T, Bosmans R, Gloor D, Hänggi A & Kropf C 2022 Spiders of Europe. Version 06.2022. – Internet: <https://araneae.unibe.ch/> (21. Jun. 2022) – doi: [10.24436/1](https://doi.org/10.24436/1)
- Oger P 2021 Familles / Anyphaenidae | Les araignées de Belgique et de France (piwigo.com) – Internet: arachno.piwigo.com (12. Jun. 2021)
- Pantini P & Isaia M 2019 Araneae.it: the online Catalog of Italian spiders with addenda on other arachnid orders occurring in Italy (Arachnida: Araneae, Opiliones, Palpigradi, Pseudoscorpionida, Scorpiones, Solifugae). – *Fragmenta Entomologica* 51: 127–152 – Internet: www.araneae.it (4. Jun. 2022)
- Platnick NI 1974 The spider family Anyphaenidae in America north of Mexico. – *Bulletin of the Museum of Comparative Zoology* 146: 205–266
- Ramirez MJ 2003 The spider subfamily Amaurobioidinae (Araneae, Anyphaenidae): a phylogenetic revision at the generic level. – *Bulletin of the American Museum of Natural History* 277: 1–261 – doi: [10.1206/0003-0090\(2003\)277<0001:TSSAAA>2.0.CO;2](https://doi.org/10.1206/0003-0090(2003)277<0001:TSSAAA>2.0.CO;2)
- Simon E 1878 Les arachnides de France. Tome VI. Roret, Paris. 334 pp.
- Simon E 1896 Descriptions d'arachnides nouveaux de la famille des Clubionidae. – *Annales de la Société Entomologique de Belgique* 40: 400–422
- Simon E 1897 Etudes arachnologiques. 27e Mémoire. XLII. Descriptions d'espèces nouvelles de l'ordre des Araneae. – *Annales de la Société Entomologique de France* 64: 465–510
- Simon E 1932 Les arachnides de France. Synopsis générale et catalogue des espèces françaises de l'ordre des Araneae. Tome VI, 4e partie, Roret, Paris. pp. 773–978
- Soyer B 1963 Contribution à l'étude éthologique et écologique des araignées de la Provence occidentale. VI: Les araignées thomisides et clubionides des collines et des terrains sales. – *Bulletin de la Société linnéenne de Provence* 23: 27–34

- Urones C 1987 Distribución y ecología de las arañas en la provincia de Zamora. – Anuario del Instituto de Estudios Zamoranos “Florián de Ocampo” 1986: 67-122
- Urones C 1996 Catálogo y atlas de las arañas de la familia Anyphaenidae en la península Ibérica e islas Baleares. – Graellsia 52: 73-80
- Urones C, Barrientos JA & Espuny A 1995 El género. *Anyphaena* Sundevall, 1833 (Araneae: Anyphaenidae) en la Península Ibérica. – Boletín de la Asociación Española de Entomología 19: 109-131
- World Spider Catalog 2022 World spider catalog. Version 23.0. Natural History Museum, Bern. – Internet: <http://wsc.nmbe.ch> (4. Jun. 2022) – doi: [10.24436/2](https://doi.org/10.24436/2)