



## **Philodendron cardosoi, a new species of Araceae from Brazil**

Author: Gonçalves, Eduardo G.

Source: Willdenowia, 34(2) : 525-527

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.34.34215>

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

doi:10.3372/wi.34.34215 (available via <http://dx.doi.org/>)

EDUARDO G. GONÇALVES

## *Philodendron cardosoi*, a new species of *Araceae* from Brazil

### Abstract

Gonçalves, E. G.: *Philodendron cardosoi*, a new species of *Araceae* from Brazil. – Willdenowia 34: 525-527. – ISSN 0511-9618; © 2004 BGBM Berlin-Dahlem.

*Philodendron cardosoi* is described as a species new to science and illustrated. It is a member of *P.* subg. *Philodendron* sect. *Polytomium* and is so far only known from the type locality in the southeast of the Brazilian state of Pará.

The largest genera in the *Araceae* are *Anthurium* and *Philodendron*, with approximately 700 and 400 species, respectively. Even though there are serious efforts to elucidate the taxonomy of both genera, new species are still added yearly. Some revisional efforts have doubled the number of known species of *Philodendron* in general areas (Grayum 1996, Croat 1997). During a brief visit to the city of Belém in 2003, I was invited by some collectors to identify their plants. Among many interesting and poorly known species, one plant was recognized as a new species of *Philodendron* and it is here described as new and illustrated.

*Philodendron cardosoi* E. G. Gonç., **sp. nova** – Fig. 1

Holotypus: Brazil, Pará, São Geraldo do Araguaia, Parque Estadual da Serra dos Martírios / Andorinhas, nas margens do Rio Araguaia, próximo à Vila de Santa Cruz, no lugar denominado Pedra Escrita, c. 200 m, 7.2003, A. L. R. Cardoso (IAN; isotypus: UB).

Ad subgenus *Philodendron* sectionem *Polytomium* (Schott) Engl. pertinens. *Herba* terricola, caudex prorepens, internodiis 2-4 × 4-5 cm; prophyllum deciduum, rectum. *Petiolus* 45-47.5 × 1.3-1.5 cm, supra leviter deplanatus; *lamina* ambitu ovata, 40-60 × 32-45 cm, pinnatipartita; lobis primariis lateralibus lanceolatis, 6-7 utroque. *Pedunculus* 16-22 cm longus, 0.6-1 cm diam.; *spatha* 9-12 cm longa, extus albo-viridis, intus albida marginibus roseis; *spadix* 8.5-9.3 cm; *inflorescentia femina* 3-3.2 × 0.6-1.1 cm, *mascula sterilis* 0.5-0.6 × 1-1.1 cm, *mascula fertilis* 5.5-6.8 × 0.8-1 cm; *ovaria* 8-9-locularia, ovula 2-3 per loculum, ad axem affixa.

Terrestrial herb. *Stem* creeping, prostrate and rooting; internodes glossy green, 2-4 × 4-5 cm, prophylls purplish, straight, up to 18-22.5 cm long. *Leaves* erect; *petioles* 45-47.5 × 1.3-1.5 cm, slightly flattened adaxially, with short wine-coloured stripes; *sheath* 2-3 cm long; *leaf blade* pinnatipartite, ovate in outline, 40-60 × 32-45 cm, fresh leaves semi-glossy on both sides, drying light brown; anterior division 27-30 × 29-32 cm, primary lateral veins 6-8 per side, arising at an




Fig. 1. *Philodendron cardosoi* E. G. Gonç. – A: spadix; B: leaf blade; C: habit; D: staminode, side view; E: male flower, side view; F: gynoecium, longitudinal section; G: ovary, cross section. – Drawn from *Cardoso s.n.* by the author.

angle of 50-60° from the midrib, lobes oblanceolate, 18-30 × 1.2-2.5 cm, 8.5-12 times longer than wide, acuminate at apex, sinus ending 3-5 mm from the midrib; posterior divisions 9-11 × 27-28 cm, sinus parabolic, basiscopic lobes 1-2 per side, acroscopic 2-3; basal veins denuded for 1-1.2 cm. *Inflorescence* erect at anthesis, one per leaf axil; *peduncle* 16-22 × 0.6-1 cm; concolourous with petiole; *spathe* 9-12 cm long, whitish green with medium green stripes outside, white in-

side, slightly constricted at middle, edged with pinkish margins; *spadix* 8.5-9.3 cm long; *fertile male portion* 5.5-6.8 × 0.8-1 cm, tapering to the apex; *sterile male portion* 0.5-0.6 × 1-1.1 cm; *female portion* 3-3.2 × 0.6-1.1 cm, extending up to  $\frac{1}{3}$  of the spadix length. *Staminate flowers* prismatic, 4-androus, 1.5-2 × 0.6-1 mm; *staminodes* (sterile flowers) subprismatic, 2 × 1 mm. *Pistillate flowers* with gynoecium 1.3-1.5 mm long, stigma rounded, almost as wide as the ovary, c. 1 mm diam., style inconspicuous; ovary elliptic to obovate, 1-1.2 mm diam., 8-9-locular, 2-3-ovulate, ovules attached to the middle of the flower axis, funicle shorter than the ovules. *Infructescence* unknown.

*Relationships.* – *Philodendron cardosoi* belongs to *P.* subg. *Philodendron* and has to be placed in sect. *Polytomium* (Schott) Engl. because of its pinnate leaves. However, it is the only species in the section that is exclusively terrestrial and it will not climb even if an appropriate vertical support is given. *P. cardosoi* seems to be closer to *P. mayoi* E. G. Gonç., another species with pinnate leaves from Central Brazil (Gonçalves 2000). However, the flowering plants of the latter species are always hemi-epiphytic. The new species may also be confused with the Amazonian *P. tortum* M. L. Soares & Mayo, but differs in having fewer primary lateral lobes (6-7 instead of 9-10) and for the straight (not curved) prophyll (Soares & Mayo 2001). Another important difference is that the leaves are held erect in *P. tortum* and inflexed in *P. cardosoi*.

*Eponymy.* – The epithet is in honor of André Cardoso, a plant enthusiast who has been collecting, growing and breeding aroids, and showed me a small group of this *Philodendron* in his home in Belém, where I was able to recognize it as a new species.

*Distribution.* – *Philodendron cardosoi* occurs in southeastern Pará and – probably – in the adjacent state of Tocantins, in central N Brazil, on quartzite outcrops. *P. cardosoi* grows as a terrestrial herb and its creeping stem usually is buried under the litter.

## Acknowledgements

I would like to thank the plant collectors João Batista Fernandes da Silva and André Cardoso for allowing me to study their plants. I am also grateful to the Instituto Plantarum de Estudos da Flora Ltda for logistic help.

## References

- Croat, T. B. 1997: A revision of *Philodendron* subgenus *Philodendron* (*Araceae*) for Mexico and Central America. – Ann. Missouri Bot. Gard. **84**: 311-704. [[CrossRef](#)]
- Gonçalves, E. G. 2000: Two new species of *Philodendron* (*Araceae*) from Central Brazil. – Kew Bull. **55**: 175-180. [[CrossRef](#)]
- Grayum, M. H. 1996: Revision of *Philodendron* subgenus *Pteromischum* (*Araceae*) for Pacific and Caribbean tropical America. – Syst. Bot. Monogr. **47**.
- Soares, M. L. C. & Mayo, S. J. 2001: Three new species of *Philodendron* (*Araceae*) from the Ducke Forest Reserve, central Amazonas, Brazil. – Feddes Repert. **112**: 42-44.

Address of the author:

Eduardo G. Gonçalves, Curso de Ciências Biológicas, Universidade Católica de Brasília, Sala M-206, QS 7, Lote 1, EPTC, CEP 72030-170, Taguatinga, DF, Brazil; e-mail: eduardog@ucb.br