



Nuclear SSR Markers for Miscanthus, Saccharum, and Related Grasses (Saccharinae, Poaceae)

Authors: Hodkinson, Trevor R., Cesare, Mariateresa de, and Barth, Susanne

Source: Applications in Plant Sciences, 1(11)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1300042>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

PRIMER NOTE

NUCLEAR SSR MARKERS FOR *MISCANTHUS, SACCHARUM*, AND RELATED GRASSES (SACCHARINAE, POACEAE)¹

TREVOR R. HODKINSON^{2,3,5}, MARIATERESA DE CESARE^{2,4}, AND SUSANNE BARTH⁴

²School of Natural Sciences, Trinity College, Dublin D2, Ireland; ³Trinity Centre for Biodiversity Research, Trinity College, Dublin D2, Ireland; and ⁴Teagasc Crops Environment and Land Use Programme, Oak Park Research Centre, Carlow, Ireland

- *Premise of the study:* We developed nuclear simple sequence repeat (SSR) markers for the characterization of the biomass crop *Miscanthus*, especially *M. sacchariflorus*, *M. sinensis*, and *M. ×giganteus*, and tested for cross-species amplification.
- *Methods and Results:* Twenty-nine SSR markers (di- and tetranucleotide repeats) were developed from DNA sequences obtained from 192 clones from an enriched genomic library of *M. sinensis*. All markers were successfully amplified in *M. sacchariflorus*, *M. sinensis*, and *M. ×giganteus*, and 19 amplified across a broad range of *Miscanthus* species. Polymorphism information content and expected heterozygosity values (19 locus sample) were 0.88 and 0.89, respectively, for *M. sinensis*, 0.48 and 0.54 for *M. sacchariflorus*, and were the lowest in *M. ×giganteus* (0.33, 0.41). Thirteen out of 19 primer pairs showed cross-species amplification in non-*Miscanthus* sensu stricto taxa.
- *Conclusions:* The new set of 29 SSR markers will be of high value for characterizing *Miscanthus* germplasm collections, for prebreeding, and for assessing variation in natural populations.

Key words: cross-species amplification; microsatellites; *Miscanthus*; Poaceae; *Saccharum*; SSRs.

Miscanthus Andersson is under development as a biomass crop and has been characterized by a wide range of markers including amplified fragment length polymorphism (AFLP; Hodkinson et al., 2002), restriction fragment length polymorphism (RFLP; Hernández et al., 2001), inter-simple sequence repeat (ISSR) PCR, and DNA sequences of nuclear and chloroplast regions generated using conventional (Hodkinson et al., 2002) and next-generation approaches including RNAseq and genotyping by sequencing (GBS; Ma et al., 2012). Simple sequence repeat (SSR) markers from maize and *Brachypodium distachyon* (L.) P. Beauv. (Hernández et al., 2001; Zhao et al., 2011) have been successfully applied to *Miscanthus*, and chloroplast SSRs have been developed by De Cesare et al. (2010).

Some nuclear SSR markers have also been developed, such as those for *M. sinensis* Andersson, *M. floridulus* (Labill.) Warb. (Ho et al., 2011), and several other *Miscanthus* species (Zhou et al., 2011). However, there is a need to develop additional SSR markers for *Miscanthus* as the total number of available markers is limited. There is also a need to test these markers on a range of species, especially *M. sacchariflorus* (Maxim.) Hack., *M. sinensis*, and *M. ×giganteus* Greef & Deuter ex Hodk. & Renvoize as these comprise the main species of germplasm collections. SSRs developed from *Saccharum officinarum* L. expressed sequence

tags (ESTs) have been recently used by Kim et al. (2012) to generate genetic maps of *M. sacchariflorus* and *M. sinensis* with genome coverage of 72.7% and 84.9%, respectively. The numbers of linkage groups found for the two maps (40 for *M. sacchariflorus* and 23 for *M. sinensis*) were higher than the basic chromosome number for *Miscanthus* ($x = 19$). Additional markers, such as those generated in this study, will be required to make more saturated maps, especially from noncoding regions that are underrepresented in current maps. Recently, single-nucleotide polymorphism (SNP) markers generated using GBS markers have been used for high-resolution mapping and identified all 19 linkage groups in *M. sinensis* (Ma et al., 2012).

METHODS AND RESULTS

DNA samples were either freshly extracted or obtained from the DNA bank at Trinity College, Dublin. Fresh leaves were frozen in liquid nitrogen and ground manually to a fine powder. Total genomic DNA was extracted following a modified cetyltrimethylammonium bromide (CTAB) method (Hodkinson et al., 2007). Total genomic DNA from the *M. sinensis* clone SW217 was used by ATG Genetics (Vancouver, British Columbia, Canada) to build a nuclear microsatellite-enriched library. After digestion with multiple 4-cutter restriction enzymes, enrichment for SSRs containing fragments was obtained through biotinylated TC_n, TG_n, and GATA_n simple sequence motifs. The selected fragments were cloned into the EcoRI site of the plasmid pUC19 and screened for positive clones using ³²P-labeled TC_n, CA_n, and GATA_n simple sequence motifs. Two 96-well microtiter plates containing single positive bacterial colonies, one selected for the presence of dinucleotide repeats and the second for the presence of tetranucleotide repeats, were produced. The 192 clones were sequenced by AGOWA GmbH (Berlin, Germany), and SSRs were identified in the clones using "find microsat Win32" (Salamini, unpublished). All 192 clones contained SSRs (96 dinucleotides and 96 tetranucleotides). Eighty primer pairs were designed equally among these sets using Primer3 software (Rozen and Skaltsky, 2000; <http://frodo.wi.mit.edu/primer3/>) and tested with PCR. Selection of the final sample of 29 primers was based on clarity of product on an agarose gel. Primer details and GenBank numbers are provided in Table 1.

¹Manuscript received 20 May 2013; revision accepted 12 July 2013.

The study was funded by the National Development Plan of Ireland through Teagasc core funding. M.d.C. was financed under the Teagasc Walsh Fellowship Ph.D. Scheme. We thank TCD Botanic Gardens Dublin; Svalöf Weibull, Sweden; Royal Botanic Gardens Kew, United Kingdom; and the University of Hohenheim, Germany, for sharing their *Miscanthus* resources.

⁵Author for correspondence: Trevor.Hodkinson@tcd.ie

TABLE 1. Characteristics of 29 primer pairs developed for microsatellite genotyping.

Locus	Clone, GenBank accession no.	Repeat motif	Fluorescent dye	Forward primer sequence (5'-3')	Reverse primer sequence (5'-3')	T _a (°C)	Sequence length (bp)	SSR size (bp)
Mis-1	SSR1A10, KF130838	(TCTA) ₂₀	FAM	CAGTCCTTGGAGCAGGCTAT	AAGATCTCAAACCTATAGTC	54	202	80
Mis-13	SSR1F10, KF130839	(TAGA) ₁₉	ROX	CGGACTAACCTGTGAATCTT	GTCCTTGGAGCAGGCTATGA	54	230	76
Mis-14	SSR1F12, KF130840	(GATA) ₁₅	FAM	GTAGCTGCAACTGCTAGTGT	ACTCGCATTGGTTGGTATGA	59	141	60
Mis-15	SSR1F2, KF130841	(ATCT) ₁₆	FAM	ACTACTGCATGCATCATGATG	TGCTTCGCCGCGAAGTTCA	59	195	64
Mis-16	SSR1F5, KF130842	(TATC) ₁₃ / (TCTA) ₁₆	VIC	ATCTTGCCTAGGATGCATTAG	TGGTCTATTACAACAAAGGCT	60	264	52+64*
Mis-20	SSR1G12, KF130843	(TCTA) ₁₇	TAMRA	TAGCTGAGCTGTCTATGGTA	TAGCCATTGAGGCTAAGGAT	54	249	68
Mis-22	SSR1G8, KF130844	(TAGA) ₁₇	VIC	CGAGCGAGCCTGCATGTGTG	TTGACGTCAGCAAGATATTG	54	173	68
Mis-23	SSR1G9, KF130845	(ATCT) ₁₅	TAMRA	CACGAACTGAATCAGCATGC	GTAGCTGCAACTGCTAGTGT	60	240	60
Mis-24	SSR1H10, KF130846	(AGAT) ₁₅	VIC	ATACACGATCCAACATGTC	ATGTGTCACCAAGAGATG	60	324	60
Mis-33	SSR2B7, KF130847	(CT) ₂₀	TAMRA	TGACATAGGCTACACATAT	CGAGTGAGGCACTATTCA	48	242	40
Mis-37	SSR2D9, KF130848	(TC) ₃₄	FAM	GAATGCAGTCATCAGCAGCT	TGGACATCTCTAGTTGATC	54	218	68
Mis-41	SSR2F5, KF130849	(GA) ₂₄	ROX	ATAATGCAGGTCAGTCAAC	CGCAGCTAGCTGCTTGTCA	54	226	48
Mis-42	SSR2F6, KF130850	(AG) ₃₁	FAM	GCCGCCAGGCTCCAAGCCT	ATCCGAGCCATGTATGCACG	54	206	62
Mis-50	SSR2H9, KF130851	(GA) ₂₁	ROX	TACGGACGATTAACCAAGCC	CGCAAGGTGCAGGACCATCA	54	230	42
Mis-51	SSR2G4, KF130852	(TC) ₂₀	FAM	GATCCATACGGATTATCA	ATCATAGGCAAACAGGATCG	60	164	40
Mis-52	SSR2C11, KF130853	(GA) ₁₉	NED	TTATTGGTGCACCAAAGGTGT	AACAAGCCCTCAAGCTTCCT	60	370	38
Mis-53	SSR2G10, KF130854	(GA) ₁₉	FAM	AGGCAGCACCTCACAAA	GGTGGAGATGCTCTTGTG	60	173	38
Mis-54	SSR2A11, KF130855	(CT) ₁₈	NED	TAAGAAACGACGACAGCAGAA	AGTCTCCGGCTTTCTCACAA	60	226	36
Mis-55	SSR2B9, KF130856	(GA) ₁₈	VIC	CGGCTCGAGTGATACCTTT	TACCGGATTTAAGGGCTTT	60	250	36
Mis-59	SSR2B3, KF130857	(GA) ₁₆	FAM	GAGCTGATCGCTAGCAAG	TTCGATAAACAGGGATTGG	60	152	32
Mis-60	SSR2C3, KF130858	(GA) ₁₆	FAM	AGATGGCAGCTTGCTCTTGT	CCATTGTTGAGCACGATGT	60	190	32
Mis-63	SSR1G3, KF130859	(TCTA) ₁₄	VIC	AGGCTAGCACTTCTCCAAA	CTGCTTGGTGACCCCTATAA	60	234	56
Mis-64	SSR1G6, KF130860	(AGAT) ₁₄	NED	TCCCCTTAGTGTCCGTGAAG	GAGGCAGGTGTAGTCGGAGA	60	236	56
Mis-66	SSR1D5, KF130861	(CTAT) ₁₃	VIC	CATGGCTACAGGCCACCTAAA	ATAACGAGAAATGGCGATG	60	165	52
Mis-69	SSR1F4, KF130862	(TCTA) ₁₃	NED	CCTCTGCGGATATGAGGTGT	GAAGTGACAACATGGCATGG	60	175	52
Mis-70	SSR1B10, KF130863	(TATC) ₁₂	NED	TCGCACCTTTAATTTTGAT	TTATGAACCCGACAGGGAGA	60	249	48
Mis-71	SSR1D3, KF130864	(TAGA) ₁₂	VIC	CAACCATGAGCACTTCTCCA	AACATAGGAGGCCAAGCAA	60	179	48
Mis-78	SSR2G11, KF130865	(CT) ₁₅	NED	TCTGCAGGTGACAAAGGAAGA	GTCAACCGGCATGTTGAT	60	167	30
Mis-79	SSR2G9, KF130866	(CT) ₁₅	VIC	GCCAACTCGTGGATTTGAGT	CGTAGCAAGAGGGAAACAA	60	248	30

Note: T_a = annealing temperature.

*Compound SSR separated by a nonpolymorphic region.

Twenty-nine primer sets provided reliable amplification, and 19 of these were selected to have a mixture of di- and tetranucleotide SSRs. A template DNA volume of 1 μL (40 ng·μL⁻¹) was amplified with an initial denaturation of 5 min at 95°C followed by 35 cycles each with a denaturation of 1 min at 95°C,

1 min at a primer-specific annealing temperature (Table 1), and an extension of 1 min at 72°C, followed by a final extension at 72°C for 10 min. The reaction mixture (final volume) contained 1× reaction buffer containing 2 mM MgSO₄, 0.125 μM dNTPs, 0.25 μM of each primer, and 0.5 U of Taq DNA polymerase

TABLE 2. Genetic properties of the newly developed markers for three *Miscanthus* species.^a

Locus	<i>M. sacchariflorus</i> (n = 9)				<i>M. sinensis</i> (n = 73)				<i>M. ×giganteus</i> (n = 15)			
	A	Size range (bp)	H _e	PIC	A	Size range (bp)	H _e	PIC	A	Size range (bp)	H _e	PIC
Mis-1	2	127–161	0.375	0.305	19	125–256	0.904	0.896	3	125–161	0.370	0.340
Mis-14	2	87–119	0.663	0.604	25	87–208	0.928	0.924	2	99–119	0.500	0.375
Mis-15	3	144–148	0.620	0.548	20	144–205	0.862	0.852	2	146–148	0.500	0.375
Mis-20	2	200–234	0.320	0.269	28	197–300	0.907	0.901	2	200–234	0.499	0.375
Mis-22	1	124	0.000	0.000	14	103–174	0.837	0.818	1	124	0.000	0.000
Mis-23	3	191–223	0.625	0.555	27	191–314	0.935	0.932	2	203–223	0.499	0.375
Mis-24	1	331	0.000	0.000	21	283–361	0.905	0.899	1	331	0.000	0.000
Mis-37	5	160–200	0.789	0.756	27	160–222	0.938	0.935	3	160–226	0.531	0.420
Mis-41	2	214–215	0.444	0.346	35	197–512	0.924	0.919	1	214	0.000	0.000
Mis-42	3	206–247	0.560	0.499	21	163–247	0.909	0.903	4	183–236	0.574	0.500
Mis-50	2	207–256	0.408	0.325	25	199–260	0.869	0.859	2	207–256	0.497	0.373
Mis-51	2	136–140	0.463	0.356	24	132–176	0.887	0.879	1	140	0.000	0.000
Mis-52	6	177–207	0.806	0.777	18	170–207	0.863	0.850	3	177–207	0.557	0.457
Mis-54	5	213–236	0.796	0.763	18	207–244	0.860	0.848	4	213–224	0.647	0.586
Mis-59	7	135–155	0.840	0.820	10	123–160	0.792	0.766	4	148–155	0.678	0.618
Mis-64	4	214–258	0.740	0.692	30	194–286	0.923	0.918	2	232–258	0.476	0.363
Mis-69	3	130–143	0.612	0.541	17	105–197	0.861	0.848	2	130–138	0.500	0.375
Mis-70	3	219–237	0.595	0.526	26	211–328	0.903	0.897	2	219–225	0.500	0.375
Mis-79	3	242–266	0.540	0.466	22	235–274	0.904	0.897	4	224–252	0.479	0.427
Mean			0.537	0.481			0.890	0.881			0.411	0.333

Note: A = number of alleles; H_e = expected heterozygosity; PIC = polymorphism information content.

^aStatistics provided for species where sample size (n) was 9 or greater.

TABLE 3. Cross-amplification of the newly developed microsatellites of *Miscanthus*.^a

Locus	A (<i>n</i> = 166) ^e	Size range (bp)	Miscanthus s.s. ^b										Saccharinae										Other Andropogoneae/Paniceae ^d									
			<i>M. sacchariflora</i>	<i>M. sinensis</i>	<i>M. sinensis</i> subsp. <i>condensata</i>	<i>M. × gigantea</i>	<i>M. transvaalica</i>	<i>M. tenuiseta</i>	<i>M. ecklonii</i>	<i>M. junccea</i>	<i>M. violacea</i>	<i>M. nepalensis</i>	<i>M. violacea</i>	<i>M. violacea</i>	<i>M. nervosa</i>	<i>M. nervosa</i>	<i>M. mucronata</i>	<i>Saccharum officinarum</i>	<i>Saccharum spontaneum</i>	<i>Spodiopogon</i> ^f	<i>Zea diploperennis</i>	<i>Cymbopogon citratus</i>	<i>Sorghum halepense</i>	<i>Pennisetum sp.</i>								
Mis-1	20	125–256	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-14	33	71–208	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-15	21	144–205	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-20	33	197–300	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-22	16	103–174	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-23	30	176–314	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-24	23	248–361	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-37	33	169–226	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-41	44	131–512	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-42	29	121–247	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-50	30	199–260	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-51	27	132–176	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-52	22	132–207	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-54	20	207–244	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-59	13	123–162	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-64	40	177–286	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-69	24	105–220	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
Mis-70	31	211–328	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+			
Mis-79	34	224–276	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+			
Average		27.5																														

^aCross-amplification in *Miscanthus* species, other Saccharinae, other Andropogoneae, and Paniceae (+ = yes; – = no).

^b*Miscanthus* s.s. (Asian *Miscanthus* with basic chromosome number of 19).

^c*Miscanthus* s.l. (GrassBase—The Online World Grass Flora [<http://www.kew.org/data/grasses-dbhtml>]).

^d*Sorghum* is classified in Sorghinae (Andropogoneae), *Cymbopogon* in Andropogoninae (Paniceae), and *Pennisetum* in Cenchrinae (Paniceae).

^eTotal allele number and size range in base pairs (bp) for 19 nuclear SSR markers across all samples.

^f*Spodiopogon rhizophorus* and *S. sibiricus*.

(New England BioLabs, Herts, United Kingdom). Five different fluorescent dyes were used for primer labeling to allow multiplexing, in pools (Table 1). A polyA treatment at 65°C was applied for 30 min to the PCR products. Undiluted PCR products were then sized using an ABI 3130xl automated DNA sequencer (Applied Biosystems, Carlsbad, California, USA) and the resulting peaks were scored with GeneMapper version 4.0 software (Applied Biosystems). All 29 primer pairs produced good amplification on eight test genotypes of *M. sacchariflorus*, *M. sinensis*, and *M. ×giganteus*, but 11 loci were not consistently amplified across our entire collection and were discarded from further analyses. Our final analysis therefore included 19 SSR markers. Allele number, size range, expected heterozygosity (H_e), and polymorphism information content (PIC) were calculated using PIC Calculator Extra (<http://www.genomics.liv.ac.uk/animal/pic.html>). H_e and PIC values were only calculated for *M. sacchariflorus*, *M. sinensis*, and *M. ×giganteus* because of sample size (Table 2).

Polymorphism at 19 microsatellite loci was studied in a collection of 166 individual grasses (Appendix 1), mostly belonging to the species *M. sinensis*, *M. sacchariflorus*, and *M. ×giganteus*. Fourteen individuals belonging to closely related genera were also included. All markers revealed considerable length polymorphism, with the number of alleles ranging from 13 to 44 per locus, with an average of 27.5 (Table 3). The loci amplified included a tetranucleotide repetition in nine cases and a dinucleotide repetition in the remaining 10. No major difference was observed between di- and tetranucleotide microsatellite loci in their ability to detect variation. Thirteen out of 19 primer pairs showed cross-amplification in non-*Miscanthus* species (Table 3). Average allele number was higher than the value of 12 found by Hernández et al. (2001) in a previous study using SSRs from maize. The higher number of clones used in our study (166 against 16 clones) and the introduction of species other than *M. sinensis*, *M. sacchariflorus*, and *M. ×giganteus* could account for the difference in allele number.

PIC and H_e values varied considerably among species (Table 2) and were the highest (0.88 and 0.89, respectively) for *M. sinensis*, 0.48 and 0.54 for *M. sacchariflorus*, and the lowest (0.33 and 0.41) in *M. ×giganteus*. The PIC value of *M. sinensis* (0.88) was consistent with the value of 0.83 in Hernández et al. (2001), both are higher than the average PIC value recently found by Zhao et al. (2011) in a study examining transferability of 49 microsatellite markers from *Brachypodium distachyon* to *M. sinensis*.

In the past few years, the first nuclear microsatellite markers for *Miscanthus* have been developed (Hung et al., 2009; Ho et al., 2011; Zhou et al., 2011). Both studies from Zhou et al. (2011) on transferability from *Brachypodium* P. Beauv. and from Hung et al. (2009) on nine new microsatellite loci specific for *Miscanthus*, were limited to *M. sinensis*, thus explaining the low level of polymorphism found compared to the markers in this study. Zhou et al. (2011) extended the test for their 14 newly developed markers to *M. floridulus*, *M. lutarioriparius* L. Liu ex S. L. Chen & Renvoize, and *M. sacchariflorus*, increasing the average number of alleles found to 16.1 and the PIC value to 0.76. A different approach was used by Ho et al. (2011) to develop 12 new SSR primer pairs for *Miscanthus*. They designed primers based on genic microsatellite loci (EST-SSRs) obtained through transcriptome sequencing and detected an average of 7.9 alleles per locus when tested on *M. floridulus* and *M. sinensis*.

CONCLUSIONS

The newly developed primers presented here were found to cross-amplify not only within *Miscanthus* species but also in other members of the Saccharinae, Andropogoneae, and

Paniceae. They amplified DNA in *Zea* L. (Tripsacinae), *Sorghum* Moench (Sorghinae), *Cymbopogon* Spreng. (Andropogoninae), and *Pennisetum* Rich. (Paniceae). The primers are of high value for characterization of *Miscanthus* species and can be applied to other closely related genera including *Saccharum* L.

LITERATURE CITED

- DE CESARE, M., T. R. HODKINSON, AND S. BARTH. 2010. Chloroplast DNA markers (cpSSRs, SNPs) for *Miscanthus*, *Saccharum* and related grasses (Panicoideae, Poaceae). *Molecular Breeding* 26: 539–544.
- HERNÁNDEZ, P., G. DORADO, D. A. LAURIE, A. MARTÍN, AND J. W. SNAPE. 2001. Microsatellites and RFLP probes from maize are efficient sources of molecular markers for the biomass energy crop *Miscanthus*. *Theoretical and Applied Genetics* 102: 616–622.
- HO, C.-W., T.-H. WU, T.-W. HSU, J.-C. HUANG, C.-C. HUANG, AND T.-Y. CHIANG. 2011. Development of 12 genic microsatellite loci for a biofuel grass, *Miscanthus sinensis* (Poaceae). *American Journal of Botany* 98: e201–e203.
- HODKINSON, T. R., M. W. CHASE, C. TAKAHASHI, I. J. LEITCH, M. D. BENNETT, AND S. A. RENVOIZE. 2002. The use of DNA sequencing (ITS and *trnL-F*), AFLP, and fluorescent in situ hybridization to study allopolyploid *Miscanthus* (Poaceae). *American Journal of Botany* 89: 279–286.
- HODKINSON, T. R., S. WALDREN, J. A. N. PARRELL, C. T. KELLEHER, K. SALAMIN, AND N. SALAMIN. 2007. DNA banking for plant breeding, biotechnology and biodiversity evaluation. *Journal of Plant Research* 120: 17–29.
- HUNG, K.-H., T.-Y. CHIANG, C.-T. CHIU, T.-W. HSU, AND C.-W. HO. 2009. Isolation and characterization of microsatellite loci from a potential biofuel plant *Miscanthus sinensis* (Poaceae). *Conservation Genetics* 10: 1377–1380.
- KIM, C., D. ZHANG, S. A. AUMLAND, L. K. RAINVILLE, K. JAKOB, B. KRONMILLER, E. J. SACKS, ET AL. 2012. SSR-based genetic maps of *Miscanthus sinensis* and *M. sacchariflorus*, and their comparison to sorghum. *Theoretical and Applied Genetics* 124: 1325–1338.
- MA, X., E. JENSEN, N. ALEXANDROV, AND M. TROUKHAN. 2012. High resolution genetic mapping by genome sequencing reveals genome duplication and tetraploid genetic structure of the diploid *Miscanthus sinensis*. *PLoS ONE* 7: e33821. <http://dx.plos.org/10.1371/journal.pone.0033821>.
- ROZEN, S., AND H. SKALETSKY. 2000. Primer3 on the WWW for general users and for biologist programmers. In S. Misener and S. A. Krawetz [eds.], *Methods in molecular biology*, vol. 132: Bioinformatics methods and protocols, 365–386. Humana Press, Totowa, New Jersey, USA.
- ZHAO, H., J. YU, F. M. YOU, M. LUO, AND J. PENG. 2011. Transferability of microsatellite markers from *Brachypodium distachyon* to *Miscanthus sinensis*, a potential biomass crop. *Journal of Integrative Plant Biology* 53: 232–245.
- ZHOU, H. F., S. S. LI, AND S. GE. 2011. Development of microsatellite markers for *Miscanthus sinensis* (Poaceae) and cross-amplification in other related species. *American Journal of Botany* 98: e195–e197.

APPENDIX 1. List of all accessions used in the study, source, and herbarium voucher number. All taxa are Andropogoneae subtribe Saccharinae unless indicated otherwise.

Taxon ^a	Source ^b	Voucher ^c
<i>M. sacchariflorus</i> 1	TCD Bot. Gardens	TCD P15
<i>M. sinensis</i> ‘Zebrinus’ 2	TCD Bot. Gardens	TCD P20
<i>M. sinensis</i> ‘Zebrinus’ 3	TCD Bot. Gardens	TCD P21
<i>M. ×giganteus</i> 4	TCD Bot. Gardens	TCD P34
<i>M. ×giganteus</i> 5	TCD Bot. Gardens	TCD P36
<i>Miscanthus</i> sp. 6	TCD Bot. Gardens	Tea-6
<i>M. sinensis</i> 7	TCD Bot. Gardens	TCD P48
<i>Miscanthus</i> sp. 8	TCD Bot. Gardens	TCD P50
<i>M. sinensis</i> 9	TCD Bot. Gardens	TCD P51
<i>M. sacchariflorus</i> 10	TCD Bot. Gardens	TCD P58
<i>Miscanthus</i> sp. 11	TCD Bot. Gardens	Tea-11
<i>M. sinensis</i> 13	TCD Bot. Gardens	TCD P73
<i>M. sinensis</i> 14	TCD Bot. Gardens	TCD P75
<i>Miscanthus</i> sp. 15	TCD Bot. Gardens	TCD P104
<i>M. transmorrisonensis</i> 16	TCD Bot. Gardens	TCD P105
<i>M. ×giganteus</i> 17	TCD Bot. Gardens	TCD P108
<i>Miscanthus</i> sp. 18	TCD Bot. Gardens	Tea-18
<i>M. sinensis</i> ‘Goliath’ 19	TCD Bot. Gardens	TCD P110, SIN-H6
<i>M. ×giganteus</i> 20	TCD Bot. Gardens	TCD P114
<i>Miscanthus</i> sp. 21	TCD Bot. Gardens	Tea-21
<i>Miscanthus</i> sp. 22	TCD Bot. Gardens	Tea-22
<i>Miscanthus</i> sp. 23	TCD Bot. Gardens	Tea-23
<i>M. sinensis</i> 24	TCD Bot. Gardens	TCD P11
<i>M. sinensis</i> 25	TCD Bot. Gardens	TCD P11
<i>M. sinensis</i> 26	TCD Bot. Gardens	TCD P11
<i>Miscanthus</i> sp. 27	TCD Bot. Gardens	Tea-27
<i>Miscanthus</i> sp. 28	TCD Bot. Gardens	Tea-28
<i>Miscanthus</i> sp. 29	TCD Bot. Gardens	Tea-29
<i>M. sinensis</i> 30	TCD Bot. Gardens	Tea-30
<i>M. ×giganteus</i> 31	TCD Bot. Gardens	Tea-31
<i>M. ×giganteus</i> 32	TCD Bot. Gardens	Tea-32
<i>M. sinensis</i> ‘Zebrinus’ 33	TCD Bot. Gardens	TCD P20
<i>Miscanthus</i> sp. 34	TCD Bot. Gardens	Tea-34
<i>M. sinensis</i> ‘Gross Fontane’ 35	TCD Bot. Gardens	TCD P30
<i>M. sinensis</i> ‘Gross Fontane’ 36	TCD Bot. Gardens	Tea-36
<i>Miscanthus</i> sp. 37	TCD Bot. Gardens	Tea-37
<i>Miscanthus</i> sp. 38	TCD Bot. Gardens	Tea-38
<i>Miscanthus</i> sp. 39	TCD Bot. Gardens	Tea-39
<i>M. sinensis</i> 40	TCD Bot. Gardens	TCD P62
<i>Miscanthus</i> sp. 42	TCD Bot. Gardens	Tea-42
<i>Miscanthus</i> sp. 43	TCD Bot. Gardens	Tea-43
<i>M. sinensis</i> subsp. <i>condensatus</i> 44	TCD Bot. Gardens	TCD P94
<i>Miscanthus</i> sp. 45	TCD Bot. Gardens	Tea-45
<i>Miscanthus</i> sp. 46	TCD Bot. Gardens	Tea-46
<i>Miscanthus</i> sp. 47	TCD Bot. Gardens	Tea-47
<i>Miscanthus</i> sp. 48	TCD Bot. Gardens	Tea-48
<i>Miscanthus</i> sp. 49	TCD Bot. Gardens	Tea-49
<i>Miscanthus</i> sp. 50	TCD Bot. Gardens	Tea-50
<i>Miscanthus</i> sp. 51	TCD Bot. Gardens	Tea-51
<i>Miscanthus</i> sp. 52	TCD Bot. Gardens	Tea-52
<i>Miscanthus</i> sp. 53	TCD Bot. Gardens	Tea-53
<i>Miscanthus</i> sp. 54	TCD Bot. Gardens	Tea-54
<i>Miscanthus</i> sp. 55	TCD Bot. Gardens	Tea-55
<i>M. sinensis</i> ‘Goliath’ 56	Teagasc Oak Park	Tea-56
<i>M. sinensis</i> ‘Goliath’ 57	TCD Bot. Gardens	Tea-57
<i>M. sinensis</i> ‘Sirene’ 58	Teagasc Oak Park	Tea-58
<i>M. sinensis</i> ‘Strictus’ 59	TRH garden	Tea-59
<i>M. sinensis</i> ‘Strictus’ 60	TCD Bot. Gardens	Tea-60
<i>M. sinensis</i> ‘Malapartus’ 61	TRH Garden	Tea-61
<i>M. sinensis</i> 62	TRH Garden	Tea-62
<i>M. sinensis</i> ‘Sirene’ 63	TCD Bot. Gardens	Tea-63
<i>M. ×giganteus</i> 64	TCD Bot. Gardens	Tea-64
<i>M. ×giganteus</i> 65	TCD Bot. Gardens	Tea-65
<i>M. ×giganteus</i> 66	TRH Garden	Tea-66
<i>Miscanthus</i> sp. 68	TCD Bot. Gardens	Tea-68
<i>Miscanthus</i> sp. 69	TCD Bot. Gardens	Tea-69
<i>Miscanthus</i> sp. 70	TCD Bot. Gardens	Tea-70
<i>Miscanthus</i> sp. 71	TCD Bot. Gardens	Tea-71

APPENDIX 1. Continued.

Taxon ^a	Source ^b	Voucher ^c
<i>Miscanthus</i> sp. 72	TCD Bot. Gardens	Tea-72
<i>Miscanthus</i> sp. 73	TCD Bot. Gardens	Tea-73
<i>M. ×giganteus</i> 74	Germany—from Denmark	Tea-M1 Lasei 1
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 75	Germany	Tea-M81 RH 81
<i>M. sinensis</i> 76	Germany—from Japan	Tea-88-110
<i>M. sinensis</i> 77	Germany—from Japan	Tea-88-111
<i>M. sinensis</i> 78	Germany—from Japan	Tea-90-5
<i>M. sinensis</i> 79	Germany—from Japan	Tea-90-6
<i>M. sinensis</i> 80	Germany—from Sweden	Tea-SW 217
<i>M. ×giganteus</i> 81	Germany—from Denmark	Tea-M53 IPL 53
<i>M. ×giganteus</i> 82	Germany	Tea-M56 HAGA 56
<i>M. ×giganteus</i> 83	Germany	Tea-M63 GREIF 63
<i>M. sacchariflorus</i> 84	Germany—from Japan	Tea-M11 MATEREC 11
<i>M. sinensis</i> 'Goliath' 85	Germany	Tea-M7 GOFAL 7
<i>M. sinensis</i> hybrid 86	Germany	Tea-M42 BERBO 42
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 87	Germany	Tea-M43RH43
<i>M. sinensis</i> hybrid 88	Germany	Tea-M78 JESEL 78
<i>Miscanthus</i> sp. 89	Oak Park	Tea-89
<i>Miscanthus</i> sp. 90	Oak Park	Tea-90
<i>Miscanthus</i> sp. 91	Oak Park	Tea-91
<i>Miscanthus</i> sp. 92	Oak Park	Tea-92
<i>M. ×giganteus</i> 93	IGER/TinPlant/Oak Park	Tea-93
<i>M. ×giganteus</i> 94	Old Trial Teagasc Oak Park	Tea-94
<i>M. sinensis</i> 95	Sweden	Tea-95
<i>M. sinensis</i> 96	Sweden	Tea-96
<i>M. sinensis</i> 97	Sweden	Tea-97
<i>M. sinensis</i> 98	Sweden	Tea-98
<i>M. sinensis</i> 99	Sweden	Tea-99
<i>M. sinensis</i> 100	Sweden	Tea-100
<i>M. sinensis</i> 101	Sweden	Tea-101
<i>M. sinensis</i> 102	Sweden	Tea-102
<i>M. sinensis</i> 103	Sweden	Tea-103
<i>M. sinensis</i> 104	Sweden	Tea-104
<i>M. sinensis</i> 105	Sweden	Tea-105
<i>M. sinensis</i> 106	Sweden	Tea-106
<i>M. sinensis</i> 107	Sweden	Tea-107
<i>M. sinensis</i> 108	Sweden	Tea-108
<i>M. sinensis</i> 109	Sweden	Tea-109
<i>M. sinensis</i> 110	Sweden	Tea-110
<i>M. sinensis</i> 111	Sweden	Tea-111
<i>M. sinensis</i> 112	Sweden	Tea-112
<i>M. sinensis</i> 113	Sweden	Tea-113
<i>M. sinensis</i> 114	Sweden	Tea-114
<i>M. sinensis</i> 115	Sweden	Tea-115
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 116	Sweden	Tea-116
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 117	Sweden	Tea-117
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 118	Sweden	Tea-118
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 119	Sweden	Tea-119
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 120	Sweden	Tea-120
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 121	Sweden	Tea-121
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 122	Sweden	Tea-122
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 123	Sweden	Tea-123
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 124	Sweden	Tea-124
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 125	Sweden	Tea-125
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 126	Sweden	Tea-126
<i>M. sacchariflorus</i> × <i>M. sinensis</i> 127	Sweden	Tea-127
<i>M. sacchariflorus</i> 128	TCD Bot. Gardens	Tea-128
<i>M. sacchariflorus</i> 129	TCD Bot. Gardens	Tea-129
<i>Miscanthus</i> sp. 130	TCD Bot. Gardens	Tea-130
<i>Miscanthus</i> sp. 131	TCD Bot. Gardens	Tea-131
<i>Saccharum officinarum</i>	TCD Bot. Gardens	TCD TRH s.n.
<i>Cymbopogon citratus</i> ^d	TCD Bot. Gardens	TCD TRH s.n.
<i>Zea diploperennis</i> ^e	TCD Bot. Gardens	TCD TRH s.n.
<i>Sorghum halepense</i> 6 ^f	RBG Kew 151 01	Kew 1966-54209
<i>Pennisetum</i> sp. ^g	TCD Bot. Gardens	TCD TRH s.n.
<i>M. sinensis</i> var. <i>variegatus</i> 1	RBG Kew 154 04	Kew 1969-19093
<i>M. sinensis</i> subsp. <i>condensatus</i> 7	RBG Kew 151	Kew 1969-19091
<i>M. oligostachyus</i> 16	RBG Kew 151 (pot)	Kew 1995-1864
<i>M. nepalensis</i> 25	RBG Kew TH 4	Kew 1985-8388

APPENDIX 1. Continued.

Taxon ^a	Source ^b	Voucher ^c
<i>M. sinensis</i> 'Goliath' 27	ADAS Steinmann nurseries	Kew MB93/02
<i>M. sinensis</i> 'Gracillimus' 28	ADAS Piccoplant, Germany	Kew MB94/05
<i>M. sinensis</i> 'Roland' 29	ADAS Piccoplant, Germany	Kew MB94/06
<i>M. sinensis</i> Anderss. 30	ADAS Wye College	Kew MB94/07
<i>M. sinensis</i> 'Gross Fontane' 31	ADAS Genft Dogels, Germany	Kew PN95/01
<i>M. sacchariflorus</i> 61	RBG Kew	Kew 1987-2727
<i>M. sinensis</i> 'Yakushimanum' 63	RBG Kew	Kew 1987-1148
<i>M. transmorrisonensis</i> 65	RBG Kew	Kew 1990-2748
<i>M. fuscus</i> 82	RBG Kew	Kew 590
<i>M. violaceus</i> 84	RBG Kew	Kew 7437
<i>M. ecklonii</i> 86	RBG Kew	Kew 2347
<i>M. junceus</i> 88	RBG Kew	Kew 1060
<i>M. junceus</i> 89	RBG Kew	Kew 2309
<i>M. ecklonii</i> 105	RBG Kew	Kew 2929
<i>M. ecklonii</i> 106	RBG Kew	Kew 247
<i>M. yunnanensis</i> 107	RBG Kew	Kew 30689
<i>M. nudipes</i> 109	RBG Kew	Kew 2007
<i>M. tinctorius</i> 112	RBG Kew	Kew 1466
<i>Saccharum spontaneum</i> 117	RBG Kew	Kew Butt, 1977
<i>Narenga porphyrocoma</i> 120	RBG Kew	Kew 2092
<i>Saccharum contortum</i> 121	RBG Kew	Kew 3797
<i>Spodiopogon rhizophorus</i> 125	RBG Kew	Kew 283
<i>Spodiopogon sibiricus</i> 128	RBG Kew	Kew 210
<i>Eulalia quadrinervis</i> 134	RBG Kew	Kew 3294
<i>M. sinensis</i> 'Morning Light' 155	RBG Kew	Kew 1996 821
<i>M. sacchariflorus</i> 159	RBG Kew	Kew 3598 1935
<i>M. sacchariflorus</i> 160	RBG Kew	Kew 1984
<i>M. tinctorius</i> 'Nana Variegata' 161	RBG Kew	Kew 1996 1065
<i>M. sinensis</i> 'Goliath' 194	ADAS	Kew PN96/30

^a Numbers accompanying species names represent the DNA extraction identifier for this study.

^b Source abbreviations: ADAS = Agricultural Development Advisory Service (now Agriculture and Environmental Consultancy); IGER = Institute of Grassland and Environmental Research (now Institute of Biological, Environmental and Rural Sciences [IBERS]); RBG Kew = Royal Botanic Gardens, Kew, Richmond, Surrey, United Kingdom; TCD Bot. Gardens = Trinity College Dublin Botanical Garden, Dublin, Ireland; Teagasc Oak Park = Teagasc Oak Park Research Centre, Carlow, Ireland; TRH Garden = personal garden of first author.

^c Voucher abbreviations: Kew = Herbarium of the Royal Botanic Gardens, Kew, Richmond, Surrey, United Kingdom; TCD = Trinity College Dublin Herbarium, Ireland; Tea = Teagasc Oak Park Research Centre, Carlow, Ireland.

^d Andropogoninae, Andropogoneae (subtribe/tribe).

^e Tripsacinae, Andropogoneae.

^f Sorghinae, Andropogoneae.

^g Cenchrinae, Paniceae.