



## **Microsatellite Primers in *Parietaria judaica* (Urticaceae) to Assess Genetic Diversity and Structure in Urban Landscapes**

Authors: Bossu, Angèle, Bertaudière-Montès, Valérie, Dubut, Vincent, and Manel, Stéphanie

Source: Applications in Plant Sciences, 2(9)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1400036>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## MICROSATELLITE PRIMERS IN *PARIETARIA JUDAICA* (URTICACEAE) TO ASSESS GENETIC DIVERSITY AND STRUCTURE IN URBAN LANDSCAPES<sup>1</sup>

ANGÈLE BOSSU<sup>2,5</sup>, VALÉRIE BERTAUDIÈRE-MONTÈS<sup>2</sup>, VINCENT DUBUT<sup>3</sup>, AND STÉPHANIE MANEL<sup>2,4</sup>

<sup>2</sup>Aix Marseille Université, Institut de recherche pour le développement (IRD), Laboratoire Population Environnement Développement (LPEd) UMR\_D151, 13331 Marseille, France; <sup>3</sup>Aix Marseille Université, Centre national de la recherche scientifique (CNRS), IRD, Univ. Avignon, IMBE—UMR7263, 13331 Marseille CEDEX 3, France; and <sup>4</sup>Centre de coopération internationale en recherche agronomique pour le développement (CIRAD), UMR AMAP, TA A51/PS2, 34398 Montpellier CEDEX 5, France

- **Premise of the study:** Urbanization is one of the main factors contributing to loss of genetic diversity, as the resulting landscape fragmentation and habitat loss induce species isolation. However, studies of genetic structure and diversity in urbanized landscapes are still rare. We characterized microsatellite primers for *Parietaria judaica* to study this environment.
- **Methods and Results:** Eleven microsatellite loci from *P. judaica*, an urban plant, were isolated using shotgun pyrosequencing, and the simple sequence repeat (SSR) markers were screened in 20 individuals of *P. judaica*. The loci were tested on 166 individuals from three populations in different cities. The number of alleles ranged from two to 19, and expected and observed heterozygosity values ranged from 0.019 to 0.912 and 0.019 to 0.448, respectively.
- **Conclusions:** The markers amplified well in the species and will be useful for examining genetic diversity and population genetic structure in this urban plant.

**Key words:** microsatellites; *Parietaria judaica*; urban area; Urticaceae.

*Parietaria judaica* L. is a perennial herb, with individual plants consisting of many shoots emerging from a common rootstock. The plant is a wind-pollinated species of the Urticaceae family that grows in urban Mediterranean areas (Fotiou et al., 2011) and is very common on urban walls. This wide occurrence will facilitate studies of genetic diversity and genetic structure in urban environments. While urbanization is one of the main factors contributing to loss of genetic biodiversity, genetic patterns in urban areas have been understudied (Manel and Holderegger, 2013). In the current study, we have developed and characterized 11 microsatellite markers for *P. judaica* that will be used for further studies to examine genetic diversity of plants in urban environments.

### METHODS AND RESULTS

Twenty individuals of *P. judaica* were collected in Marseilles, France (43.296346°N, 5.369889°E; voucher specimen available at the Herbarium of

<sup>1</sup>Manuscript received 18 April 2014; revision accepted 18 June 2014.

This research was carried out as part of a Ph.D. thesis (PACA regional research grant no. 2011-019) and financially supported by the Agence Nationale de la Recherche (ANR). This research is a contribution to the “Trame Verte Urbaine” (TVU, NVD08\_321105) program entitled “Assessment of urban green corridors and development of a reference frame: An infrastructure between aesthetic and ecology for a new urbanity.” S.M. was supported by the Institut Universitaire de France.

<sup>5</sup>Author for correspondence: angele.bossu@etu.univ-amu.fr

doi:10.3732/apps.1400036

Muséum d’Histoire Naturelle d’Aix-en-Provence [AIX], barcode number AIX036092). Samples were sent to Genoscreen (Lille, France) to develop and characterize microsatellites. Total genomic DNA was isolated from individuals’ leaf tissues using NucleoSpin Plant II (Macherey-Nagel, Duren, Germany). A total of 1 µg of DNA was used for the development of microsatellite libraries through 454 GS FLX Titanium pyrosequencing of enriched DNA libraries as described in Malausa et al. (2011). Total DNA was enriched for TG, TC, AAC, AAG, AGG, ACG, ACAT, and ACTC repeat motifs and subsequently amplified. PCR products were purified, quantified, and GS FLX libraries were then constructed following the manufacturer’s protocols (Roche Diagnostics, Meylan, France) and sequenced on a GS FLX PicoTiterPlate Kit (Roche Diagnostics). Sequences with a minimum of five repeats were selected. The bioinformatics program QDD version 1 with default parameters (Megléc et al., 2010) was used to filter for redundancy, resulting in a final set of sequences from which it was possible to design primers. Finally, for a total of 43.9 Mbp read in the run, 10,895 sequences comprised microsatellite motifs with an average length of 304 bp. A total of 1114 primer sets were designed with QDD using BLAST, ClustalW, and Primer3 (Altschul et al., 1990; Larkin et al., 2007; Rozen and Skaletsky, 2000). A subgroup of 47 primer pairs was then tested for amplification. Primer sets were discarded if they failed to amplify or led to multiple fragments.

Each primer pair was assessed in eight individuals of *P. judaica*. Microsatellite loci were amplified in 25-µL reactions containing 20 ng of template DNA, 1 unit of *Taq* polymerase (FastStart *Taq* DNA polymerase, Roche Diagnostics), 0.6 mM of dNTPs, 3.75 mM of MgCl<sub>2</sub>, and 0.1 µM of each forward and reverse primer. The amplification profiles included initial denaturation at 95°C for 10 min; followed by 40 cycles of denaturation at 95°C for 30 s, annealing at 55°C for 30 s, and extension at 72°C for 1 min; followed by a final extension at 72°C for 10 min. For each PCR, a negative control was included. A total of 2 µL of PCR product was separated on 2% agarose gel. Primer pairs that amplified in at least five individuals were validated. Among 47 primer pairs tested, 26 pairs were validated on a subset of eight individuals from Marseilles, France. Polymorphism of 24 microsatellite loci was tested on the same subset of individuals

TABLE 1. Characteristics of the microsatellite loci designed for *Parietaria judaica*.

Locus	Primer sequences (5'–3')	$T_a$ (°C)	Repeat motif	Fluorescent dye	Allele size (bp)	GenBank accession no.
ParJ_20	F: AGTTCAGTTGGGTTGGCAG R: TATGGGTGTGAGTGGCTACG	56 56	(AG) <sub>11</sub>	PET	146	KJ747956
ParJ_26	F: CCTCCAATCGTTGAGAGGAA R: CCGCCATTGCTGTCTAGC	54 56	(GAA) <sub>8</sub>	PET	157	KJ747957
ParJ_27	F: TCCGATTATCCTCTTCGCAT R: GCAGAGAAATGAAAGAGAGTCCG	53 53	(TTC) <sub>12</sub>	6-FAM	159	KJ789106
ParJ_31	F: CTTACGCGCCACGACTACAAC R: TCAAAAATCGAGATTGGCTCA	56 52	(GTAT) <sub>5</sub>	PET	184	KJ789103
ParJ_33	F: AGCCGTTAAGGTCCCAAAGT R: GGGACATGCGAAAGAAAAGA	57 53	(TTC) <sub>18</sub>	VIC	190	KJ872579
ParJ_34	F: AACGGAATTCATTCTCCGA R: TTGATTCTCCAAACACAACG	52 51	(TGA) <sub>10</sub>	NED	190	KJ872580
ParJ_37 <sup>a</sup>	F: GAAGCCTCGTCGAACCAAGT R: CGACACCTTCACCAACAAGA	57 55	(TC) <sub>9</sub>	6-FAM	201	
ParJ_42	F: CGTATTTTGGCGGAAAGCTA R: CCCATTCGTGAAGTTGGTAGA	53 54	(CT) <sub>10</sub>	6-FAM	235	KJ789107
ParJ_43	F: GTGGCTCTCGACTCGGAC R: AACTCGAGAGCTTCATCGTTTC	58 55	(CTT) <sub>14</sub>	VIC	240	KJ789105
ParJ_44	F: GACAGTGTCCCTATTGAGCG R: TCGGGAGTTGGAACATATCA	56 53	(CTT) <sub>11</sub>	NED	243	KJ789108
ParJ_45	F: GTAGGAAAAGGGTTTGTCTCG R: TTTCTCTGATGTGGAAGCCC	53 55	(CT) <sub>10</sub>	PET	245	KJ789104

Note:  $T_a$  = annealing temperature.

<sup>a</sup>No GenBank accession number is available for locus ParJ\_37 because the region between the primer and the repeat region has too many undetermined nucleotides.

that was used for amplification tests. PCR cycles and mixture were the same as those mentioned previously except that we used 10 pmol of each forward and reverse primer (instead of 1 pmol). One microliter of each PCR product was run in an ABI 3730xl DNA Analyzer (Applied Biosystems, Waltham, Massachusetts, USA). Results were analyzed with GeneMapper 4.0 (Applied Biosystems).

Eleven markers were selected, and amplifications were tested for individuals sampled from three different cities: 85 individuals in Marseilles (43.296346°N, 5.369889°E), 29 individuals in Lauris (43.747778°N, 5.313611°E), and 52 individuals in Aix-en-Provence (43.531127°N, 5.454025°E).

The characteristics of the 11 novel microsatellite loci in *P. judaica* are summarized in Table 1. We calculated the expected heterozygosity using an R package (adegenet version 1.3-9.2; Jombart and Ahmed, 2011). The number of alleles per locus ranged from two to 13 in Lauris, two to 15 in Aix-en-Provence, and three to 16 in Marseilles (Table 2). Expected heterozygosity ranged from 0.128 to 0.874, 0.019 to 0.882, and 0.240 to 0.912 in Lauris, Aix-en-Provence, and Marseilles, respectively. Observed heterozygosity ranged from 0.068 to 0.448, 0.019 to 0.231, and 0.059 to 0.423 in

Lauris, Aix-en-Provence, and Marseilles, respectively. All loci showed significant linkage disequilibrium (except loci ParJ\_31 in the Lauris population and ParJ\_37 in the Lauris and Aix-en-Provence populations).

## CONCLUSIONS

The molecular markers described here are the first microsatellite loci isolated for *P. judaica*. Most markers showed median levels of polymorphism, while ParJ\_33 and ParJ\_43 showed high levels of polymorphism. Most loci were at Hardy–Weinberg disequilibrium potentially due to genetic drift caused by landscape fragmentation. The set of microsatellite markers will be useful for studying population genetics of *P. judaica* in urban environments.

TABLE 2. Polymorphism in 11 microsatellite loci developed for three populations of *Parietaria judaica*.<sup>a</sup>

Locus	Total		Lauris (N = 29)			Aix-en-Provence (N = 52)			Marseilles (N = 84)		
	N	A	A	$H_e$	$H_o$	A	$H_e$	$H_o$	A	$H_e$	$H_o$
ParJ_20	166	8	5	0.610 <sup>b</sup>	0.345	6	0.712 <sup>b</sup>	0.212	7	0.636 <sup>b</sup>	0.294
ParJ_26	166	7	3	0.324 <sup>b</sup>	0.069	4	0.627 <sup>b</sup>	0.173	7	0.654 <sup>b</sup>	0.317
ParJ_27	166	7	4	0.509 <sup>b</sup>	0.310	4	0.620 <sup>b</sup>	0.180	7	0.636 <sup>b</sup>	0.321
ParJ_31	166	4	2	0.328	0.207	2	0.158 <sup>b</sup>	0.019	4	0.628 <sup>b</sup>	0.282
ParJ_33	166	19	13	0.874 <sup>b</sup>	0.448	15	0.868 <sup>b</sup>	0.212	16	0.912 <sup>b</sup>	0.423
ParJ_34	166	11	6	0.561 <sup>b</sup>	0.310	6	0.592 <sup>b</sup>	0.173	9	0.551 <sup>b</sup>	0.294
ParJ_37	166	3	2	0.128	0.068	2	0.019	0.019	3	0.315 <sup>b</sup>	0.072
ParJ_42	166	6	4	0.604 <sup>b</sup>	0.138	4	0.580 <sup>b</sup>	0.137	6	0.661 <sup>b</sup>	0.256
ParJ_43	166	19	9	0.815 <sup>b</sup>	0.345	15	0.882 <sup>b</sup>	0.231	15	0.695 <sup>b</sup>	0.289
ParJ_44	166	10	4	0.329 <sup>b</sup>	0.138	7	0.633 <sup>b</sup>	0.154	8	0.650 <sup>b</sup>	0.376
ParJ_45	166	4	3	0.558 <sup>b</sup>	0.172	2	0.464 <sup>b</sup>	0.115	4	0.240 <sup>b</sup>	0.059

Note: A = number of alleles;  $H_e$  = expected heterozygosity;  $H_o$  = observed heterozygosity; N = total number of samples analyzed.

<sup>a</sup>Geographic coordinates of populations used in the study: Lauris (43.747778°N, 5.313611°E), Aix-en-Provence (43.531127°N, 5.454025°E), Marseilles (43.296346°N, 5.369889°E).

<sup>b</sup>Markers deviating from Hardy–Weinberg equilibrium after Bonferroni correction ( $P < 0.004$ ).

#### LITERATURE CITED

- ALTSCHUL, S. F., W. GISH, W. MILLER, E. W. MYERS, AND D. J. LIPMAN. 1990. Basic local alignment search tool. *Journal of Molecular Biology* 215: 403–410.
- FOTIOU, C., A. DAMIALIS, N. KRIGAS, J. M. HALLEY, AND D. VOKOU. 2011. *Parietaria judaica* flowering phenology, pollen production, viability and atmospheric circulation, and expansive ability in the urban environment: Impacts of environmental factors. *International Journal of Biometeorology* 55: 35–50.
- JOMBART, T., AND I. AHMED. 2011. adegenet 1.3-1: New tools for the analysis of genome-wide SNP data. *Bioinformatics (Oxford, England)* 27: 3070–3071.
- LARKIN, M. A., G. BLACKSHIELDS, N. P. BROWN, R. CHENNA, P. A. McGETTIGAN, H. McWILLIAM, F. VALENTIN, ET AL. 2007. Clustal W and Clustal X version 2.0. *Bioinformatics (Oxford, England)* 23: 2947–2948.
- MALAUSSA, T., A. GILLES, E. MEGLÉCZ, H. BLANQUART, S. DUTHOY, C. COSTEDOAT, V. DUBUT, ET AL. 2011. High-throughput microsatellite isolation through 454 GS-FLX Titanium pyrosequencing of enriched DNA libraries. *Molecular Ecology Resources* 11: 638–644.
- MANEL, S., AND R. HOLDEREGGER. 2013. Ten years of landscape genetics. *Trends in Ecology & Evolution* 28: 614–621.
- MEGLÉCZ, E., C. COSTEDOAT, V. DUBUT, A. GILLES, T. MALAUSSA, N. PECH, AND J.-F. MARTIN. 2010. QDD: A user-friendly program to select microsatellite markers and design primers from large sequencing projects. *Bioinformatics (Oxford, England)* 26: 403–404.
- ROZEN, S., AND H. SKALETSKY. 2000. Primer3 on the WWW for general users and for biologist programmers. In S. Misener and S. A. Krawetz [eds.], *Methods in molecular biology*, vol. 132: *Bioinformatics methods and protocols*, 365–386. Humana Press, Totowa, New Jersey, USA.