

Development of Highly Variable Microsatellite Markers for the Tetraploid *Silene stellata* (Caryophyllaceae)

Authors: Zhou, Juannan, Dudash, Michele R., Fenster, Charles B., and Zimmer, Elizabeth A.

Source: Applications in Plant Sciences, 4(12)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1600117>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

DEVELOPMENT OF HIGHLY VARIABLE MICROSATELLITE MARKERS FOR THE TETRAPLOID *SILENE STELLATA* (CARYOPHYLLACEAE)¹

JUANNAN ZHOU^{2,3,6}, MICHELE R. DUDASH⁴, CHARLES B. FENSTER⁵, AND ELIZABETH A. ZIMMER³

²Behavior, Ecology, Evolution and Systematics Graduate Program, University of Maryland, College Park, Maryland 20742 USA;

³Department of Botany, National Museum of Natural History, MRC 166, Smithsonian Institution, Washington, DC 20013-7012 USA; ⁴Department of Natural Resource Management, South Dakota State University, Brookings, South Dakota 57007 USA; and

⁵Department of Biology and Microbiology, South Dakota State University, Brookings, South Dakota 57007 USA

- **Premise of the study:** We designed and tested microsatellite markers for the North American native species *Silene stellata* (Caryophyllaceae) to investigate its population genetic structure and identify selection on floral design through male reproductive success.
- **Methods and Results:** A total of 153 candidate microsatellite loci were isolated based on next-generation sequencing. We identified 18 polymorphic microsatellite loci in three populations of *S. stellata*, with di- or trinucleotide repeats. Genotyping results showed the number of alleles per locus ranged from six to 45 and expected heterozygosity ranged from 0.511 to 0.951. Five of these loci were successfully amplified in *S. virginica* and *S. caroliniana* and were also polymorphic.
- **Conclusions:** The microsatellite markers reported here provide a valuable tool for paternity analysis in *S. stellata*. They will also be useful for investigating the population genetic structures of *S. stellata* and related species.

Key words: Caryophyllaceae; microsatellites; plant–animal interaction; polyploidy; *Silene stellata*.

There are nearly 700 species in the genus *Silene* L. (Caryophyllaceae). *Silene* species exhibit diverse pollination specialization as manifested by their floral diversity (Kephart et al., 2006). Approximately one third of all *Silene* species in the Old and New World exhibit nocturnal pollination syndromes and usually form close interactions with noctuid moths from the genus *Hadena* (Noctuidae) (Kephart et al., 2006). *Silene stellata* (L.) W. T. Aiton is an infrequent native perennial herb that is distributed throughout the eastern part of the United States. The flowers are pollinated by *Hadena ectypa* as well as by a number of generalist nocturnal moths (Reynolds et al., 2009; Kula et al., 2013). In addition to the positive effect of pollination, oviposition by female *H. ectypa* inside the calyx results in strong negative effects through larval predation on the reproductive tissues of *S. stellata* (Kula et al., 2013). Depending on the amount of pollination service provided by the generalist moths, the net outcome of the *Silene*–*Hadena* interaction can range from mutualism to parasitism (Reynolds et al., 2012), making it a valuable system for understanding the evolutionary dynamics of interspecific interactions (Kephart et al., 2006; Bernasconi et al., 2009).

Ecological and evolutionary studies utilizing genetic markers of North American *Silene* species are relatively few compared

with the European species (Moyle, 2006), potentially due to technical complications caused by the prevalent polyploidy: most North American *Silene* species have been shown to be polyploids, including tetra-, hexa-, and octoploids, with tetraploidy being the most common ploidy level (Popp and Oxelman, 2007). Prior to this study, we tested nine microsatellite markers developed for *S. latifolia* Poir. (Magalhaes et al., 2011) in 20 individuals of *S. stellata* and did not identify any polymorphic loci. We report the development of 18 highly variable microsatellite markers for the tetraploid *S. stellata*. These markers are being used to quantify the population genetic structure of *S. stellata*. We will also use these markers to assess individual siring abilities through paternity assignment to investigate selection on floral design through the male function of the hermaphroditic *S. stellata*. We also tested transferability of these markers to two closely related tetraploid *Silene* species: the specialist hummingbird-pollinated *S. virginica* L. and the primarily *Bombus*- and hawk moth-pollinated *S. caroliniana* Walter (Reynolds et al., 2009).

METHODS AND RESULTS

Genomic DNA was extracted from fresh leaf tissue of a *S. stellata* individual collected from a natural population near Mountain Lake Biological Station in Giles County, Virginia, USA (37.348296°N, 80.544301°W, elevation ca. 1100–1300 m; Appendix 1), using the DNeasy Plant Mini Kit (QIAGEN, Valencia, California, USA) following the manufacturer's protocol. DNA libraries were prepared using the KAPA Library Preparation Kit version 2.13 (KR0453; Kapa Biosystems, Wilmington, Massachusetts, USA) following the manufacturer's protocol. Libraries were quantified with a Qubit Fluorometric Quantitation instrument (Life Technologies, Carlsbad, California, USA) and then sequenced using an Illumina MiSeq version 3 kit (Illumina, San Diego, California, USA) to produce paired-end reads of ≤301 bases. A total of 8,557,438 reads were imported and paired by name in Geneious 7.0.6 (Biomatters, Auckland,

¹Manuscript received 19 September 2016; revision accepted 27 October 2016.

The authors thank G. Bernasconi, D. Taylor, and their laboratory groups for initial help in microsatellite marker development, and T. Glenn and his laboratory group for help in next-generation sequencing and primer design of *Silene stellata*.

⁶Author for correspondence: jnzhou@umd.edu

doi:10.3732/apps.1600117

TABLE 1. Characteristics of 18 microsatellite loci developed in *Silene stellata*.^a

Locus	Primer sequences (5'–3')	Fluorescent dye	Repeat motif	Allele size range (bp)	T _a (°C)	GenBank accession no.
SS1010	F: ACACAGCGGATGGTGTCTG R: CTCTCTTTCAATCCTGGCCC	HEX	(AG) ₂₆	174–248	60	KX712873
SS62979	F: CTTTGGGTGTGTGCTCAGAATAG R: TTGTGTCTGCCATCCTTGC	HEX	(AG) ₈	155–239	55	KX712874
SS63719	F: GTACGCGTCGCTTTGTTAAC R: TGATACTGCTCACCGGATGC	FAM	(AT) ₈	135–154	58	KX712875
SS64450	F: CAAGTGGTATCAGAGCCAGG R: ACCTGTAACCTTGTGGGATGC	FAM	(AT) ₉	220–254	60	KX712876
SS64692	F: TGGTACCCTCAAGACTAGGC R: CCTCCCTGAACGCTCTTC	HEX	(AAT) ₁₄	148–265	60	KX712877
SS68587	F: TCAGGTCAATAGGCTGGGG R: CTGGGTGGTTGATGAAATTTGG	HEX	(AAC) ₈	191–281	55	KX712878
SS71250	F: TAGACGGTTCGCGAATGGG R: TTTCTCACAACTCACGCTC	FAM	(AG) ₁₄	125–203	58	KX712879
SS73434	F: GAGAAGGGAAGACCGGTG R: TGACCCGAGACCTACTGATTG	FAM	(AG) ₁₀	184–210	60	KX712880
SS78777	F: ACGTACAGTAGCATAGGTGGG R: TCAGGTCATAGTAGGCCAC	HEX	(AC) ₈	156–248	60	KX712881
SS85007	F: CAGGGTCAACATCAACAGGC R: AGATTGTCTCTCGGCTCCC	FAM	(AAT) ₄₁	149–356	60	KX712882
SS89178	F: TCACTACCCACCTTTACGG R: ATCCCTAGATGCCCGCAC	FAM	(AG) ₈	254–374	55	KX712883
SS100056	F: GCATCTTGTGTTCCCGAGC R: GGGCTATCTTGATCTCCACTC	FAM	(AAG) ₃₇	91–205	60	KX712884
SS115569	F: CCAGCTATCCCAACGAATC R: ACCCAACCAACGAATCAAAC	HEX	(AAC) ₁₅	377–399	55	KX712885
SS122722	F: ACACAATGTTTTCTGACTTGCG R: ATTGTATCTCGGCTGCCAC	FAM	(AAT) ₁₅	176–287	60	KX712886
SS129528	F: AGTGACGGATACCATGGAGC R: CATCCAATCCGCTTTCTCTC	HEX	(AAG) ₃₁	96–226	60	KX712887
SS131968	F: ACCCAACATACATCAACCTCTC R: GAGCGGTTGAGAATGGTGTG	FAM	(AAT) ₉	154–205	55	KX712888
SS131990	F: TCAGCTACCCACCAATTTCTTG R: CCCACCCTTAATACGTACACC	HEX	(AT) ₁₉	186–282	55	KX712889
SS141347	F: GGAAGAATGGAGGTGCTC R: GAGAGATCAGACCGAGGCTG	HEX	(ACC) ₁₆	132–192	60	KX712890

Note: T_a = annealing temperature.

^aAll values are based on 95 samples representing three populations in Virginia.

New Zealand). A de novo assembly was performed on the first 1,000,000 sequences for which both reads of any pair were ≥200 bases.

A total of 99,506 consensus sequences between 200 and 400 bp were extracted and screened for potential microsatellite loci using MSATCOMMANDER 1.0.8 beta (Faircloth, 2008) with default settings. Primers were designed for sequences with perfect di-, tri-, and tetranucleotide repeats in Primer3 software (Rozen and Skaletsky, 1999). We designed primers for 153 out of a total of 946 loci identified as containing microsatellites. One hundred fourteen primer pairs were first tested with seven randomly selected *S. stellata* samples for amplification success and polymorphism. For each primer pair, we modified the forward primer by attaching a CAG tag (5'-CAGTCGGGCGTCATCA-3') preceding the 5' end to enable the cost-efficient fluorescent labeling system of PCR products described by Boutin-Ganache et al. (2001) and Glenn (2001). Ten microliter PCR reactions were performed using the QIAGEN Type-it Microsatellite PCR Kit. Each reaction contained the following components: ~10 ng of genomic DNA, 5 μL of the 2× Multiplex PCR Master Mix, 0.02 μM of the modified forward primer, 0.2 μM of reverse primer, and 0.2 μM of the fluorescently labeled CATG primer (5'-6FAM, 5'-HEX). A touchdown PCR protocol was used to test all primer pairs: 5 min of denaturing at 95°C; five cycles of 95°C for 30 s, 60°C for 1.5 min, and 72°C for 30 s; followed by 28 cycles of 95°C for 30 s, 55°C for 1.5 min, and 72°C for 30 s; and a final extension at 60°C for 30 min.

Of the 114 primer pairs tested, 50 produced bands consistently on an agarose gel. Amplicons of these primers were analyzed using an ABI Prism 3730 Genetic Analyzer (Applied Biosystems, Foster City, California, USA), then visualized and scored in Geneious 7.0.6 (Biomatters). Eighteen loci showed clear peak patterns and were polymorphic (Table 1). These 18 polymorphic loci were further characterized using a total of 95 *S. stellata* individuals collected from three local populations near Mountain Lake Biological Station (Meadow,

Woodland, and Windrock) within 8 km of one another. To investigate marker transferability, these 18 loci were also tested on six individuals from one population each of *S. virginica* from Newport, Virginia, and *S. caroliniana* from Potomac, Maryland (Appendix 1).

We report the following parameters for the three populations of *S. stellata*: sample size, number of alleles, number of private alleles, observed heterozygosity, and expected heterozygosity. The parameters were estimated using GenoDive version 2.0b27 (Meirmans and Van Tienderen, 2004) while correcting for unknown dosage of alleles for partial heterozygotes. Across the three populations, the number of alleles ranged from six to 45, and expected heterozygosity ranged from 0.511 to 0.951 (Table 2).

Of the 18 loci tested in *S. virginica* and *S. caroliniana*, 10 loci were successfully amplified in both species. Genotyping results showed two loci were monomorphic in both species, one locus was polymorphic in *S. virginica* but monomorphic in *S. caroliniana*, five loci were polymorphic in both species, and two loci showed multiple bands (Table 3). Vouchers for the *Silene* species were deposited at the Norton-Brown Herbarium (University of Maryland, College Park, Maryland, USA; Appendix 1).

CONCLUSIONS

We developed 18 novel microsatellite loci for *S. stellata*. These loci showed high variability in *S. stellata* and therefore are suitable for future paternity analysis. Five of these markers are polymorphic in the related *S. virginica* and *S. caroliniana*. These microsatellites will also be useful for studying the

TABLE 2. Genetic diversity of the 18 polymorphic microsatellites of *Silene stellata*.^a

Locus	Total (n = 95)			Meadow (n = 55)				Windrock (n = 20)				Woodland (n = 20)			
	A	H _o	H _e	A	A _p	H _o	H _e	A	A _p	H _o	H _e	A	A _p	H _o	H _e
SS1010	31	0.847	0.894	28	10	0.824	0.903	17	0	0.840	0.912	19	2	0.886	0.876
SS62979	36	0.828	0.913	28	12	0.664	0.925	17	4	0.823	0.918	15	4	0.996	0.894
SS63719	18	0.468	0.877	14	2	0.527	0.858	12	2	0.364	0.891	12	2	0.512	0.885
SS64450	15	0.486	0.712	10	4	0.393	0.626	7	2	0.474	0.724	9	3	0.591	0.782
SS64692	38	0.707	0.906	29	11	0.658	0.909	17	4	0.712	0.879	19	4	0.737	0.915
SS68587	27	0.866	0.932	25	5	0.883	0.931	17	1	0.835	0.932	16	1	0.879	0.933
SS71250	26	0.999	0.918	24	4	0.999	0.921	21	1	0.989	0.912	18	1	1.010	0.922
SS73434	13	0.210	0.875	10	2	0.305	0.890	7	2	0.154	0.842	9	0	0.172	0.889
SS78777	34	0.892	0.919	33	12	0.975	0.931	17	0	0.770	0.911	17	1	0.933	0.915
SS85007	45	0.621	0.921	38	17	0.716	0.922	18	3	0.585	0.926	20	3	0.555	0.902
SS89178	37	0.826	0.887	26	9	0.903	0.866	19	5	0.677	0.866	18	3	0.931	0.951
SS100056	35	0.821	0.951	33	8	0.799	0.961	22	1	0.883	0.943	21	1	0.781	0.948
SS115569	6	0.394	0.511	6	4	0.458	0.551	2	0	0.397	0.538	2	0	0.328	0.442
SS122722	38	0.733	0.944	31	9	0.807	0.957	18	4	0.547	0.916	19	2	0.856	0.959
SS129528	35	0.896	0.937	29	7	0.825	0.924	20	3	0.982	0.933	23	2	0.883	0.954
SS131968	18	0.779	0.761	17	3	0.699	0.730	14	1	0.760	0.704	11	0	0.869	0.844
SS131990	30	0.440	0.939	27	8	0.458	0.937	15	0	0.407	0.943	15	3	0.455	0.937
SS141347	17	0.993	0.891	16	2	0.998	0.884	15	0	0.989	0.895	14	1	0.990	0.891

Note: A = number of alleles; A_p = number of private alleles; H_e = expected heterozygosity; H_o = observed heterozygosity; n = number of individuals sampled.

^aLocality and voucher information are provided in Appendix 1.

population genetics of *S. stellata* and related North American species in this genus.

LITERATURE CITED

- BERNASCONI, G., J. ANTONOVICS, A. BIÈRE, D. CHARLESWORTH, L. F. DELPH, D. FILATOV, T. GIRAUD, ET AL. 2009. *Silene* as a model system in ecology and evolution. *Heredity* 103: 5–14.
- BOUTIN-GANACHE, I., M. RAPOSO, M. RAYMOND, AND C. F. DESCHÉPPER. 2001. M13-tailed primers improve the readability and usability of microsatellite analyses performed with two different allele-sizing methods. *BioTechniques* 31: 24–28.
- FAIRCLOTH, B. C. 2008. MSATCOMMANDER: Detection of microsatellite repeat arrays and automated, locus-specific primer design. *Molecular Ecology Resources* 8: 92–94.
- GLENN, T. C. 2001. 5' tags: A cheap way to fluorescently tag PCR products. Website <http://dna.uga.edu/wp-content/uploads/sites/11/2013/12/5-Primer-Tags-PCR-products.pdf> [accessed 21 November 2016].
- KEPHART, S., R. J. REYNOLDS, M. T. RUTTER, C. B. FENSTER, AND M. R. DUDASH. 2006. Pollination and seed predation by moths on *Silene* and allied Caryophyllaceae: Evaluating a model system to study the evolution of mutualisms. *New Phytologist* 169: 667–680.
- KULA, A. A. R., M. R. DUDASH, AND C. B. FENSTER. 2013. Choices and consequences of oviposition by a pollinating seed predator, *Hadena ectypa* (Noctuidae), on its host plant, *Silene stellata* (Caryophyllaceae). *American Journal of Botany* 100: 1148–1154.
- MAGALHAES, I. S., G. GLEISER, A.-M. LABOUCHE, AND G. BERNASCONI. 2011. Comparative population genetic structure in a plant–pollinator/seed predator system. *Molecular Ecology* 20: 4618–4630.
- MEIRMANS, P. G., AND P. H. VAN TIENDEREN. 2004. GENOTYPE and GENODIVE: Two programs for the analysis of genetic diversity of asexual organisms. *Molecular Ecology Notes* 4: 792–794.
- MOYLE, L. C. 2006. Correlates of genetic differentiation and isolation by distance in 17 congeneric *Silene* species. *Molecular Ecology* 15: 1067–1081.
- POPP, M., AND B. OXELMAN. 2007. Origin and evolution of North American polyploid *Silene* (Caryophyllaceae). *American Journal of Botany* 94: 330–349.
- REYNOLDS, R. J., M. J. WESTBROOK, A. S. ROHDE, J. M. CRIDLAND, C. B. FENSTER, AND M. R. DUDASH. 2009. Pollinator specialization and pollination syndromes of three related North American *Silene*. *Ecology* 90: 2077–2087.
- REYNOLDS, R. J., A. A. R. KULA, C. B. FENSTER, AND M. R. DUDASH. 2012. Variable nursery pollinator importance and its effect on plant reproductive success. *Oecologia* 168: 439–448.
- ROZEN, S., AND H. SKALETSKY. 1999. Primer3 on the WWW for general users and for biologist programmers. In S. Misener and S. A. Krawetz [eds.], *Methods in molecular biology*, vol. 132: Bioinformatics: Methods and protocols, 365–386. Humana Press, Totowa, New Jersey, USA.

TABLE 3. Genetic diversity of six microsatellite loci developed in *Silene stellata* in two related native *Silene* species.^a

Locus	Allele size range (bp)	<i>S. virginica</i> (n = 6)			<i>S. caroliniana</i> (n = 6)			
		A	H _o	H _e	Allele size range (bp)	A	H _o	H _e
SS64692	154–196	7	1	0.94	151–214	8	1	0.861
SS68587	188–191	2	0.5	0.369	161–191	5	0.667	0.614
SS71250	113–129	5	0.833	0.754	117–165	6	1	0.662
SS78777	198–254	11	0.833	0.969	184–244	12	1	0.939
SS131968 ^b	—	—	—	—	157–169	3	0.333	0.489
SS141347	144–180	9	0.5	0.867	147–168	7	1	0.826

Note: A = number of alleles; H_e = expected heterozygosity; H_o = observed heterozygosity; n = number of individuals sampled.

^aLocality and voucher information are provided in Appendix 1.

^bLocus SS131968 was monomorphic in *S. virginica*.

APPENDIX 1. Geographic location and voucher information of *Silene* populations used in this study.

Species	Location	Population	Geographic coordinates	Voucher no. ^a
<i>S. stellata</i> (L.) W. T. Aiton ^b	Newport, VA, USA	Meadow	37.348296°N, 80.544301°W	MARY1012978
<i>S. stellata</i>	Newport, VA, USA	Meadow	37.348296°N, 80.544301°W	MARY1012979
<i>S. stellata</i>	Newport, VA, USA	Windrock	37.413889°N, 80.519444°W	MARY1012974
<i>S. stellata</i>	Newport, VA, USA	Windrock	37.413889°N, 80.519444°W	MARY1012975
<i>S. stellata</i>	Newport, VA, USA	Woodland	37.355415°N, 80.553469°W	MARY1012977
<i>S. stellata</i>	Newport, VA, USA	Woodland	37.355415°N, 80.553469°W	MARY1012978
<i>S. caroliniana</i> Walter	Potomac, MD, USA	Great Falls	38.989511°N, 77.246713°W	MARY1012980
<i>S. caroliniana</i>	Potomac, MD, USA	Great Falls	38.989511°N, 77.246713°W	MARY1012981
<i>S. virginica</i> L.	Newport, VA, USA	Windrock	37.413889°N, 80.519444°W	MARY1012982
<i>S. virginica</i>	Newport, VA, USA	Windrock	37.413889°N, 80.519444°W	MARY1012983

^aHerbarium vouchers were deposited at the University of Maryland Herbarium (MARY).

^bIndividual used for DNA extraction.