

Development of Polymorphic Microsatellite Markers for the Okinawa Pit Viper, *Ovophis okinavensis*

Authors: Kadota, Yohei, Kurita, Kazuki, and Inoue, Eiji

Source: Current Herpetology, 38(2) : 169-172

Published By: The Herpetological Society of Japan

URL: <https://doi.org/10.5358/hsj.38.169>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Development of Polymorphic Microsatellite Markers for the Okinawa Pit Viper, *Ovophis okinavensis*

YOHEI KADOTA¹, KAZUKI KURITA^{2*}, AND EIJI INOUE³

¹Department of Zoology, Graduate School of Science, Kyoto University, Sakyo, Kyoto, 606–8502, JAPAN

²Division of Forest and Biomaterials Science, Graduate School of Agriculture, Kyoto University, Sakyo, Koyo, 606–8502, JAPAN

³Department of Biology, Faculty of Science, Toho University, Funabashi City, Chiba, 274–8510, JAPAN

Abstract: We isolated and characterized 11 microsatellite markers for *Ovophis okinavensis*, a viperid species endemic to the Ryukyu Archipelago. These markers showed polymorphism among 25 individuals from Okinawajima Island. The number of alleles per locus was 3–11, and observed and expected heterozygosity were 0.240–0.960 and 0.218–0.853, respectively. Cross-amplification confirmed that at least seven out of the 11 microsatellite markers were applicable to a putative relative, *Trimeresurus gracilis*. The markers provided here are expected to be valuable for population- and/or individual-level genetic studies of not only *O. okinavensis*, but also its relatives.

Key words: *Ovophis okinavensis*; Primer sequence; Short tandem repeat; *Trimeresurus gracilis*; Viperidae

INTRODUCTION

Ovophis okinavensis is a short, stout-bodied viperid snake that inhabits forest areas, especially near streams, ponds, and marshes, on many subtropical islands of the Okinawa and Amami groups, Ryukyu Archipelago, Japan. This snake is a typical ambush forager and exhibits a foraging strategy that is adjusted to spatial and temporal fluctuations of the emergence of two frog species with different breeding periods (Kadota, 2011).

The specialized foraging behavior together with its broad geographic range suggests that this snake has a heterogeneous population structure. However, there are currently no useful genetic markers (including those of the relatives) to obtain fine-scale genetic information on this species. Here, we report newly developed 11 microsatellite primers of *O. okinavensis* and examine the suitability of these markers for *Trimeresurus gracilis*, which is the putative sister taxon to *O. okinavensis* (Castoe and Parkinson, 2006; Malhotra et al., 2010).

* Corresponding author. Tel: +81–75–753–6129;
FAX: +81–75–753–6129;
E-mail address: kurita@zoo.zool.kyoto-u.ac.jp

MATERIALS AND METHODS

We isolated microsatellite loci from ventral

TABLE 1. Characteristics of 11 microsatellite loci developed for *Ovophis okinavensis* and the results of cross-amplification in *Trimeresurus gracilis*.

Locus (Accession no.)	Primer sequence (5'-3')	Repeat motif	T _a	<i>Ovophis okinavensis</i> (n=25)			<i>Trimeresurus gracilis</i> (n=4)				
				Size range (bp)	A	H _O	H _E	Size range (bp)	A	H _O	H _E
OvoP1 (LC414977)	F: GCCCATCGACTTTGTTTGC R: GGTTTTCTTGCCCTCAGCAC	(AC) ₂₃	64	168-192	10	0.96	0.85	168	1	0.00	0.00
OvoP7 (LC414978)	F: GGTGCATACAACCATCAACAGG R: GAACAACGGAGGCAAGAAG	(AC) ₁₂	64	243-249	4	0.68	0.56	240-252	3	0.50	0.41
OvoP11 (LC414979)	F: TCTGACATAGTGGAGGATGG R: GTGCCACAGGCAATGTTTT	(GT) ₁₃	62	198-204	4	0.24	0.22	191-204	4	0.75	0.72
OvoP12 (LC414980)	F: AAAGAGGCAGTGGAGCATGT R: GATGAGACTGAGAGGATAGGG	(TG) ₁₂	64	248-264	4	0.64	0.7	256-268	5	1.00	0.78
OvoP17 (LC414981)	F: GTGGAAAGAAAGTAAGCATTG R: ATGAAAACCCATGGAAGGTG	(GT) ₈	62	305-321	5	0.52	0.43	318-328	4	0.25	0.72
OvoP24 (LC414982)	F: AAGGGACATGTGTGTGTG R: CGAGTGCGGTAICTTCTTCC	(TG) ₁₃	62	208-212	3	0.56	0.65	190-208	3	0.25	0.59
OvoP29 (LC414983)	F: CACATACACACACGCACAGG R: TGGGGCAAGCTGACTTAAC	(AC) ₉	64	200-210	5	0.56	0.71	191-217	4	1.00	0.69
OvoP30 (LC414984)	F: GGTTGTCCAATTCCTTCTGG R: GTTTGGCCCAAAATAGCC	(GT) ₁₈	64	258-266	5	0.76	0.66	—	—	—	—
OvoP38 (LC414985)	F: GTTACACATTGCTCGCTTGC R: CTTCCTGGTGTACAGGTTTT	(CA) ₂₂	62	265-383	11	0.88	0.83	—	—	—	—
OvoP40 (LC414986)	F: CCTCTTCACTCCACCCAGTC R: CGCTAGCGACCCAAAGTTAC	(TG) ₁₆	62	306-320	8	0.56	0.66	311-312	2	—*	—*
OvoP45 (LC414987)	F: AAAAACAGTCGGGTTTGTGG R: CCAGTCTCCCTTTAICTCTTTGA	(TG) ₁₂	62	248-266	4	0.36	0.33	250-260	5	0.5	0.75

T_a, annealing temperature (°C); A, number of alleles; H_O, observed heterozygosity; H_E, expected heterozygosity; * only one individual successfully amplified.

scale tissue samples from Okinawajima Island snakes using the method slightly modified from Glenn and Schable (2005). The detailed methods for library preparation and primer design are described in Kurita et al. (2013). For amplification trials, 25 individuals of *O. okinavensis* from the northern part of Okinawajima Island were used (DNA identification numbers by Y. Kadota: 4, 100–102, 106, 107, 111, 114, 117, 128, 129, 133, 135, 138, 139, 145, 149, 151, 152, 154, 158, 164, 170, 175, 177). In addition, we tested cross-species amplification of developed markers on four individuals of *T. gracilis* from Taiwan (temporal specimen numbers by Ming-Chung Tu: 79804–805 and 79807–808). PCR amplification was conducted using fluorescently (FAM, HEX, NED) labeled M13(-21) universal primers (Schuelke, 2000) with the TaKaRa Ex Taq kit (Takara Bio, Otsu, Japan). Fragment sizes were measured with GeneScan 400HD ROX size standard (Applied Biosystems, Foster City, CA, USA) on ABI 3130xl Genetic Analyzer (Applied Biosystems) and analyzed by the Peak Scanner software (Applied Biosystems). Number of alleles (A), and observed (H_o) and expected heterozygosities (H_e) were calculated using GenAlEx 6.503 (Peakall and Smouse, 2006, 2012). Deviation from Hardy–Weinberg equilibrium and linkage disequilibrium between loci were tested using GENEPOP 4.7.0 (Rousset, 2008). Null allele frequency was estimated using CERVUS 3.0.7 (Kalinowski et al., 2007).

RESULTS AND DISCUSSION

The characteristics of the 11 isolated microsatellite loci are summarized in Table 1. These loci exhibited high or moderate allelic polymorphism in *O. okinavensis*. The number of alleles per locus ranged from 3 to 11. The observed and expected heterozygosity ranged from 0.240 to 0.960 and from 0.218 to 0.853, respectively. There was no evidence of deviations from the Hardy–Weinberg equilibrium at any locus and significant linkage disequilibrium at any pair of loci after the sequential

Bonferroni correction (Rice, 1989). The estimated frequency of null alleles ranged from -0.13 to 0.12 . In *T. gracilis*, eight of 11 loci were successfully amplified for all individuals examined, and seven loci showed polymorphism. As such, the microsatellite markers developed here will be useful for population- and/or individual-level genetic studies of *O. okinavensis* and its relatives, in the fields of ecology and population genetics (e.g. population structure, genetic variation, demography, and relatedness analyses including paternity).

ACKNOWLEDGMENTS

We would like to thank M.-C. Tu and C.-F. Lin for collecting and providing valuable tissue samples of *T. gracilis*. The samples were from protected animals in Taiwan. They were permitted to collect by Dr. Tu who obtained his permission from Council of Agriculture, Executive Yuan; and T. Hikida and T. Okamoto for their technical support. We also thank anonymous reviewers for their comments in improving our manuscript. This research was financially supported in part by the Global COE Program A06 to Kyoto University.

LITERATURE CITED

- CASTOE, T. A. AND PARKINSON, C. L. 2006. Bayesian mixed models and the phylogeny of pitvipers (Viperidae: Serpentes). *Molecular Phylogenetics and Evolution* 39: 91–110.
- GLENN, T. C. AND SCHABLE, N. A. 2005. Isolating microsatellite DNA loci. *Methods in Enzymology* 395: 202–222.
- KADOTA, Y. 2011. Is *Ovophis okinavensis* active only in the cool season? Temporal foraging pattern of a subtropical pit viper in Okinawa, Japan. *Zoological Studies* 50: 269–275.
- KALINOWSKI, S. T., TAPER, M. L., AND MARSHALL, T. C. 2007. Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology* 16: 1099–1106.
- KURITA, K., HIKIDA, T., AND TODA, M. 2013.

- Development and characterization of polymorphic microsatellite marker for East Asian species of the genus *Plestiodon*. *Conservation Genetics Resources* 5: 355–357.
- MALHOTRA, A., CREER, S., POOK, C. E., AND THORPE, R. S. 2010. Inclusion of nuclear intron sequence data helps to identify the Asian sister group of New World pitvipers. *Molecular Phylogenetics and Evolution* 54: 172–178.
- PEAKALL, R. AND SMOUSE, P. E. 2006. GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6: 288–295.
- PEAKALL, R. AND SMOUSE, P. E. 2012. GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics* 28: 2537–2539.
- RICE, W. R. 1989. Analyzing tables of statistical tests. *Evolution* 43: 223–225.
- ROUSSET, R. 2008. GENEPOP'007: a complete re-implementation of the genepop software for Windows and Linux. *Molecular Ecology Resources* 8: 103–106.
- SCHUELKE, M. 2000. An economic method for the fluorescent labeling of PCR fragments. *Nature Biotechnology* 18: 233–234.
-
- Accepted: 7 May 2019*