

## **PARASITES OF WATERFOWL FROM SOUTHWEST TEXAS: I. THE NORTHERN CINNAMON TEAL, *Anas cyanoptera septentrionalis***

Authors: WILKINSON, JAMES N., CANARIS, ALBERT G., and BRODERSON, DOROTHY

Source: Journal of Wildlife Diseases, 13(1) : 62-63

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-13.1.62>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## PARASITES OF WATERFOWL FROM SOUTHWEST TEXAS: I. THE NORTHERN CINNAMON TEAL, *Anas cyanoptera septentrionalium*

JAMES N. WILKINSON, ALBERT G. CANARIS and DOROTHY BRODERSON,  
Department of Biological Sciences, The University of Texas at El Paso,  
El Paso, Texas 79968, USA

**Abstract:** Seventeen northern cinnamon teal, *Anas cyanoptera septentrionalium* were collected in Hudspeth County, Texas: 10 in January, 1974, and seven in September, 1974. Fifteen parasite species were recovered: 11 were new host records. Three species of parasites were recorded from spring migrants and 15 from fall migrants.

### INTRODUCTION

The northern cinnamon teal, *Anas cyanoptera septentrionalium*, is a sub-species native to western North America. The bird is an early fall and spring migrant. Although not abundant, some utilize the Rio Grande Valley in El Paso and Hudspeth Counties, Texas, on their annual migrations.

Not much is known about parasites of waterfowl in the Southwest and even less about the parasites of the northern cinnamon teal. McDonald<sup>3,4,5,6</sup> and Larios<sup>2</sup> have published references to parasites from this duck.

### MATERIALS AND METHODS

Seventeen cinnamon teal were collected in the Rio Grande Valley, Hudspeth County, Texas; 10 spring migrants returning from the South in January, 1974, and seven fall migrants in September, 1974. Ducks were placed in individual plastic bags and on ice immediately after collecting. Those that could not be examined within six hrs were frozen and examined later.

Trematodes, cestodes, and acanthocephalons were fixed in Alcohol-Formalin-Acetic Acid, stained in Harris' hematoxylin, and mounted in Canada balsam.

### Acknowledgment

We wish to thank Dr. Malcolm E. McDonald for his assistance in identification of helminth parasites.

Nematodes and ectoparasites were preserved in 70% ethanol and mounted in lactophenol and Hoyer's, respectively.

### RESULTS AND DISCUSSION

All 17 northern cinnamon teal were infected with at least one species of parasite. Of the 15 species of parasites recorded only three were recorded from the spring migrants while all 15 were recorded from the fall migrants.

The most common parasites were *Cloacotaenia megalops* (88%), *Echinostoma revolutum* (47%), and *Trichobilharzia physellae* (41%), (Table 1).

Data presented by Buscher<sup>1</sup> for the blue-winged teal (*Anas discors* L.) and Turner and Threlfall<sup>7</sup> for the blue-winged teal and green-winged teal (*Anas crecca* L.) indicated that the juvenile ducks were infected with more species of helminths than the adults. The fall cinnamon teal in our study consisted of one adult and six juveniles. Although the sample was small, the greater number of species of parasites observed in the fall migrants may have been influenced by the preponderance of juveniles.

All parasites recorded in this study have been reported from various species of North American ducks, but 11 are new host records for cinnamon teal.

TABLE 1. Parasites of the Northern Cinnamon Teal, *Anas cyanoptera septentrionalium*, from Southwest Texas.

	Spring (10 Ducks) No. Infected	Fall (7 Ducks) No. Infected
<b>TREMATODA</b>		
* <i>Echinostoma revolutum</i>	6	2
<i>Notocotylus attenuatus</i>	—	2
* <i>Trichobilharzia physellae</i>	4	3
* <i>Bilharziella polonica</i>	—	2
<b>CESTODA</b>		
<i>Cloacotaenia megalops</i>	8	7
* <i>Microsomacanthus hopkinsi</i>	—	2
* <i>Sobolevicanthus filumferens</i>	—	1
* <i>Echinocotyle</i> sp.	—	1
* <i>Diorchis nyrocae</i>	—	3
<b>NEMATODA</b>		
<i>Capillaria anatis</i>	—	2
<i>Streptocara crassicauda</i>	—	1
<b>ACANTHOCEPHALA</b>		
* <i>Corynosoma constrictum</i>	—	1
* <i>Fillicollis anatis</i>	—	2
<b>MALLOPHAGA</b>		
* <i>Trinoton querquedulae</i>	—	3
* <i>Anaticola crasicornis</i>	—	1

\*New host records.

**LITERATURE CITED**

- BUSCHER, H. N. 1966. Intestinal helminths of the blue-winged teal, *Anas discors* L. at Delta, Manitoba, Can. J. Zool. 44: 113-116.
- LARIOS, I. 1944. Description de un Cestoda del Genero *Hymenolepis* Encontrado en los Patos Silvestres del hago de Texcoco, Mexico. An. Biol. Inst. Univ. Nac. Mexico 15: 73-78.
- MCDONALD, M. E. 1969. Annotated Bibliography of Helminths of Waterfowl (Anatidae). Bureau of Sport Fisheries and Wildlife. Special scientific report. Wildlife No. 125. Washington, D.C.
- . 1969. Catalog of Helminths of Waterfowl (Anatidae). Bureau of Sport Fisheries and Wildlife. Special scientific report. Wildlife No. 126. Washington, D.C.
- . 1974. Key to Nematodes Reported in Waterfowl. Department of the Interior, Bureau of Sport Fisheries and Wildlife. Resource Publication 122, Washington, D.C.
- . 1974. Nematode Parasites of Waterfowl (Anseriformes) from Western United States. U.S. Fish and Wildlife Service. Bear River Research Station. 19 p.
- TURNER, B. C. and W. THRELFALL. 1976. The metazoan parasites of green-winged teal (*Anas crecca* L.) and blue-winged teal (*Anas discors* L.) from Eastern Canada. Proc. Helm. Soc. Wash. 42: 157-159.

Received for publication 30 March 1976