

## **SEASONAL TRENDS OF HELMINTH PARASITES OF BOBWHITE QUAIL 1**

Authors: DAVIDSON, WILLIAM R., KELLOGG, FOREST E., and DOSTER, GARY L.

Source: Journal of Wildlife Diseases, 16(3) : 367-375

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-16.3.367>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## SEASONAL TRENDS OF HELMINTH PARASITES OF BOBWHITE QUAIL <sup>□</sup>

WILLIAM R. DAVIDSON, FOREST E. KELLOGG and GARY L. DOSTER, Southeastern Cooperative Wildlife Disease Study, Department of Parasitology, College of Veterinary Medicine, University of Georgia, Athens, Georgia 30602, USA.

**Abstract:** Helminthologic examination of 120 adult and 65 juvenile bobwhites (*Colinus virginianus*) from Leon County, Florida, during a one-year period revealed seven common (>30% prevalence) species including *Raillietina cesticillus*, *R. colinia*, *Cheilospirura spinosa*, *Cyrnea colini*, *Heterakis bonasae*, *Tetrameres pattersoni*, and *Trichostrongylus tenuis*. Less frequently found helminths included *Hymenolepis* sp., *Rhabdometra odiosa*, *Mediorhynchus papillosus*, *Aproctella stoddardi*, *Dispharynx nasuta*, *Gongylonema ingluvicola*, *Strongyloides avium*, and *Subulura* sp. Juvenile bobwhites had acquired infections of 6 of the 7 common helminths by July and all seven species by August. A shift from a predominance of immature to mature parasites was noted with increasing age of juvenile bobwhites. Patterns of acquisition of common helminths by juvenile bobwhites followed both linear and non-linear (plateau effect) trends when compared to age of the host. By mid-winter total helminth burdens of juvenile birds approached levels in adults. *Cheilospirura spinosa*, *C. colini* and *T. pattersoni* showed marked peaks in transmission between June and September. The two cecal nematodes, *H. bonasae* and *T. tenuis*, showed seasonal shifts in relative abundance with *H. bonasae* predominating during the summer and *T. tenuis* predominating during the winter. Lesions attributable to helminths were rare and involved minimal tissue damage.

### INTRODUCTION

The popularity of the bobwhite (*Colinus virginianus*) as a game bird has promoted considerable research on this species. Numerous parasitologic studies have been published on bobwhites and have revealed that they harbor a great variety and number of helminths.<sup>1,2,3,12,14,18,20,21</sup>

Previous investigations, however, have provided little information on the seasonal aspects of helminth infections in these birds. In the present study, bobwhites were collected on a monthly basis for 12 consecutive months in a high-density population<sup>13</sup> to determine seasonal trends in helminth infections.

### METHODS

The study was conducted on a 447 ha area of Tall Timbers Research Station, which is located in a limestone region of broken terrain in Leon County, Florida. The topography, vegetation, and bobwhite population density were described previously.<sup>13</sup>

Ten adult birds were shot each month from March, 1968, through February, 1969. Five juvenile birds were obtained during July of 1968, and 10 juvenile birds per month were collected from August, 1968, through January, 1969. All birds were collected between the 10th and 20th of each month. Not more than one adult and one immature bobwhite per covey

<sup>□</sup> This study was supported by Tall Timbers Research, Inc., Tallahassee, Florida, and by an appropriation from the Congress of the United States. Funds were administered and research coordinated under the Federal Aid in Wildlife Restoration Act (50 Stat. 917) and through Contract Nos. 14-16-0008-676 and 14-16-0008-209, Fish and Wildlife Service, U.S. Department of the Interior.

were utilized, and an attempt was made to obtain equal numbers of male and female birds.

Bobwhites were tagged immediately after death, placed in individual plastic bags, and chilled until examined later in the day. Procedures for examination of the viscera and identification of the parasites were those used by Kellogg and Prestwood.<sup>14</sup> When feasible, helminths were classified as immature or mature stages.

Monthly parasite burdens were examined for changes that could be attributed to time of collection by calculation of rank correlation coefficients and comparison with critical  $r$  values. Prevalence data were tested for differences attributable to stratifying parameters by the Chi square test.

## RESULTS AND DISCUSSION

One hundred and twenty adult and 65 juvenile bobwhites were examined for helminths during the one-year period.

Four cestode, one acanthocephalan, and 10 nematode species were found (Table 1). All of these species have been found previously in bobwhites from several localities except for *Raillietina colinia* which has been reported only in Texas.<sup>11,12,19,20</sup> Specimens from this study which earlier were described as *Mediorhynchus bakeri*<sup>3</sup> have been synonymized with *M. papillosis*.<sup>17</sup> Re-examination of specimens collected previously from Tall Timbers Research Station and reported as *Raillietina* sp. and *Tetrameres americana*<sup>14</sup> revealed these helminths to be *R. colinia* and *T. pattersoni*.

All birds harbored at least one species of helminth, and seven species, *R. cesticillus*, *R. colinia*, *Cheilospirura spinosa*, *Cyrnea colini*, *Heterakis bonasae*, *Tetrameres pattersoni*, and *Trichostrongylus tenuis*, were considered to be common (>30% prevalence) in bobwhites on the study area. The remaining species were judged to be infrequent or to represent accidental in-

TABLE 1. Helminths recovered from 120 adult and 65 juvenile bobwhites from Tall Timbers Research Station, Leon County, Florida.

Helminth	Adult Bobwhites	Juvenile Bobwhites
<b>CESTODA</b>		
<i>Hymenolepis</i> sp.	1 (1) 1*	—
<i>Raillietina cesticillus</i>	32 (14) 1-137	32 (20) 1-124
<i>Raillietina colinia</i>	43 (8) 1-98	52 (6) 1-40
<i>Rhabdometra odiosa</i>	3 (4) 2-8	8 (3) 1-8
<b>ACANTHOCEPHALA</b>		
<i>Mediorhynchus papillosis</i>	1 (2) 2	—
<b>NEMATODA</b>		
<i>Aproctella stoddardi</i>	3 (1) 1	2 (5) 5
<i>Cheilospirura spinosa</i>	72 (5) 1-35	45 (3) 1-26
<i>Cyrnea colini</i>	72 (4) 1-21	85 (5) 1-18
<i>Dispharynx nasuta</i>	2 (1) 1	9 (1) 1
<i>Gongylonema ingluvicola</i>	2 (1) 1	—
<i>Heterakis bonasae</i>	100 (47) 3-198	97 (20) 1-86
<i>Strongyloides avium</i>	22 (5) 1-17	5 (5) 1-10
<i>Subulura</i> sp.	1 (1) 1	2 (1) 1
<i>Tetrameres pattersoni</i>	78 (7) 1-58	71 (10) 1-65
<i>Trichostrongylus tenuis</i>	98 (59) 1-361	42 (16) 1-123

\*Percent prevalence (average/per infected bird) range.

fections. The monthly prevalences and intensities of common helminths in juvenile and adult bobwhites are presented in Tables 2 and 3.

By July, juvenile bobwhites had acquired infections of all common helminths except *R. cesticillus* which was first detected in August. During July, infections with *C. spinosa*, *C. colini*, *H. bonasae*, and *T. pattersoni* in juvenile bobwhites were comprised solely or primarily of immature helminths, but thereafter the proportion of immature helminths was lower with infections comprised mainly of mature helminths

(Fig. 1). This shift from a predominance of immature to mature helminths was attributed to the time required for helminth growth and maturation. Some reported periods required for maturation of these or closely related species are: *C. spinosa*, 45 days;<sup>4</sup> *C. colini*, 41 days;<sup>4</sup> *T. americana*, 45 days;<sup>4</sup> and *H. bonasae*, 27 days.<sup>6</sup>

From July to August, juvenile bobwhites had sharp increases in the prevalence and intensity of all mature helminths except *T. tenuis*, but thereafter the pattern of acquisition of helminth populations differed among




FIGURE 1. The relative abundance (% of population) of immature *Cheilospirura spinosa*, *Cryneae colini*, *Tetrameres pattersoni* and *Heterakis bonasae* in juvenile bobwhites during 7 consecutive months.

TABLE 2. Monthly prevalences and intensities of common helminths in 65 juvenile bobwhites.

Month Number examined	Jul 5	Aug 10	Sep 10	Oct 10	Nov 10	Dec 10	Jan 10
<b>HELMINTH</b>							
<i>Railletina cesticillus</i>	0 (0)*	40 (40)	30 (10)	40 (5)	40 (6)	40 (46)	20 (20)
<i>Railletina colinia</i>	40 (4)	60 (6)	70 (4)	80 (6)	30 (17)	40 (3)	40 (6)
<i>Cheilosporira spinosa</i> (ad.)†	0 (0)	40 (7)	40 (3)	30 (3)	30 (1)	50 (2)	30 (2)
<i>Cheilosporira spinosa</i> (im.)	60 (4)	30 (2)	10 (3)	10 (2)	10 (1)	0 (0)	10 (1)
<i>Cyrnea colini</i> (ad.)	20 (5)	60 (8)	90 (4)	60 (8)	90 (3)	90 (4)	100 (3)
<i>Cyrnea colini</i> (im.)	80 (7)	60 (5)	70 (2)	10 (1)	0 (0)	30 (2)	40 (1)
<i>Heterakis bonasae</i> (ad.)	0 (0)	80 (7)	80 (9)	100 (16)	100 (20)	100 (14)	100 (31)
<i>Heterakis bonasae</i> (im.)	60 (2)	50 (3)	70 (2)	90 (25)	30 (7)	20 (2)	10 (2)
<i>Tetrameres pattersoni</i> (ad.)	20 (4)	80 (8)	70 (18)	60 (11)	70 (9)	70 (5)	70 (7)
<i>Tetrameres pattersoni</i> (im.)	40 (8)	0 (0)	40 (3)	10 (1)	20 (2)	0 (0)	30 (2)
<i>Trichostrongylus tenuis</i>	40 (6)	10 (2)	30 (3)	40 (8)	40 (4)	50 (14)	80 (35)

\*Percent prevalence (average per infected bird).

†Designates developmental stage of helminth: ad = adult and im = immature. Those without designation were not classified as to developmental stage.

TABLE 3. Monthly prevalences and intensities of common helminths in 120 adult bobwhites.

Month Number examined	Mar		Apr		May		Jun		Jul		Aug		Sep		Oct		Nov		Dec		Jan		Feb	
	10		10		10		10		10		10		10		10		10		10		10		10	
<b>HELMINTH</b>																								
<i>Railletina cesticillus</i>	30 (3)*		20 (75)		20 (12)		50 (29)		30 (6)		40 (15)		10 (3)		20 (2)		30 (2)		30 (11)		50 (3)		50 (7)	
<i>Railletina colinia</i>	60 (12)		90 (11)		50 (3)		60 (9)		60 (25)		50 (4)		20 (3)		0 (0)		20 (1)		0 (0)		60 (2)		50 (3)	
<i>Cheliosporira spinosa</i> (ad.)†	60 (3)		50 (4)		50 (1)		60 (7)		60 (6)		90 (5)		60 (3)		80 (5)		90 (3)		80 (4)		100 (6)		70 (3)	
<i>Cheliosporira spinosa</i> (im.)	0 (0)		0 (0)		0 (0)		10 (1)		60 (4)		70 (8)		20 (5)		0 (0)		10 (1)		0 (0)		0 (0)		20 (1)	
<i>Cyrnea colini</i> (ad.)	80 (3)		70 (3)		90 (5)		100 (3)		80 (4)		50 (3)		80 (3)		30 (4)		60 (2)		60 (5)		40 (4)		60 (4)	
<i>Cyrnea colini</i> (im.)	10 (1)		0 (0)		20 (1)		40 (1)		40 (4)		30 (6)		70 (1)		10 (2)		20 (1)		30 (2)		10 (1)		0 (0)	
<i>Heterakis bonasae</i> (ad.)	100 (39)		100 (48)		100 (19)		100 (36)		100 (38)		100 (59)		100 (43)		100 (73)		100 (45)		100 (42)		100 (39)		100 (17)	
<i>Heterakis bonasae</i> (im.)	100 (21)		70 (6)		60 (6)		100 (24)		70 (3)		60 (2)		30 (3)		30 (5)		40 (2)		20 (1)		40 (2)		0 (0)	
<i>Tetrameres pattersoni</i> (ad.)	80 (5)		70 (5)		60 (5)		80 (3)		80 (11)		100 (5)		60 (6)		80 (9)		80 (7)		70 (7)		90 (6)		80 (12)	
<i>Tetrameres pattersoni</i> (im.)	0 (0)		0 (0)		0 (0)		60 (1)		40 (9)		0 (0)		10 (1)		10 (2)		10 (1)		0 (0)		10 (1)		10 (1)	
<i>Trichostrongylus tenuis</i>	100 (140)		100 (109)		100 (61)		100 (110)		100 (51)		100 (24)		100 (12)		90 (22)		100 (33)		90 (50)		100 (63)		100 (32)	

\*Percent prevalence (average per infected bird).

†Designates stage of helminth: ad = adult and im = immature. Species without designation were not classified as to developmental stage.

species. *Cyrnea colini*, *H. bonasae*, and *T. tenuis* showed continued increases in prevalence that were correlated ( $P < 0.05$ ) with time, eventually reaching or approaching a prevalence of 100%. In contrast, *C. spinosa*, *T. pattersoni* and *R. cesticillus* exhibited a plateau effect with relatively stable prevalence values from August through January. The prevalence of *R. colinia* showed considerable variation, although a case could be made for a trend of an initial increase followed by a decline to a lower stable level.

As with the prevalence of infection, the intensities of *H. bonasae* and *T. tenuis* showed increases that were correlated ( $P < 0.05$ ) with time, although *T. tenuis* did not increase sharply until December and January. The intensities of the remaining species did not increase with time ( $P < 0.05$ ) but rather exhibited tendencies to (1) peak and decline (*C. spinosa*, *C. colini*, and *T. pattersoni*), (2) remain relatively stable (*R. colinia*), or (3) fluctuate erratically (*R. cesticillus*).

Mean total helminth burdens in juvenile bobwhites increased from 18 in July to 76 in January, although the increase was not significantly correlated with time ( $P < 0.05$ ). The general trend of increasing helminth burdens with age in juvenile birds apparently is a function of length of exposure and was influenced primarily by *H. bonasae* and *T. tenuis* which continued to increase in intensity. By January, the total mean helminth burden in juvenile bobwhites approached levels found in adults.

Each month adult bobwhites harbored infections of all common helminths except for *R. colinia* which was not found in October or December. All differences between adult and juvenile bobwhites in prevalences or intensities of infection with common helminths (Table 1) are attributable to the initial limited exposure of juvenile birds. Adult bobwhites had a significantly lower ( $X^2=5.83$ ;  $df=1$ ) prevalence of *Dispharynx nasuta* and a significantly

higher ( $X^2=9.27$ ;  $df=1$ ) prevalence of *Strongyloides avium* than juvenile bobwhites. Fewer *D. nasuta* in adult bobwhites is consistent with reports for other avian species.<sup>7</sup> The higher prevalence of *S. avium* in adults resulted from the frequent occurrence of the parasite from March to July (before collection of juveniles), and thus, whether this difference was due to host age or to a general decline of *S. avium* on the study area is unknown. The occurrence of two species, *Hymenolepis* sp. and *M. papillosis*, in adult but not juvenile bobwhites was not considered significant since both species probably represent accidental infections.

In adult bobwhites, three species of heteroxenous (intermediate host required) helminths, *C. spinosa*, *C. colini*, and *T. pattersoni*, had distinct peaks in the proportion of immature helminths in the population (Fig. 2). These single annual peaks in acquisition of larvae indicate that transmission of these species occurs primarily during the summer. Summer peaks in transmission

have obvious survival advantages for these species since this time coincides with an abundance of arthropod intermediate hosts and entrance of juvenile bobwhites into the population. The diet of young bobwhite chicks consists mainly of animal foods<sup>8</sup> and hence chances of infection with heteroxenous helminths are enhanced.

The cestodes, *R. cesticillus* and *R. colinia*, showed erratic fluctuations in prevalence and intensity without any indication of peaks in transmission or seasonal trends in the levels of infection. Analyses of crop contents of birds collected during this study (Whitcomb, unpubl.) revealed three genera of beetles (*Anisodactylus*, *Harpalus*, and *Selenophorus*) which are known intermediate hosts of *R. cesticillus*.<sup>15</sup> The genera *Anisodactylus* and *Selenophorus* accounted for 85% of the arthropods recovered from May through August and probably are important intermediate hosts of *R. cesticillus* on the study area.

In adult bobwhites, *H. bonasae* and *T. tenuis* differed from all other helminths




FIGURE 2. The relative abundance (% of population) of immature *Cheilospirura spinosa* (●), *Cyrnea colini* (○) and *Tetrameres pattersoni* (▲) in adult bobwhites during 12 consecutive months.

by consistently maintaining high prevalences of infection (90-100%) throughout the period of study. The most plausible reason for the higher prevalences of infection with these species is their monoxenous life cycle (intermediate hosts not required). Although acquisition probably occurred to some extent each month, *H. bonasae* and *T. tenuis* were acquired in substantial numbers during different periods of the year. For *H. bonasae* this period was between March and July as evidenced by the higher proportions ( $P < 0.05$ ) of immature worms. The much higher intensity of *H. bonasae* larvae during certain collection periods (March and June) is similar to the peaks in acquisition of *H. gallinarum* larvae by free-ranging

galliform birds that have been attributed to consumption of earthworm transport hosts.<sup>16</sup> In contrast to *H. bonasae*, burdens of *T. tenuis* declined from March to September ( $P < 0.01$ ) and rose from September through February ( $P < 0.05$ ). Hon *et al.*<sup>10</sup> reported a similar trend of a winter peak for *T. tenuis* in wild turkeys (*Meleagris gallopavo*) in south central Florida. These opposing trends of *H. bonasae* and *T. tenuis* produced a dramatic shift in the composition of cecal worm populations with *H. bonasae* predominating during the summer and *T. tenuis* predominating during the winter (Fig. 3).

During a three-year, fall-to-spring study on helminthiasis in bobwhites from Kansas, Hanson and Robel<sup>9</sup>




FIGURE 3. The relative abundance of cecal nematodes, *Heterakis bonasae* (■) and *Trichostrongylus tenuis* (●), in adult bobwhites during 12 consecutive months.



measured seasonal trends by changes in the total prevalences of all cestodes and all nematodes. Their study revealed a peak prevalence of cestodes in the spring and peak prevalences of nematodes in November and February. The total prevalence of helminthiases in their study was only 23% compared to 100% every month in our study, and therefore prevalence was not a valuable basis for comparison of the two sets of data. In addition, the helminth fauna of bobwhites in the Kansas study differed markedly in species composition. Equivalent prevalences of helminthiases in juvenile and adult birds during fall and winter in the Kansas study, however, does suggest a similarity to our

study where helminth infections in juveniles approached levels in adults during the winter.

Lesions attributable to helminths were rare. Minor proliferation of the gizzard lining due to *C. spinosa* and *C. colini* occurred occasionally. Ulceration of the proventriculus at the point of attachment of *D. nasuta* also was noted infrequently. Two birds had greatly enlarged proventriculi due to heavy infections of *T. pattersoni*, and large numbers of *R. colinia* in two birds caused marked distention of the small intestine. Based on the severity and frequency of associated lesions helminths were not considered to be important mortality factors during the present study.

#### Acknowledgements

The authors wish to thank Dr. J.D. Webster, Department of Biology, Hanover College, Hanover, Indiana 47243 for lending specimens of *Raillietina colinia*.

#### LITERATURE CITED

1. BLAKENEY, W.C., Jr. and R.W. DIMMICK. 1971. Gizzard and intestinal helminths of bobwhite quail in Tennessee. *J. Wildl. Manage.* 35: 559-562.
2. BURNETTE, D.W. 1963. Endoparasites of the bobwhite quail, *Colinus virginianus* L. in southern Illinois. M.S. Thesis. Southern Illinois Univ., Carbondale. 19p.
3. BYRD, E.E. and F.E. KELLOGG. 1971. *Mediorhynchus bakeri*, a new acanthocephalan (Gigantorhynchidae) from the bobwhite, *Colinus virginianus virginianus* (L.). *J. Parasit.* 57: 137-142.
4. CRAM, E.B. 1931. Developmental stages of some nematodes of the Spiruroidea parasitic in poultry and game birds. USDA Tech. Bull. No. 227. 27p.
5. ———, M.F. JONES and E.A. ALLEN. 1931. Internal parasites and parasitic diseases of the bobwhite. Pp. 229-310. In: *The Bobwhite Quail: Its Habits, Preservation, and Increase*. H.L. Stoddard, Ed. Charles Scribner's Sons, New York. 559p.
6. DAVIDSON, W.R., G.L. DOSTER and M.B. MCGHEE. 1978. Failure of *Heterakis bonasae* to transmit *Histomonas meleagridis*. *Avian Dis.* 22: 627-632.
7. GOBLE, F.C. and H.L. KUTZ. 1945. The genus *Dispharynx* (Nematoda: Acuariidae) in galliform and passeriform birds. *J. Parasit.* 31: 323-331.
8. HANDLEY, C.O. 1931. Food of the young. Pp. 159-165. In: *The Bobwhite Quail: Its Habits, Preservation, and Increase*. H.L. Stoddard, Ed. Charles Scribner's Sons, New York. 559p.
9. HANSEN, M.F. and R.J. ROBEL. 1972. Seasonal changes and habitat influencing helminthiasis in bobwhite quail. Pp. 350-356. In: *Proceedings of*

- the First National Bobwhite Quail Symposium*. J.S. Morrison and J.C. Lewis, Eds. Oklahoma State University, Research Foundation, Stillwater. 390p.
10. HON, L.T., D.J. FORRESTER and L.E. WILLIAMS, Jr. 1978. Helminth acquisition by wild turkeys (*Meleagris gallopavo osceola*) in Florida. Proc. Helm. Soc. Wash. 45: 211-218.
  11. KELLOGG, F.E. and J.P. CALPIN. 1971. A checklist of parasites and diseases reported from the bobwhite quail. Avian Dis. 15: 704-715.
  12. ——— and G.L. DOSTER. 1972. Diseases and parasites of the bobwhite. Pp. 233-267. In: *Proceedings of the First National Bobwhite Quail Symposium*. J.S. Morrison and J.C. Lewis, Eds. Oklahoma State University, Research Foundation, Stillwater. 390p.
  13. ———, ——— and L.L. WILLIAMSON. 1970. A bobwhite density greater than one bird per acre. J. Wildl. Manage. 34: 464-466.
  14. ——— and A.K. PRESTWOOD. 1968. Gastrointestinal helminths from wild and pen-raised bobwhites. J. Wildl. Manage. 32: 468-475.
  15. REID, W.M. 1962. *Chicken and turkey tapeworms*. Univ. Georgia Coll. Agric. 71p.
  16. ———. 1967. Etiology and dissemination of blackhead disease syndrome in turkeys and chickens. Exp. Parasit. 21: 249-275.
  17. SCHMIDT, G.D. and R.E. KUNTZ. 1977. Revision of *Mediorhynchus* Van Cleave 1916 (Acanthocephala) with a key to species. J. Parasit. 63: 500-507.
  18. VERNARD, C. 1933. Helminths and coccidia from the Ohio bobwhite. J. Parasit. 19: 205-208.
  19. WEBSTER, J.D. 1944. A new cestode from the bobwhite. Trans. Am. Microsc. Soc. 63: 44-45.
  20. ———. 1947. Helminths from the bob-white in Texas, with descriptions of two new cestodes. Trans. Am. Microsc. Soc. 66: 339-343.
  21. ——— and C.J. ADDIS. 1945. Helminths from the bobwhite quail in Texas. J. Parasit. 31: 286-287.

*Received for publication 9 March 1977*

---