



## **GASTROINTESTINAL PARASITES OF THE EASTERN COTTONTAIL (*Sylvilagus floridanus*) IN CENTRAL PENNSYLVANIA 1**

Authors: WIGGINS, J.P., COSGROVE, M., and ROTHENBACHER, H.

Source: Journal of Wildlife Diseases, 16(4) : 541-544

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-16.4.541>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## GASTROINTESTINAL PARASITES OF THE EASTERN COTTONTAIL (*Sylvilagus floridanus*) IN CENTRAL PENNSYLVANIA <sup>□</sup>

J.P. WIGGINS, M. COSGROVE and H. ROTHENBACHER, Department of Veterinary Science, The Pennsylvania State University, University Park, Pennsylvania 16802, USA.

**Abstract:** During a 3 year period, 186 eastern cottontails (*Sylvilagus floridanus*) were trapped from two areas and examined for helminth and protozoan parasites. Fecal samples from 139 were evaluated for coccidia and helminth ova. Nine species of coccidia were identified: *Eimera audubonii*, *E. azul*, *E. environ*, *E. honessi*, *E. maior*, *E. minima*, *E. neoirresidua*, *E. neoleporis*, and *E. sylvilagi*. Ova from 5 helminth species were found: *Cittotaenia* sp., *Hasstilesia tricolor*, *Passalurus* sp., a trichostrongyle-type nematode species, and *Trichuris* sp. Five helminths were recovered from stomachs and small intestines: *Cittotaenia* sp. *H. tricolor*, *Obeliscoides cuniculi*, *Passalurus ambiguus*, and *Trichostrongylus calcaratus*.

### INTRODUCTION

Several detailed reports of internal parasites of *Sylvilagus* spp. and other leporids have been published in recent years.<sup>1,2,4,6,7,8,11,18</sup> While these reports indicated that the parasites of rabbit populations are similar in different areas, with coccidia, cestodes, gastrointestinal nematodes, and trematodes being present in almost all areas, the particular genera and species do have regional variations. This study was undertaken to identify the gastrointestinal parasites of the eastern cottontail in Pennsylvania, their characteristic dimensions, and prevalence.

### MATERIALS AND METHODS

Rabbits were trapped from two areas: a 4.1 ha plot on The Pennsylvania State University Wastewater Renovation Facility on State Game Lands 176 (SGL 176) and, approximately 8 km from this area, a 162 ha area at the State Correctional Institute at Rockview (Rockview), Centre County, Pennsylvania. Unbaited box traps were set at a density of 25

traps/ha with a grid interval of 20 m in early September, 1975-1977; and in March, 1976-1978. Vegetative cover at the SGL 176 area consisted of old field, cultivated field with grain and forage crops, multiflora rose hedge, and scattered pine and hardwood trees. Vegetation on the Rockview area was typified by orchards and old fields.

A total of 186 cottontails was trapped and then euthanized in a CO<sub>2</sub> chamber. At necropsy, animals were examined for lesions and parasites. Fecal samples were analyzed for coccidia and helminth ova, contents of stomachs and small intestines were examined for gastrointestinal parasites. A modification of the McMaster technique<sup>17</sup> was used for egg and oocyst counts. Oocysts were sporulated in 2.5% potassium dichromate for one week and then examined qualitatively.

Stomachs and small intestines were examined with the aid of a dissecting microscope for all animals except those trapped in September, 1975 and March, 1978. Helminths recovered from the stomachs and all intestinal nematodes

<sup>□</sup> Approved by the Director, The Pennsylvania Agricultural Experiment Station as paper 5643, Journal Series.

TABLE 1. Species of *Eimeria* recovered, their prevalence, and oocyst dimensions.

	<i>E. audubonii</i>	<i>E. azul</i>	<i>E. environ</i>	<i>E. honessi</i>	<i>E. maior</i>	<i>E. minima</i>	<i>E. neoirresidua</i>	<i>E. neoleporis</i>	<i>E. Sylvilagi</i>
Oocyst prevalence (%)	14.5	10.1	73.4	17.3	7.2	2.2	33.1	24.5	9.4
Oocyst length									
Mean	20.9	21.9	25.8	27.1	42.6	13.0	25.2	38.1	29.6
Range	17.5-24.5	18.0-27.0	19.0-35.1	21.0-35.0	27.0-48.0	12.0-14.0	20.0-35.0	28.0-46.0	24.0-37.0
Oocyst width									
Mean	16.1	16.6	17.4	17.6	24.8	12.0	16.8	18.4	18.3
Range	12.0-21.0	14.5-19.0	11.0-24.5	14.0-25.5	17.5-30.0	11.0-12.5	12.5-23.0	11.7-26.5	16.0-26.5
Oocyst residuum <sup>a</sup>	-	-	-	3.7	4.6	-	-	1.6	±b
Microspyle width	-	-	4.9	4.8	6.2	-	4.7	5.5	5.4
Sporocyst length									
Mean	11.0	11.9	13.1	13.4	18.1	6.1	12.6	15.7	15.1
Range	10.5-13.0	7.8-15.0	10.0-17.5	11.5-17.0	14.0-26.0	4.0-11.0	10.0-17.0	13.5-20.0	12.0-19.5
Sporocyst width									
Mean	5.7	5.6	6.4	6.4	8.0	4.0	6.4	7.2	7.0
Range	5.5-7.0	3.3-6.5	4.5-10.0	4.0-8.5	5.5-11.0	3.0-5.0	5.0-7.5	6.0-10.0	5.5-7.5
Sporocyst residuum <sup>a</sup>	-	2.7	-	-	4.2	+	2.8	5.1	4.3
Refractile globule <sup>a</sup>	3.5	3.8	4.0	4.5	5.5	+	4.0	5.0	5.0
Polar granule	-	-	-	-	-	+	-	-	-

- = Absent; + = Present.

<sup>a</sup>The means represent the diameter of the residua or refractile globule.<sup>b</sup>Mean residuum size is not presented since not all oocysts had a residuum.

were fixed in formalin for at least 24 h, then mounted with CMCP<sup>□</sup> mounting media for microscopic examination and species identification. Since this work was part of a larger project, stomach contents were removed and the intestinal sections were received after use in bacteriology and were not sufficiently complete to quantitatively evaluate the total number of helminths present. Means and ranges presented are in micrometers.

## RESULTS

Nine species of coccidia (Table 1), three species of nematode ova, one species of cestode ova, and one species of trematode ova were found (Table 2). Five helminths were recovered and identified from stomachs and small intestines: *Cittotaenia* sp., *Hasstilesia tricolor*, *Obeliscoides cuniculi*, *Passalurus ambiguus*, and *Trichostrongylus calcaratus*.

## DISCUSSION

Mean dimensions for all *Eimeria* spp., and ova of *Cittotaenia* sp., *H. tricolor*, and *Trichuris* sp. were similar to earlier studies.<sup>6,10,11,13,15,18</sup> Mean dimensions of

*Passalurus* sp. ova were slightly greater (108.7 × 50.6) than those reported previously (95.0 - 103.0 × 43.0).<sup>10</sup> Since *P. ambiguus* was consistently identified as the only oxyurid parasite present this suggests that the *Passalurus* sp. ova were from this one species. The mean dimensions of the trichostrongyle-type nematode ova were within the range for *Obeliscoides cuniculi* and *Trichostrongylus* spp. ova;<sup>10</sup> therefore, the ova could not be differentiated.

The prevalence of some helminth ova and coccidia recovered was rather typical of recent surveys.<sup>1,3,8,11,12,18</sup> However, low prevalence was noted for several *Eimeria* spp. and ova of *Cittotaenia* sp., *H. tricolor*, and *Passalurus* sp.<sup>1,3,6,8,12,16</sup> Since most tapeworm ova frequently are passed in proglottids, fecal egg flotation procedures may not reflect true prevalence of a tapeworm infection unless a gravid segment is completely crushed.<sup>5</sup> Thus, absence of tapeworm eggs does not always indicate lack of infection. A low prevalence for *H. tricolor* ova was probably a result of the quantification method used. A more accurate count of trematode ova requires the analysis of a fecal plug formed by centrifugation.<sup>14</sup>

TABLE 2. Species of helminth ova recovered, their prevalence, and ova dimensions.

Ova type	Ova prevalence (%)	Ova length		Ova width	
		Mean	Range	Mean	Range
<i>Cittotaenia</i> sp.	27.0	69.1	50.0-90.0	63.8	40.0-83.2
<i>Hasstilesia tricolor</i>	14.6	22.0	18.2-26.0	14.0	13.0-19.5
<i>Passalurus</i> sp.	11.7	108.7	98.0-123.0	50.6	34.2-74.1
Trichostrongyle-type	46.0	74.7	36.4-85.0	44.4	16.9-65.0
<i>Trichuris</i> sp.	16.1	63.9	63.8-75.0	29.4	23.1-39.6

## Acknowledgements

The authors would like to thank L. Frankhouser for technical assistance. This work was supported in part by grant matching funding by the U.S. Department of Interior,

<sup>□</sup> Turtox, Chicago, Illinois 60620, USA.

Office of Water Research and Technology and by the Institute for Research on Land and Water Resources, The Pennsylvania State University.

#### LITERATURE CITED

1. BOOKOUT, T.A. 1971. Helminth parasites in snowshoe hares from northern Michigan. *J. Wildl. Dis.* 7: 246-248.
2. BRITTAIN, P.C. and D.R. VOTH. 1975. Parasites of the black-tailed jackrabbit in northcentral Colorado. *J. Wildl. Dis.* 11: 269-271.
3. CARVALHO, J.C.M. 1942. *Eimeria neoleporis* n. sp., occurring naturally in the cottontail and transmissible to the tame rabbit. *Iowa St. Coll. J. Sci.* 16: 409-410.
4. ———. 1943. The coccidia of the wild rabbits of Iowa. I. Taxonomy and host specificity. *Iowa St. Coll. J. Sci.* 18: 103-133.
5. CHANDLER, A.C. 1949. *Introduction to Parasitology*, 9th Ed., Wiley and Sons, Inc., New York, 799 pp.
6. DUSZYNSKI, D.W. and W.C. MARQUARDT. 1969. *Eimeria* (Protozoa: Eimeridae) of the cottontail rabbit, *Sylvilagus audubonii*, in northeastern Colorado with descriptions of three new species. *J. Protozool.* 16: 128-137.
7. FLYNN, R.J. 1973. *Parasites of Laboratory Animals*, 5th Ed., The Iowa State University Press, Ames, 884 pp.
8. HOLLOWAY, H.L. 1966. Helminths of rabbits and opossums at Mountain Lake, Virginia. *J. Wildl. Dis.* 2: 38-39.
9. JACOBSON, H.A., R.L. KIRKPATRICK and B.S. MCGINNES. 1978. Disease and physiologic characteristics of two cottontail populations in Virginia. *Wildl. Monog. No. 60*, 53 pp.
10. LEVINE, N.D. 1968. *Nematode Parasites of Domestic Animals and Man*. Burgess Publishing Co., Minn., 600 pp.
11. ——— and V. IVENS. 1972. Coccidia of the Leporidae. *J. Protozool.* 19: 572-581.
12. NOVLESKY, M.A. and W.G. DYER. 1970. Helminths of the eastern cottontail rabbit, *Sylvilagus floridanus*, from North Dakota. *Am. Midl. Nat.* 84: 267-269.
13. ROWAN, W.B. 1955. The life cycle and epizootiology of the rabbit trematode, *Hasstilesia tricolor* (Stiles and Hassall, 1894) Hall, 1916 (Trematoda: Brachylaemidae). *Trans. Am. Microsc. Soc.* 74: 1-21.
14. SLOSS, M.W. and R.L. KEMP. 1978. *Veterinary Clinical Parasitology*, 5th Ed., Iowa St. Univ. Press, Ames, 274 pp.
15. SOULSBY, E.J.L. 1968. *Helminths, Arthropods, and Protozoa of Domesticated Animals*, 6th Ed., The Williams and Wilkins Co., Baltimore, 824 pp.
16. STRINGER, R. 1967. Parasites from rabbits from North Carolina. *J. Wildl. Dis.* 3: 93.
17. WHITLOCK, H.V. 1948. Some modifications of the McMaster helminth egg-counting technique and apparatus. *Austral. Counc. Sci. Ind. Res. J.* 21: 177-180.
18. WIGGINS, J.P. and H. ROTHENBACHER. 1979. *Eimeria azul* sp. n. from the Eastern Cottontail in central Pennsylvania. *J. Parasit.* 65: 393-394.

Received for publication 26 June 1979