



## Tuberculosis in a Ruffed Grouse

Author: SNOEYENBOS, G. H.

Source: Bulletin of the Wildlife Disease Association, 2(1) : 9

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-2.1.9>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Tuberculosis in a Ruffed Grouse

G. H. SNOEYENBOS

*Department of Veterinary & Animal Sciences  
University of Massachusetts, Amherst, Mass.*

Received for publication 24 August 1965

During November of 1958, a partially dressed ruffed grouse (*Bonasa umbelus umbelus*) which had been shot in Massachusetts was submitted for examination because it had grossly abnormal viscera. The cadaver was emaciated which was consistent with the hunter's report of a weak flight. Examination revealed an enlarged liver and spleen with numerous small yellowish-grey caseous nodules in these organs as well as in the lungs and in many areas of the peritoneum. Large numbers of slender acid-fast bacilli were found in smears made from a number of nodules. Lesions of caseation necrosis with peripheral giant cells and encapsulation were observed in sections stained with hematoxylin and eosin.

The presence of classical lesions and the demonstration of acid-fast bacilli was considered justification for a diagnosis of tuberculosis. Unfortunately no attempt was made to isolate and classify the organism.

Practically all species of birds are considered to be susceptible to

tuberculosis, and in almost all instances except in the parrot, the infection is due to the avian type of tubercle bacillus. The relatively few recorded instances of tuberculosis in free-living wild birds are thought to be a result of exposure to material derived from infected domestic chickens. Feldman (Feldman, W. H.: Avian tuberculosis infections. Baltimore, The Williams and Wilkins Company, 1938, pp. 234-242) cited reports of the disease in the free-living crow (*Corvus brachyrhynchos*), raven (*Corvus corax*), barn owl (*Strix pratincola*), cowbird (*Molothrus a. ater*), eastern sparrow hawk (*Falco s. sparverius*), in an unidentified owl and in a great variety of captive wild birds.

The ruffed grouse referred to in this report was killed in a woodlot near several homes which may have maintained a few free-ranging "backyard" chickens. Although avian tuberculosis is rare in commercial poultry in Massachusetts, it is occasionally found, and may be common in "backyard" flocks.