

IXODES TEXANUS IN EASTERN CANADA

Author: WEBSTER, W. A.

Source: Bulletin of the Wildlife Disease Association, 2(3) : 78-79

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-2.3.78.a>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

**NEW HOST AND DISTRIBUTION
RECORDS OF ACANTHOCEPHALA
FROM NORTH AMERICAN BIRDS**

During the past few years the authors have surveyed the helminth parasites of birds in several localities of the United States. This paper reports some host and

distribution records of *Acanthocephala* which appear to be new. New host and state records are indicated by an asterisk in the following table. The worms are retained in the authors' collections.

G. D. SCHMIDT
K. A. NEILAND

TABLE 1. Some records of *Acanthocephala* from birds

Parasite	Host and State
Polymorphidae	
<i>Polymorphus trochus</i>	<i>Fulica americana</i> Montana*
Van Cleave, 1945	
<i>Polymorphus</i> sp.	<i>Ixoreus naevius</i> * Oregon
<i>Arhythmorhynchus longicollis</i> (Villot, 1875)	<i>Larus delawarensis</i> * Oregon*
Plagiorhynchidae	
<i>Plagiorhynchus formosus</i>	<i>Ixoreus naevius</i> * Oregon*, Washington
Van Cleave 1918	<i>Colaptes cafer</i> Washington
<i>Plagiorhynchus paulus</i>	<i>Passerella iliaca</i> Oregon*
Van Cleave and Williams, 1951	<i>Ixoreus naevius</i> * Washington*
Centrorhynchidae	
<i>Sphaerovirostris lancea</i>	<i>Ixoreus naevius</i> Washington*
Westrumb, 1821) Golvan, 1960	<i>Perisoreus obscurus</i> * Washington*
Gigantorhynchidae	
<i>Mediorhynchus grandis</i>	<i>Quiscalus quiscula</i> <i>Sternella neglecta</i> * Colorado*
Van Cleave, 1916	
<i>Mediorhynchus papillosus</i>	<i>Turdus migratorius</i> * Colorado*
Van Cleave, 1916	<i>Junco caniceps</i> * Colorado*
	<i>Penthestes gambeli</i> * Colorado*
<i>Mediorhynchus</i> sp.	<i>Sciurus noveboracensis</i> * New York

Dept. of Zoology
Colorado State College
Greeley, Colo.
Alaska Dept. Fish and Game
Fairbanks, Alaska, 24 February, 1966

IXODES TEXANUS IN EASTERN CANADA

The presence of *Ixodes texanus* Banks has been reported only twice in Ontario, all other records indicate that this tick is restricted in Canada to the west coast area (Gregson, 1956, Can. Dept. Agric. Publ. 930). Reports from the United States, however, record the distribution of *I. texanus* in Maryland, Delaware, Virginia, North Carolina, South Carolina, Georgia and Florida in the eastern area (Clifford, Anastos, and Elbe, 1961, Entomol. Soc. Amer., Miscell. Publ. 2, 215-237).

The author has made several collections of *I. texanus* from Ontario, the records of which are listed below.

Ontario.

South March, 6.II.64, ex *Mephitis mephitis* 1L
Cookstown, 14.II.64, ex *Procyon lotor* 1 ♀
Strathroy, 27.IV.64, ex *Procyon lotor* 30 ♀ ♀, 1 ♂, 6LL
Fletcher, 21.V.64, ex *Procyon lotor* 1 ♀
Mactier, 5.VI.64, ex *Procyon lotor* 3 ♀ ♀

Moscow, 7.VII.64, ex *Procyon*
lotor 1 ♀
 Toronto, 13.VIII.64, ex *Pro-*
cyon lotor 3 ♀ ♀
 Barrie, 29.IX.64, ex *Procyon*
lotor 1 ♀
 Moffat, 6.IV.65, ex *Procyon*
lotor 1 ♀
 London, 30.IV.65, ex *Procyon*
lotor 3 ♀ ♀
 Weston, 13.V.65, ex *Procyon*
lotor 18 ♀ ♀, 7LL
 Winona, 20.V.65, ex *Procyon*
lotor 1 ♀
 Hyndford, 7.VII.65, ex *Vul-*
pes fulva 1 ♀
 Monkton, 21.VII.65, ex *Pro-*
cyon lotor 1 ♀

The distribution of hosts, from which the ticks were collected, would indicate that this species is fairly well established in Southern Ontario. It is the purpose of this note to confirm the presence of *I. texanus* in this area.

The author expresses his appreciation to Dr. J. D. Gregson, Research Station, Canada Department Agriculture, Kamloops, B. C. for confirming the identification of this tick. The assistance of Mr. J.E.H. Martin and Miss I. S. Creelman, Research Branch, Ottawa, Ont. with regards to records is gratefully acknowledged.

W. A. WEBSTER

Animal Disease Research Inst.
 Canada Dept. of Agriculture
 Hull, Quebec, 25 February, 1966

**PNEUMOSTRONGYLUS TENUIS FROM ELK
(CERVUS CANADENSIS) IN MINNESOTA**

A new host record for the parasitic nematode, *Pneumostrongylus tenuis* Dougherty 1945, has been established for Minnesota by the re-

covery of several first stage larvae from elk droppings collected on February 1, 1966, approximately nine miles north of Grygla, Marshall County, in northwestern Minnesota. The pellets were frozen at the time of collection and were kept frozen until February 5 when they were introduced into the Baerman apparatuses at the Winton Game and Fish Headquarters. Twenty apparatuses were in use.

After 48 hours five first-stage larvae were observed on 102 slides. After 72 hours a single larva was found on 94 slides and one larva was found after 120 hours on ten slides. Measurements of the larvae fell within the limits as described by R. C. Anderson in the Canadian Journal of Zoology 41:775-792. A photograph of one of these larvae is shown in Figure 1.




FIGURE 1. First-stage larva of *Pneumostrongylus tenuis* found in droppings of Minnesota Elk.

P. tenuis is common in the white-tailed deer (*Odocoileus virginianus borealis*) of this region, occurring in about 35 percent of the adult deer (Karns, unpublished data). It has also been reported