

BLOOD PARASITES OF BIRDS COLLECTED IN FOUR SUCCESSIVE YEARS IN PANAMA 1

Authors: HUFF, CLAY G., and WETMORE, ALEXANDER

Source: Bulletin of the Wildlife Disease Association, 3(4) : 178-181

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-3.4.178>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

BLOOD PARASITES OF BIRDS COLLECTED IN FOUR SUCCESSIVE YEARS IN PANAMA¹

Although knowledge of the blood parasites of nondomesticated birds is being advanced along special lines, the geographic and systematic frequency and distribution of these parasites are still not well known. This is particularly true of the parasites of birds in tropical regions. There is a great need for field studies, in both temperate and tropical regions, which would contribute to our knowledge about such parasites as *Plasmodium* and *Leucocytozoon* (which are transmitted by biting, flying insects) in reference to (1) the seasonal acquisition of parasites by the birds and (2) the interrelations between the parasite populations in the migratory and nonmigratory hosts.

This report is based on the study of blood smears made in Panama by one of us (AW) during the months of February and March of 1952, 1954, and 1955 and during May and June of 1953. The birds were collected for the Smithsonian Institution. Blood smears were made in the field at the time the birds were collected and were numbered to correspond to the specimens from which they were taken. Later in the laboratory the dried smears were fixed in methyl alcohol and stained with Giemsa's stain. Examination was made first by a high dry objective to detect the presence of the larger parasites such as microfilaria and then by an oil immersion objective to detect the smaller parasites.

Table 1 summarizes the distribution of the birds in which no blood parasites were found, according to their geograph-

ic distribution and their systematic classification. Table 2 lists the parasites found and their hosts. The provinces of Panama from which specimens were taken were Chiriquí, Veraguas, Coclé, Colon, and Panama; also Taboga and Taboguilla islands and the Canal Zone. Smears were examined from 183 individual birds. Twenty-eight of these contained one species of parasite (in 3 cases there were double infections). *Plasmodium* was found in 5 birds belonging to 5 families; *Haemoproteus* occurred in 11 birds belonging to 6 families; only one bird was found to be infected with *Leucocytozoon*; *Trypanosoma* occurred in 4 birds belonging to 3 families; and microfilaria were found in 11 birds belonging to 8 families.

No attempt is made here to determine the species of any of these parasites with the exception of some of those belonging to the genus *Plasmodium*. The stages of the latter found in *Eucometis penicillata* would appear to place it close to, or in the *hexamerium vaughani* group of *Plasmodium*, and the stages found in *Tangavius aeneus* probably are *P. cathemerium* or *P. relictum*. Precise determination of species of *Plasmodium* usually can be made only by isolation of the parasites in susceptible laboratory animals and a careful study of the biological as well as the morphological characteristics of the resulting infections. Specific determinations of the parasites belonging to *Haemoproteus*, *Leucocytozoon*, *Trypanosoma* and of microfilaria cannot be made, except in rare cases, upon the study of the very few parasites found in a blood smear.

One is inclined to expect that birds living in tropical areas might exhibit a higher incidence of infection of those parasites such as *Plasmodium* and *Leucocytozoon* which, in more temperate climates, have their transmission restricted to the warmer months of the year. The results from this small group of smears from Panama do not meet

¹ From Bureau of Medicine and Surgery, Navy Department, Research Task MR-005.09.0026. The opinions and assertions contained herein are those of the authors and are not to be construed as official or reflecting the views of the Navy Department or the Naval service at large.

TABLE 1. *Distribution of Birds Examined and Found Not To Have Blood Parasites*

Order and Family	Number of Specimens from								Number species	Number genera
	Code'	Chiriqui'	Veraguas	Panama'	Colon'	Canal Zone	Taboga Is.	Taboguilla Is.		
Tinamiformes										
Tinamidae	1							1	1	1
Podicipediformes										
Podicipedidae				2				2	2	2
Ciconiiformes										
Ardeidae	1							1	1	1
Falconiformes										
Cathartidae			1					1	1	1
Buteonidae	1		2					3	2	2
Falconidae	1							1	1	1
Galliformes										
Phasianidae			1					1	1	1
Columbiformes										
Columbidae	1	5	1					7	5	4
Psittaciformes										
Psittacidae		3						3	1	1
Cuculiformes										
Cuculidae		1						1	1	1
Apodiformes										
Trochilidae	3	9	3	1			1	17	14	11
Trogoniformes										
Trogonidae		1						1	1	1
Coraciiformes										
Alcedinidae			1					1	1	1
Piciformes										
Capitonidae	1	1						2	2	2
Ramphastidae		1						1	1	1
Picidae	1	6		1				8	6	5
Passeriformes										
Dendrocolaptidae	2	1	1					4	3	3
Furnariidae		1	1					2	2	2
Formicariidae	2	3	3					8	7	5
Cotingidae		4						4	3	3
Pipridae		2	2					4	3	3
Tyrannidae	5	13	7	2	1		2	31	23	19
Hirundinidae	3	1	3			1		8	4	4
Troglodytidae	1	1	2	4		1		9	5	3
Turdidae	1							1	1	1
Vireonidae		1						1	1	1
Coerebidae			2			1		3	2	2
Parulidae	1	1		1		1		4	4	3
Icteridae		2	1					4	4	4
Thraupidae	3	5	2					10	8	5
Fringillidae	5	3	2	1				11	8	8
TOTALS	33	65	35	12	1	4	3	2 155	119	102

TABLE 2. Blood Parasites of Birds of Panama¹

<i>Plasmodium</i>		
<i>Thryothorus modestus</i> , TROGLODYTIDAE (wrens) ♀ Sona', Veraguas	(449,100)	23 Jun '53
<i>Tyrannus melancholicus</i> , TYRANNIDAE (tyrant flycatchers) ♀ El Uracillo, Coclé ²	(445,237)	27 Feb '52 ¹
<i>Eucometis penicillata</i> , THRAUPIDAE (tanagers) ♂ Chilar, Colo'n	(445,528)	18 Mar '52 ² ³
<i>Tangavius aeneus</i> , ICTERIDAE (blackbirds) ♂ Sona', Veraguas	(449,234)	27 May '53 ⁴
<i>Myioborus torquatus</i> , PARULIDAE (wood warblers) ♂ Cerro Punta, Chiriquí	(455,959)	23 Mar '54
<i>Haemoproteus</i>		
<i>Nystalus radiatus</i> , BUCCONIDAE (puffbirds) ♀ El Uracillo, Coclé	(445,701)	25 Feb '52
<i>Gymnostinops montezuma</i> , ICTERIDAE (blackbirds) ♂ Taboga Is.	(445,479)	17 Mar '52 ²
<i>Muscivora tyrannus</i> , TYRANNIDAE (tyrant flycatchers) ♂ Sona', Veraguas	(448,979)	27 May '53
<i>Vireo flavoviridis</i> , VIREONIDAE (vireos) ♀ Sona', Veraguas	(449,152)	1 Jun '53
<i>Vireo flavoviridis</i> , VIREONIDAE (vireos) ♀ Sona', Veraguas	(449,155)	5 Jun '53 ²
<i>Vireo flavoviridis</i> , VIREONIDAE (vireos) ♂ Taboga Is.	(445,438)	17 III '52
<i>Vireo leucophrys chiriquensis</i> , VIREONIDAE (vireos) ♂ El Volca'n, Chiriquí	(455,877)	22 III '54
<i>Dendroica petechia equatorialis</i> , PARULIDAE (woodwarblers) ♂ Puerto Vidal, Veraguas	(449,215)	8 VI '53
<i>Dendroica petechia equatorialis</i> , PARULIDAE (woodwarblers) ♂ Puerto Mutis, Veraguas	(449,210)	11 VI '53
<i>Myioborus miniatus acceptus</i> , PARULIDAE (woodwarblers) ♂ El Volca'n, Chiriquí	(455,955)	3 III '54
<i>Falco s. sparverius</i> , FALCONIDAE (falcons) ♀ El Volca'n, Chiriquí	(455,527)	5 III '54
<i>Leucocytozoon</i>		
<i>Pharomachrus mocino costaricensis</i> , TROGONIDAE (trogon) ♀ El Volca'n, Chiriquí	(455,626)	5 III '54
<i>Trypanosoma</i>		
<i>Tyrannus melancholicus</i> , TYRANNIDAE (tyrant flycatchers) ♀ El Uracillo, Coclé	(455,237)	27 II '52
<i>Cyclarhis guianensis subflavescens</i> , CYCLARHIDAE (pepper shrikes) ♀ El Volca'n, Chiriquí	(455,866)	29 III '54
<i>Phloeocastes g. guatemalensis</i> , PICIDAE (woodpeckers) ♀ El Volca'n, Chiriquí	(458,607)	12 II '55
<i>Microfilaria</i>		
<i>Tanagra inornata</i> , THRAUPIDAE (tanagers) ♀ El Uracillo, Coclé	(445,547)	25 II '52
<i>Chlorophanes spiza</i> , COEREVIDAE (honey-creepers) ♀ El Uracillo, Coclé	(445,221)	II '52
<i>Cyanocorax affinis zeledoni</i> , CORVIDAE (crows, jays) ♂ Sona, Veraguas	(449,135)	21 V '53
<i>Eubucco bourcierii salvini</i> , CAPITONIDAE (barbets) ♂ El Volca'n, Chiriquí	(455,612)	11 III '54
<i>Turdus grayi casius</i> , TURDIDAE (thrushes) ♂ El Volca'n, Chiriquí	(455,858)	8 III '54
<i>Thryothorus rufalbus castanonotus</i> , TROGLODYTIDAE (wrens) ♂ Cerro Chame, Panama	(458,963)	27 Mar '55
<i>Gymnostinops montezuma</i> , ICTERIDAE (blackbirds) ♂ El Uracillo, Coclé	(445,479)	2 III '52
<i>Eucometis penicillata</i> , THRAUPIDAE (tanagers) ♂ Chilar, Colo'n	(445,528)	18 III '52

¹ also *Trypanosoma*² also microfilaria³ probably in *hexamerium-vaughani* group⁴ probably *cathemerium* or *relictum*

The numbers in parentheses are the U.S. National Museum catalog numbers.

that expectation. When compared with the smears made over a 10-year period from birds caught in the United States for banding purposes (Huff, 1939, J. Amer. Vet. Med. Assoc. 94: 615-620), the percentage of birds with blood infections is 15.2 for the smears from the birds of Panama, whereas it was 44.4 for the birds in the United States. It must, however, be noted that there are major differences between the two studies. First, collections were confined to the late winter and summer months in the Panamanian group while those in the United States were taken over a larger portion of the year. Second, the distributions among taxonomic groups were very different in the two studies. Although there were nearly seven times as many smears in the group from the birds in the United States as in those from Panama, the latter were taken from about four times as many species as the former. Likewise, in the higher taxonomic groups, there were 31 families and 13 orders of birds represented in the group from Panama as compared with 13 families and 6 orders represented in those from the United States.

Among the birds which were infected the percentages in the different genera of parasites were as follows:

	Panama birds	U.S. birds
<i>Haemoproteus</i>	39	80
<i>Plasmodium</i>	17	15
<i>Leucocytozoon</i>	3.5	2.7
<i>Trypanosoma</i>	10.7	2.7
<i>Microfilaria</i>	28.5	(not recorded)

In the present report all birds with infections of *Plasmodium* and microfilaria belonged to the order Passeriformes; the only *Leucocytozoon* was found in the order Trogoniformes; those with *Haemoproteus* were distributed in the orders Passeriformes (7 families), and Piciformes (1 family); and *Trypanosoma* were found in two families of the order Passeriformes and one family in the order Piciformes.

SUMMARY

During February and March of 1952, 1954, and 1955 and May and June, 1953, a total of 183 blood smears were made from birds collected for the Smithsonian Institution. Blood parasites were found in 28 of these; *Plasmodium* (5), *Haemoproteus* (11), *Leucocytozoon* (1), *Trypanosoma* (4), and microfilaria (11). The frequency for all blood parasites was 15.2 per cent as contrasted with 44.4 per cent in a previous survey of birds collected in the United States. *Haemoproteus* was less frequent and *Trypanosoma* more frequent in the birds from Panama than those in an earlier study from the United States.

CLAY G. HUFF
and ALEXANDER WETMORE
Naval Medical Research Institute,
Bethesda, Maryland
and the Smithsonian Institution,
Washington, D.C.
7 July, 1967

OBITUARY

Dr. Gordon M. Clark died June 20, 1967 from electrocution. He was working alone late at night in his new laboratory which was nearing completion. He was a research entomologist at the Rocky Mountain Laboratory of the U.S. Public Health Service in Hamilton, Montana.

Dr. Clark was a member of the Wildlife Disease Association since receiving his Ph.D. from the University of Maryland in 1956. Following graduation he was a member of the staff of the disease section of the Patuxent Wildlife Research Center until 1962 when he transferred to Montana. His main research interest was on arthropod vectors of disease and, at the untimely catastrophe of his death when only 37 years old, was a recognized authority on nasal mites of birds.

The Wildlife Disease Association mourns his loss and extends sympathies to his wife and son.