

## **Immobilization of Rocky Mountain Elk with Telazol® and Xylazine Hydrochloride, and Antagonism by Yohimbine Hydrochloride**

Authors: Millspaugh, Joshua J., Brundige, Gary C., Jenks, Jonathan A., Tyner, C. Lee, and Hustead, David R.

Source: Journal of Wildlife Diseases, 31(2) : 259-262

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-31.2.259>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Immobilization of Rocky Mountain Elk with Telazol® and Xylazine Hydrochloride, and Antagonism by Yohimbine Hydrochloride

Joshua J. Millspaugh,<sup>1,5</sup> Gary C. Brundige,<sup>2</sup> Jonathan A. Jenks,<sup>1</sup> C. Lee Tyner,<sup>3</sup> and David R. Hu-  
stead,<sup>4</sup> <sup>1</sup> Department of Wildlife and Fisheries Sciences, Box 2140B, South Dakota State University, Brookings, South  
Dakota 57007, USA; <sup>2</sup> Custer State Park, HC 83 Box 70, Custer, South Dakota 57730, USA; <sup>3</sup> Veterinary Technology  
Program, UPO 995, Morehead State University, Morehead, Kentucky 40351, USA; <sup>4</sup> Fort Dodge Laboratories, P.O.  
Box 518, Fort Dodge, Iowa 50501, USA; <sup>5</sup> Present address: College of Forest Resources, University of Washington,  
AR-10, Seattle, Washington 98195, USA

**ABSTRACT:** Ten trapped Rocky Mountain elk (*Cervus elaphus nelsoni*) were successfully im-  
mobilized with a combination of 500 mg Tel-  
azol® and 60 mg xylazine hydrochloride (HCl)  
from 9 July to 25 August 1993 in Custer State  
Park, South Dakota (USA). Mean (SD) dosages  
of 2.5 (0.6) mg/kg Telazol® and 0.3 (0.1) mg/  
kg xylazine HCl, respectively, were adminis-  
tered, resulting in a mean (SD) induction time  
of 4.6 (0.8) min. Induction time varied with  
weight and dosage. Respiratory rate (breaths/  
min) increased following injection of Telazol®  
and xylazine HCl and remained elevated or con-  
tinued to increase through 10 min post-injection  
and then declined. There were no mortalities in  
this study. Forty mg of yohimbine HCl was used  
as an antagonist in eight elk, resulting in a mean  
(SD) recovery time of 14.0 (9.9) min when ad-  
ministered intravenously ( $n = 6$ ), and 124.7 (9.5)  
min when given intramuscularly ( $n = 2$ ). Re-  
covery time varied with weight and dosage of  
yohimbine. Elk given 2.1 to 2.6 mg/kg Telazol®  
and 0.1 to 0.3 mg/kg xylazine HCl responded  
to yohimbine HCl when administered intrave-  
nously.

**Key words:** *Cervus elaphus nelsoni*, elk,  
immobilization, Telazol®, xylazine hydrochlo-  
ride, yohimbine hydrochloride.

Elk (*Cervus elaphus*) have been im-  
mobilized with succinylcholine chloride  
(Flook et al., 1962), etorphine hydrochlo-  
ride (HCl) (Magonigle et al., 1977), car-  
fentanil (Meulman et al., 1984), xylazine  
HCl (Thurmon et al., 1972), and a com-  
bination of xylazine HCl and ketamine HCl  
(Golightly and Hofstra, 1989). However,  
numerous complications resulted from the  
use of these chemical agents or their com-  
binations (Golightly and Hofstra, 1989).

Telazol® is a 1:1 mixture by weight of  
tiletamine HCl and zolazepam HCl. Tile-  
tamine HCl has similar pharmacologic  
activity to ketamine HCl (Taylor et al.,

1989), but is more potent (Short et al.,  
1989). Zolazepam HCl is a benzodiazepine  
tranquilizer and similar in pharmacologi-  
cal activity to diazepam. Xylazine HCl  
has been used successfully with ketamine  
HCl to immobilize white-tailed deer (*Odo-  
coileus virginianus*) (Mech et al., 1985)  
and moose (*Alces alces*) (Garner and Ad-  
dison, 1994).

Our objective was to determine the ef-  
ficacy and standard dosages of Telazol®  
and xylazine HCl to immobilize trapped  
elk and to determine the feasibility of re-  
versing the action of Telazol® and xylazine  
HCl with yohimbine HCl.

We trapped Rocky Mountain elk (*C.  
elaphus nelsoni*) in modified Clover traps  
in Custer State Park, South Dakota (USA)  
(43°45'N, 103°22'W) from 1 July to 30 Au-  
gust 1993 (Millspaugh et al., 1994). Cap-  
tured elk were given an intramuscular (IM)  
injection of 500 mg Telazol® (Fort Dodge  
Laboratories, Inc., Fort Dodge, Iowa, USA)  
reconstituted in 3 ml (20 mg/kg) xylazine  
HCl (TranquiVed®, Vedco Inc., St. Joseph,  
Missouri, USA) into either the gluteus max-  
imus, gluteus medius, or vastus lateralis  
with a hand syringe.

Following sedation, an ophthalmic oint-  
ment (Vetropolycin®, Pitman-Moore, Inc.,  
Mundelein, Illinois, USA) was applied to  
the elk's eyes to prevent corneal drying. A  
blindfold was used to help calm the ani-  
mal. Each elk was fitted with a mortality  
sensing radio transmitter (Lotek Engi-  
neering, Inc., Newmarket, Ontario, Can-  
ada) and ear-tagged. Ages of elk were es-  
timated by tooth wear and replacement  
(Quimby and Gaab, 1957). Ten ml of Dur-

apen® (Vedco, Inc., Overland Park, Kansas, USA), a combination antibiotic containing penicillin benzathine and penicillin procaine and 5 ml of selenium and vitamin E (E-SE®, Schering-Plough Animal Health Corp., Kenilworth, New Jersey, USA) were administered IM. Respiration rate (breaths/min) was recorded for 2 min at 10 min intervals when feasible.

We attempted to reverse the Telazol® and xylazine HCl combination with 40 mg yohimbine HCl (5 mg/ml) (Antagonil®, Lloyd Laboratories, Shenandoah, Iowa) by injection into either the jugular vein or IM about 20 min post-immobilization. All elk were monitored visually until their departure from the capture site, and were monitored three to five times per week until 1 January 1994. Weight estimations for immobilized elk were derived using a regression (SAS Institute Inc., 1990) based on chest girth and weight of hunter killed elk in Custer State Park (J. Millspaugh, unpubl.).

Induction time was defined as the period from injection to the point when elk were sedated sufficiently to be handled. Reversal time was defined as the period from yohimbine HCl administration until the elk was walking. Regression analysis (SAS Institute Inc., 1990) was used to determine the relationship between recovery time and dose, induction time and dose, recovery time and weight, and induction time and weight.

Ten 1- to 7-yr-old elk (five cows and five bulls) were successfully immobilized with 500 mg Telazol® and 60 mg xylazine HCl over a wide range of estimated weights (141 to 239 kg). One elk received an additional dose of 100 mg Telazol® and 40 mg xylazine HCl and was not included in the analyses. No mortality was observed.

Mean (SD) dose to achieve an effective level of anesthesia was 2.5 (0.6) mg/kg Telazol® (range 2.1 to 3.8,  $n = 10$ ) and 0.35 (0.01) mg/kg xylazine HCl (range = 0.3 to 0.5,  $n = 10$ ). Mean (SD) induction time was 4.6 (0.8) min (range = 3.5 to 6.2,  $n = 10$ ). Induction time was linearly re-

lated to weight ( $y = 0.02x - 0.37$ ,  $y$  = induction time in min and  $x$  = weight in kg,  $r^2 = 0.65$ ) and dosage ( $y = 7.59 - 1.19x$ ,  $y$  = induction time in min and  $x$  = dose in mg/kg,  $r^2 = 0.58$ ). Mean (SD) respiration rate was 53.0 (15.9) breaths/min immediately post-injection (range = 42 to 60,  $n = 4$ ) and 44.6 (14.5) breaths/min at 30 min post-injection (range = 36 to 64,  $n = 5$ ). Typically, at 2 to 3 min post-injection, elk began to lose coordination of leg and neck muscles and by 5 min were positioned sternal. Depth of anesthesia was consistent among all immobilized elk and no significant adverse side effects were noted.

Six of 10 elk received 40 mg yohimbine HCl intravenously (IV), two received 4 mg IM and two elk did not receive yohimbine HCl. Mean (SD) recovery time when yohimbine HCl was administered IV was 14.0 (9.9) min (range = 0.5 to 28.5 min,  $n = 6$ ). Recovery time was 115.2 min and 134.1 min for the two elk that received yohimbine HCl IM. Recovery time was related to weight ( $y = 56.35 - 0.23x$ ,  $y$  = recovery time in min and  $x$  = weight in kg,  $r^2 = 0.82$ ) and dosage ( $y = 14.23x - 26.32$ ,  $y$  = recovery time in min and  $x$  = dosage in mg/kg,  $r^2 = 0.81$ ). Recovery time was 145 min and 32 min for the two elk that did not receive yohimbine HCl.

Telazol® (500 mg) and xylazine HCl (60 mg) doses for our trapped elk resulted in rapid induction times and provided satisfactory anesthesia. Mean induction time in our study was less than for elk immobilized with xylazine HCl and ketamine HCl (Golightly and Hofstra, 1989), xylazine HCl alone (McCorquodale et al., 1988), succinylcholine chloride (Amstrup et al., 1982), etorphine (Magonigle et al., 1977), or carfentanil (Meulman et al., 1984).

Both xylazine HCl (Jessup and Clark, 1986) and Telazol® (Haigh et al., 1985) have wide safety margins. In our study, one elk required an additional dose to attain effective sedation possibly due to its large size and greater level of excitement.

This elk received a total dose of 600 mg Telazol® and 100 mg xylazine HCl and no adverse side effects were noted.

The use of Telazol® and xylazine HCl minimizes risks of accidental self-injection associated with the use of etorphine (Parker and Haigh, 1982) and carfentanil (Meulman et al., 1984). Neither xylazine HCl (Carruthers et al., 1979) or Telazol® (Gibbeau and Paquet, 1991) is lethal to humans in small doses.

A disadvantage of Telazol® is the lack of a known antagonist and potential for lengthy recovery (Haigh et al., 1985). Based on our analyses, elk given low doses of Telazol® (2.1 to 2.5 mg/kg) and xylazine HCl (0.3 mg/kg) responded to yohimbine HCl administered IV. Yohimbine HCl may be efficacious when administered IM, but only at much higher dosages. Following administration of yohimbine HCl, the pattern of recovery was predictable. First signs of recovery in our study were ear twitching, elevation of the head followed by sternal recumbency and standing and walking. Coordination was poor for several minutes.

The combination of Telazol® and xylazine HCl reported herein provided an effective level of anesthesia for trapped elk. Furthermore, we recommend the use of about 2.4 mg/kg Telazol® with 0.3 to 0.4 mg/kg of xylazine HCl to immobilize trapped elk. Yohimbine HCl was most effective as an antagonist when administered IV at low dose Telazol® and xylazine, resulting in shorter recovery, making Telazol® combined with xylazine HCl more efficacious in field studies. Advantages of Telazol® and xylazine HCl included ease of preparation, rapid induction, small dosage volume, safety to handler, minimal side effects, and predictable handling time and behavior.

Special thanks are extended to P. Whittenburg and R. Baughman for advice and in obtaining equipment. D. Garner provided advice on immobilization procedures and reviewed the manuscript. We are grateful to G. Bear for loaning us Clo-

ver traps. J. Baldwin, D. Havens, B. Rainville, K. Steiner, D. Welch, and C. Willmott assisted with field operations. L. Flake, K. Higgins, B. Kernohan, and two anonymous reviewers provided helpful comments on this manuscript. Our study was funded by the Rocky Mountain Elk Foundation and Federal Aid to Wildlife Restoration Fund, Project W-75-R (Study No. 7559) administered through South Dakota Department of Game, Fish, and Parks.

#### LITERATURE CITED

- AMSTRUP, S. C., B. W. O'GARA, AND H. MUSGRAVE. 1982. Immobilizing elk with powdered succinylcholine chloride. *Wildlife Society Bulletin* 10: 333-340.
- CARRUTHERS, S. G., M. NELSON, H. R. WEXLER, AND C. R. STILLER. 1979. Xylazine hydrochloride (Rompun) overdose in man. *Clinical Toxicology* 15: 281-285.
- FLOOK, D. R., J. R. ROBERTSON, O. R. HERMANRUDE, AND H. K. BUECHNER. 1962. Succinylcholine chloride for immobilization of North American elk. *The Journal of Wildlife Management* 26: 334-336.
- GARNER, D. L., AND E. M. ADDISON. 1994. Postpartum immobilization of adult female moose using xylazine, ketamine, and yohimbine hydrochlorides. *Journal of Wildlife Diseases* 30: 123-125.
- GIBBEAU, M. L., AND P. C. PAQUET. 1991. Evaluation of Telazol® for immobilization of black bears. *Wildlife Society Bulletin* 19: 400-402.
- GOLIGHTLY, R. T., AND T. D. HOFSTRA. 1989. Immobilization of elk with ketamine-xylazine mix and rapid reversal with yohimbine hydrochloride. *Wildlife Society Bulletin* 17: 53-58.
- HAIGH, J. C., I. STIRLING, AND E. BROUGHTON. 1985. Immobilization of polar bears with a mixture of tiletamine hydrochloride and zolazepam hydrochloride. *Journal of Wildlife Diseases* 21: 43-47.
- JESSUP, D. A., AND W. E. CLARK. 1986. *Wildlife restraint handbook*. Wildlife Investigation Laboratory, California Department of Fish and Game, Rancho Cordova, California, 162 pp.
- MAGONIGLE, R. A., E. H. STAUBER, AND H. W. VAUGHN. 1977. The immobilization of wapiti with etorphine hydrochloride. *Journal of Wildlife Diseases* 13: 258-261.
- MCCORQUODALE, S. M., L. E. EBERHARDT, AND S. E. PETRON. 1988. Helicopter immobilization of elk in southcentral Washington. *Northwest Science* 62: 49-54.
- MECH, L. D., G. D. DELGIUDICE, P. D. KARNS, AND U. S. SEAL. 1985. Yohimbine hydrochloride as an antagonist to xylazine hydrochloride-ketam-

- ine hydrochloride immobilization of white-tailed deer. *Journal of Wildlife Diseases* 48: 405–410.
- MEULMAN, T., J. D. PORT, T. H. STANLY, K. F. WILLIARD, AND J. KIMBALL. 1984. Immobilization of elk and moose with carfentanil. *The Journal of Wildlife Management* 48: 258–262.
- MILLSPAUGH, J. J., G. C. BRUNDIGE, AND J. A. JENKS. 1994. Summer elk trapping in South Dakota. *The Prairie Naturalist* 26: 125–129.
- PARKER, J. B. R., AND J. C. HAIGH. 1982. Human exposure to immobilizing agents. In *Chemical immobilization of North American wildlife*, L. Nielsen, J. C. Haigh, and M. E. Fowler (eds.). Wisconsin Humane Society, Milwaukee, Wisconsin, pp. 119–136.
- QUIMBY, D. C., AND J. E. GAAB. 1957. Mandibular dentition as an age indicator in Rocky Mountain elk. *The Journal of Wildlife Management* 21: 435–451.
- SAS INSTITUTE INC. 1990. SAS procedures guide, version 6, 3rd ed. SAS Institute Inc., Cary, North Carolina, 705 pp.
- SHORT, C. E., C. H. TRACY, AND E. SANDERS. 1989. Investigating xylazine's utility when used with Telazol® in equine anesthesia. *Veterinary Medicine* 86: 228–233.
- TAYLOR, W. P., JR., H. V. REYNOLDS, III, AND W. B. BALLARD. 1989. Immobilization of grizzly bears with tiletamine hydrochloride and zolazepam hydrochloride. *The Journal of Wildlife Management* 53: 978–981.
- THURMON, J. C., D. R. NELSON, AND J. O. MOZIER. 1972. A preliminary report on the sedative effect of Bay VA 1470 in elk (*Cervus elaphus canadensis*). *Journal of Zoo and Wildlife Medicine* 3: 9–14.

*Received for publication 3 May 1994.*